

x 5347

MAGNETIC LOW


FLIGHT LINE SPACING 3/4 MILE FLIGHT ALTITUDE NOMINALLY 1000 FEET ABOVE GROUND REGIONAL MAGNETIC FIELD SW SHEET CORNER: 56,756 GAMMAS REGIONAL FIELD REMOVED. THE FIELD INCREASES APPROXIMATELY 4.7 GAMMAS/MILE, N 50° E APPROXIMATE FIELD INCLINATION: +77.0°



FAIRBANKS (D-3), ALASKA

STATE OF ALASKA DEPARTMENT OF NATURAL RESOURCES DIVISION OF GEOLOGICAL AND GEOPHYSICAL SURVEYS

Copies of this map may be obtained from the Division at: 3001 Porcupine Drive, Anchorage, Alaska, 99501, and Box 80007, College, Alaska, 99701.



The magnetic contours shown on this map represent the total anomalous magnetic field of the earth. Variations in this field are caused by the variable magnetic character of rock units crossed by the survey flights, and hence, can be used to estimate the apparent location of rocks rich in magnetic minerals. Such rock units may be either at the surface of the ground or buried beneath it. Anomalies show both positive and negative variations depending on the shape, attitude, and constituents of local rocks. Geophysical interestation will be helpful in determining boundaries or death of buriel of depending on the snape, arrivate, and constituting in tooth local scales, seephysical interpretation will be helpful in determining boundaries or depth of buriel of anomally-causing rock units. Some anomalies may be impossible to interpret without further geologic information. Basic profile data is retained at the Division of Geological and Geophysical Surveys and should be consulted for detailed analysis.

Base map from U.S.G.S. I: 63 360 Topographic map series. Flown and compiled in 1973 by: LOCKWOOD, KESSLER & BARTLETT, INC., 2476 Huntington Drive, San Marino, California, 91108.