

Public-data File 83-55

LIVENGOOD STUDIES RESUMED: A SUMMARY OF RECENT MINERAL-RESOURCE
INVESTIGATIONS BY THE
ALASKA DIVISION OF GEOLOGICAL AND GEOPHYSICAL SURVEYS

By

M.D. Albanese, T.E. Smith, M.S. Robinson, and T.K. Bundtzen

Alaska Division of
Geological and Geophysical Surveys

November 1985

THIS REPORT HAS NOT BEEN REVIEWED FOR
TECHNICAL CONTENT (EXCEPT AS NOTED IN
TEXT) OR FOR CONFORMITY TO THE
EDITORIAL STANDARDS OF DGGS.

794 University Avenue, Basement
Fairbanks, Alaska 99701



Livengood Studies Resumed: A Summary of Recent
Alaska Division of Geological and Geophysical Surveys
Mineral Resource Investigations

By

M.D. Albanese, T.E. Smith, M.S. Robinson, and T.K. Bundtzen

INTRODUCTION

As part of a program to reexamine mineral resource potential of Interior Alaska Mining Districts, Alaska Division of Geological and Geophysical (DGGS) personnel mapped and geochemically sampled 1,000 square miles including the Livengood B-3, B-4, C-3, and C-4 Quadrangles during May-June of 1982. Maps have been produced to show bedrock geology at 1:40,000 scale (Robinson, 1983; Smith, 1983; Bundtzen, 1983; and Albanese, 1983), analyses of over 2,000 stream sediment pan-concentrate, and rock samples (Albanese, 1983), surficial geology at 1:63,360 scale (Waythomas, 1984), and detailed geology map (and report) of the Old Smoky gold-antimony prospect at 1:120 scale (Allegro, 1984). The previously published Livengood geologic map at 1:250,000 scale by Chapman, Weber, and Foster (1971) greatly assisted this investigation. Highlights of the DGGS Livengood study are included in this report.

Geologic Units

The bedrock geology of the study area is generalized in figure 1. The rock units generally occur in NE trending belts and range from Late Precambrian(?) and Lower Cambrian to Cretaceous. Bedrock units are complexly deformed and outcrop exposure is poor. Contacts, when exposed, are often fault bounded and stratigraphic relationships are difficult to determine.

The oldest rocks comprise an interlayered sequence of maroon and green argillite, black shale and limestone, vitreous quartzite and several types of bimodal quartz sandstone (informally known as the 'Grit' unit of Chapman and others, 1971). This unit contains Oldhamia, a trace fossil of probable Late Precambrian(?) to Early Cambrian age (Churkin and Brabb, 1965). These rocks are bordered on the north and perhaps overlain by a northeast trending belt of orthoquartzite, chert, maroon or green shale to phyllite, amygdaloidal greenstone, diorite, and gabbro. These younger rocks may represent a sediment-rich facies of the Fossil Creek Volcanics of Middle Ordovician age (Church and Dufree, 1961) which occurs northeast of the map area in the White Mountains (Bundtzen, 1983).

The 'Grit unit' appears to be in thrust contact with an extensive Chert terrane composed of variegated chert, gray and tan chert, chert conglomerate, quartzite, dolomitic limestone, and minor felsic tuff and greenstone. This unit includes the Livengood Dome Chert of Chapman and others (1980). The age of the chert terrane is believed to be Ordovician to Silurian based primarily on stratigraphic and fossil evidence including graptolites (Chapman and others, 1980), conodonts, brachiopods, and corals (Florence Weber, U.S. Geological Survey, personal commun., 1982, and Robinson, 1983).

The Chert Terrane is bordered by a Paleozoic clastics and chert unit consisting of quartzite, maroon and green argillite, lithic sandstone and siltstone, gray chert, limestone, and limestone breccia. This unit may

Figure 1. Geology of the Livengood B-3, B-4, C-3 and C-4 Quadrangles, Alaska. Generalized from Albanese, 1983; Bundtzen, 1983; Robinson, 1983, and Smith, 1983. (continued on following page)





Figure 1 (continued).



EXPLANATION

Intrusives (Cretaceous-Tertiary)

- Qtz porphyry (small bodies controlled by N-S trending faults)
- Syenite
- Qtz monzonite

Mesozoic flysch (Jurassic-Cretaceous)

- Siltstone, sandstone, conglomerate

Rampart group (Permian-Triassic)

- Basalt, andesite
- Greenstone
- Diorite
- Chert, shale, siltstone

Tolovana Limestone (Silurian-to-Middle Devonian)

- Limestone

Devonian clastics (mid-Devonian) and mafic complex

- Sandstone, shale, conglomerate, chert
- Volcanic flows, agglomerate
- Diorite, gabbro, greenstone
- Serpentinized ultramafic mafic rx

Paleozoic clastics and chert (Silurian-Devonian)

- Chert, maroon and green slate, sandstone, siltstone, limestone
- Quartzite

Chert terrane (Ordovician-Silurian)

- Gray and tan chert
- Variegated chert
- Greenstone, mafic volcanics
- Limestone, dolomitic limestone

Cambro-Ordovician unit

- Orthoquartzite, mafic igneous rocks, phyllite, limestone, chert, and slate

Grit unit (Cambrian-Precambrian)

- Quartzite, maroon-green slate, chert, sandstone
- Limestone, limestone breccia
- Bimodal quartzite (grit)
- Bimodal quartzite, maroon and green slate

represent a transitional phase between the underlying Ordovician and Silurian Chert terrane and a Devonian clastic sequence which occurs stratigraphically or structurally above the Paleozoic clastics and chert unit.

The Devonian clastic sequence, informally called the Cascaden Ridge Unit (Weber and others, 1985, p. 11) is a turbidic sequence composed of a basal conglomerate containing locally derived rounded chert and pebble clasts in a sandy matrix overlain by a thick section of thinly layered shale, siltstone and sandstone with very minor carbonate layers. Although turbidite features such as flute clasts and graded bedding are very rare, debris flows occur near the base of the conglomerate (Weber and others, 1985, p. 22-23). This unit is associated with a dismembered mafic-ultramafic complex containing a wide variety of rock types including gabbro, diorite, serpentine, agglomerate, and mafic volcanics. This complex appears to be tectonically mixed with the clastic rocks (Robinson and others, 1983). Fossil evidence from the Cascaden Ridge unit (primarily gastropods and pelecypods) indicates a Middle Devonian (Eifelian) age and represents a near-shore subtidal or intertidal(?) environment (R. Blodgett, personal commun., 1982). The upper part of the Tolovana Limestone, exposed as a thrust slice in the Livengood B-3 Quadrangle may represent a shelf-facies equivalent of the Devonian clastic sequence.

The Rampart Group, a heterogenous assemblage of rocks consisting of gabbro, pillow basalt, diorite, chert, conglomerate, sandstone and shale is present in the northern part of the map area. The Rampart Group is considered to be Permian to Triassic based on Permian pelecypods and bryozoans (Brosge and others, 1969) and a recent Upper Triassic radiolarian age date (D. Jones, personal commun.). In addition to the fossil evidence is a K-Ar age data average of 205 ± 6 m.y. on hornblende from gabbro in the unit (Brosge and others, 1969). The Rampart Group appears to be in fault contact with older rocks to the south (Robinson and others, 1983) and is not considered to be part of the Livengood stratigraphic section.


A thick sequence of clastic Jurassic to Cretaceous flysch unconformably overlies the Grit unit in the southeastern part of the map area, and the Devonian clastic sequence near Livengood (Robinson and others, 1983). Paleocurrent and point count data collected by Bundtzen (1983) suggest that the Devonian clastic sequence was, in part, the source rock for the Mesozoic flysch.

Felsic to intermediate plutons and dikes of presumed Tertiary to Cretaceous age (Chapman and others, 1971) intrude rocks of both the Mesozoic flysch sequence and the Devonian clastic sequence near Livengood. The intrusives generally range from monzonite to quartz monzonite in composition, with minor syenite phases north of Tolovana Hot Springs Dome. Some of the intrusive outcrops appear to be structurally controlled along north-south trending faults paralleling the Tolovana River Valley.

Regional Correlations

Chapman and others (1979) proposed correlations between Livengood stratigraphy and stratigraphy in the Charlie River Quadrangle, and suggested right lateral displacement along the Tintina fault of about 300 km (fig. 2). The correlation of the Livengood Dome chert (part of the chert terrane) with the

Figure 2. Inferred Stratigraphic Relationships



(1) indicates correlation of Champan and others, 1979

(2) indicates correlation of Robinson and others, 1983

Road River Formation extends this correlation from Livengood to the Selwyn basin in the east-central Yukon Territory (Chapman and others, 1979). Additional correlations between Livengood and Yukon Territory stratigraphy have more recently been proposed. A comparison of the restored stratigraphic column of Livengood with Yukon stratigraphy is shown in figure 2.

The Hadrynian 'Grit unit' in the east-central Yukon Territory of Gabrielse and others (1973) can be compared to the Livengood grit unit. In the east-central Yukon Territory, the grit unit consists of gritty quartz arenite, pale brown slate, and minor limestone overlain by maroon, purple, and green slates with lenses of fine-grained quartz-arenite and brown slate (Gordy, 1978). This lithology is notably similar to the grit unit of Livengood. The east-central Yukon Territory Grit Unit and the Livengood grit are underlain by the Road River Formation and the equivalent Livengood Dome chert of the chert terrane, respectively. In addition to the lithologic and stratigraphic similarities of these two grit units, the age assigned to these units is also similar.

In addition to this correlation, the Lower Earn Group of east-central Yukon Territory which consists of siliceous shale, chert, quartz, sandstone, grit, and chert pebble conglomerate and spans most of the Devonian (Gordy and others, 1982) exhibits lithologic and chronologic similarities to the Cascaden Ridge unit of Livengood. Stratigraphic similarities are also observed as the Lower Earn group in part, conformably overlies the Road River Formation (Gordy and others, 1982), while the Cascaden Ridge unit overlies the unit that is transitional to the chert terrane. These correlations, first suggested by Robinson and others (1983) have been supported by Templeman-Kluit (1984) and Hall and others (1984).

Mineral Potential

Gold placers in the Livengood area have produced over 375,000 ounces of gold. There are three types of auriferous or potentially auriferous placer deposits in the Livengood area. These are: 1) gold-bearing fluvial gravels overlain by frozen colluvial silt such as those currently being mined in the lower Livengood bench on Livengood Creek; 2) poorly preserved, locally derived, potentially auriferous, high-level terrace deposits found along Hess, Fish and Lost Creeks; and 3) gold-bearing alluvial- and colluvial-fan deposits and gravels, such as those on Amy Dome and Money Knob, east of Livengood (Waythomas and others, 1984).

The best exposed lode gold occurrence in the area is the Old Smoky Prospect on Money Knob, 3 km southeast of Livengood. Gold-antimony mineralization occurs in sheared and hydrothermally altered contact zones between Devonian clastics and an intrusive suite composed of biotite monzonite, feldspar porphyry, and felsic dikes. Reported gold values of rock samples from the Old Smoky prospect range from 0.1 to 29.8 ppm (Allegro, 1984).

Although these altered intrusives and associated vein deposits probably contributed heavily to the placer deposits, their volume at the present erosion surface may not be sufficient to constitute a viable source for all of the placer deposits, suggesting that at least a portion of the lode source for the gold was above the present exposures and has since been removed by erosion.

Other units containing mineral potential include the Chert Terrane which contains stratabound limonitic chert breccia zones with gold concentrations up to 1.3 ppm; the Paleozoic clastics and chert unit which contains zones of layered pyritic massive sulfides in quartzites in the northern part of the map area; the Rampart Group which contains minor copper mineralization in mafic igneous rocks; and the contact zone between the chert terrane and the Cascadian Ridge which contains significant stream-sediment zinc anomalies. It is important to note that these zinc anomalies are hosted within the units that are correlated with the lower and middle Paleozoic units of the Yukon Territory which host all of the known shale-hosted stratiform lead-zinc deposits of the Yukon and northern British Columbia.

BIBLIOGRAPHY

- Albanese, M.D., 1983a, Geochemical Reconnaissance of the Livengood B-3, B-4, C-3 and C-4 Quadrangles, Alaska; Summary of data on stream-sediment, pan-concentrate, and rock samples: Alaska Division of Geological and Geophysical Surveys Report of Investigations 83-1, plates, scale 1:63,360.
- Albanese, M.D., 1983b, Bedrock Geologic Map of the Livengood B-4 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys, Report of Investigations 83-3, 1 plate, scale 1:40,000.
- Allegro, G.L., 1984. Geology of the Old Smoky Prospect, Livengood C-4 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys, Report of Investigations 84-1, 9 p., 1 plate, scale 1:120.
- Brosigé, W.P., Lanphere, M.A., Reiser, H.N., and Chapman, R.M., 1969, Probable Permian age of the Rampart Group, Central Alaska: U.S. Geological Survey Bulletin 1294-B, p. B1-B18.
- Bundtzen, T.K., 1983, Bedrock Geologic Map of the Livengood B-3 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys, Report of Investigations 83-6, 1 plate, scale 1:40,000.

- Chapman, R.M., Weber, F.R., and Taber, Bond, 1971. Preliminary Geologic Map of the Livengood Quadrangle, Alaska: U.S. Geological Survey Open-file Report 71-66(483), 2 sheets, scale 1:250,000.
- Chapman, R.M., Weber, F.R., Churkin, Michael Jr., and Carter, Claire, 1980, The Livengood Dome Chert, a new Ordovician Formation in Central Alaska, and its Relevance to Displacement on the Tintina Fault, in Shorter Contributions to Stratigraphy and Structural Geology, 1979: U.S. Geological Survey Professional Paper 1126, p. F1-F13.
- Church, R.E., and Durfee, M.C., 1961, Geology of the Fossil Creek area, White Mountains, Alaska: Fairbanks, University of Alaska, M.S. thesis, 96 p.
- Churkin, Michael, Jr., and Brabb, E.E., 1965, Occurrence and Significance of Oldhamia, a Cambrian Trace Fossil, in East Central Alaska: U.S. Geological Survey Professional Paper 525D, p. D120-D124.
- Gabrielse, H., Blusson, S.L., and Roddick, J.A., 1973, Geology of Flat River, Glacier Lake, and Wrigley Lake Map-areas, District of Mackenzie and Yukon Territory: Geological Survey of Canada, Mem. 366, 153 p.
- Gordy, S.P., 1978, Stratigraphy and Structure of the Summit Lake Area, Yukon and Northwest Territories: in Current Research Part A, Geological Survey of Canada paper 78-1A, p. 43-48.
- Gordy, S.P., Abbott, J.G., Orchard, M.J., 1982, Devonian-Mississippian (Earn Group) and Younger Strata in East-central Yukon: in Current Research Part B, Geological Survey of Canada paper 82-1B, p. 93-100.
- Hall, M.H., Smith, T.E., and Weber, F.R., 1984, Geologic Guide to the Fairbanks-Livengood Area, East-Central Alaska, A Guidebook for the Geology of the Yukon-Tanana Uplands Field trip of the 80th Annual Meeting of the Cordilleran Section of the Geological Society of America, Anchorage, Alaska 1984. Unpublished report, 30 p.

- Robinson, M.S., 1983, Bedrock Geologic Map of the Livengood C-4 Quadrangle Alaska: Alaska Division of Geological and Geophysical Surveys. Report of Investigations 83-4, 1 plate, scale 1:40,000.
- Robinson, M.S., Smith, T.E., Bundtzen, T.K., and Albanese, M.D., 1982, Geology and Metallogeny of the Livengood Area, East-central Alaska; A progress report: Presented at the Alaska Miners Association Convention, Anchorage, October 1982.
- Robinson, M.S., Smith, T.E., Bundtzen, T.K., and Albanese, M.D., 1983, Geology and Metallogeny of the Livengood Area, East-central Alaska: Presented as a poster session at the Alaskan Geological Society Symposium, Anchorage, Alaska, May 1983.
- Smith, T.E., 1983, Bedrock Geologic Map of the Livengood C-3 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys. Report of Investigations 83-5, 1 plate, scale 1:40,000.
- Templeman-Kluit, Dirk, 1984, Counterparts of Alaska's Terranes in Yukon: Cordilleran Section of the Geological Association of Canada Symposium, Vancouver, B.C., Canada, February 1984, 4 p.
- Waythomas, C.F., Ten Brink, N.W., and Ritter, D.F., 1984, Surficial Geology of the Livengood B-3, B-4, C-3 and C-4 Quadrangles, Alaska: Alaska Division of Geological and Geophysical Surveys, Report of Investigations 84-6.
- Weber, F.R., Smith, T.E., Hall, M.H., and Forbes, R.B., 1985, Geologic guide to the Fairbanks-Livengood area, east-central Alaska. Prepared in conjunction with the meeting of the Pacific Section of the American Association of Petroleum Geologists, the Society of Economic Paleontologists and Mineralogists, and the Society of Exploration Geophysicists, May 22-24, 1985: Alaska Geological Society, Anchorage, Alaska, 44 p.