FE Mt. McKinky 66

PE 66-1

PE-066-01

## REPORT ON

### SHANNON LODE PROPERTIES

### Slippery Creek Area

## Kantishna Recording District, Alaska.

Ву

Eerl R. Pilgrim

October 24, 1929

Location	n	•	•	•	•	•	٠	•	•	•	•	•	•	•	-	•	•	•	Т
Transpor	rtati	on																	1
Timber A	Avail	.abl	.e	fo	r	Mı	n	ine	*						•				2
Fuel .				-									•	•					2
Water Po	ower																		2
Geology	and	Kin	er	el	.i 2	at	10	n								-			3
Native 0	egao <mark>.</mark>	r P	ro	sp	ec	t										•			7
Cinnibar	r-Ant	1=0	ny	D	er	00	1 <b>1</b> 1	វន											8
Stibnite	e Vei	n.													-				9
Photogra	aphs,	Мв	פֿסַו	3,	et	c.													

or ancione size is top principle to Employee

· .

Fairbanks, Alaska October 24, 1929

Mr. B. D. Stewart Territorial Mining Department Alaska Territory Juneau, Alaska NOTED NOV1 4 1929

C B. STEWART

Dear Sir:

I herewith present a brief report on the Shannon Lode
Properties located in the Slippert Creek Area, Kantishna
Recording District, Alaska. These claims were examined by me
in September of this year. Five days were spent at the properties which was ample time considering the small amount of development work done on the claims. A careful examination was made of
all outcrops and open cuts and a rough triangulation established
using a brunton compass and an aneroid barometer. A map constructed from this work accompanies this report.

The Shannon Properties consist of the three groups listed below:

Slippery Creek- Iron Creek Copper Group 12+66-10

Terminus Claim, Green Back Claim, Magnetite Claim, Old Sourdough

Native Copper Prospect

- K 66-38

Question Mark Claim

Cinnibar-Antimony Deposite \_ / 66 37

Merinser Claim

#### Location

These claims are located in the Slippery Creek Ares of Mt. McKinley National Park immediately to the north of Mt. McKinley at the heads of Iron Creek, Slippery Creek and the East Fork of Birch Creek. Shannon's Camp at the head of Slippery Creek is approximately 9½ miles from the center of Mt. McKinley and at an elevation of 3200 feet. The claims lie at elevations varying between 3400 feet and 4310 feet.

#### Transportation

This area can be reached at present by aeroplane or by pack animal in summer and by aeroplane or by dog team in winter.

From the end of the Park Highway, which is now completed to the East Fork of the Toklat River, it is approximately 70 miles by trail to this area. The Park Road is now completed 41 miles from McKinley Park Station on the Alaska Railroad. By aeroplane the property can be reached from Fairbanks within 2½ hours. A railroad constructed by the shortest available route would be approximately 108 miles in length to connect with the Alaska Railroad at Lignite or at Kobe. A highway would follow the present trail and be about 110 miles from McKinley Park Station.

#### Timber Available For Mining

The nearest timber available to this area is about 6 miles north in the Slippery Creek Valley and with an easy grade for hauling. This timber is within the park but extends alongsthe valley for many miles and better timber can be found outside of the park line which is about 11 miles from Shannon's Camp. This timber is principally spruce and suitable for mining purposes only.

#### Fue1

There is no fuel available other than the possibility that a commercial seam of coal might be found in the coal formation, reported by the U.S.G.S and various prospectors, on Coal Creek a fork of Clear Water Creek a tributary of Mc Kinley River. No details are known as to the size of any deposites but coal has been reported. It can be reached from Slippery Creek by a road approximately 20 miles in length.

#### Water Power

Most of the rivers and creeks on the north slope of the Alaska Range are fed from glaciers and carry such a great amount of gravel, sand, and mud that their waters could not be used for power development. Clear Water Creek as its name implies is a swift stream of clear water about 11 miles east of Slippery Creek. Clear Water Creek is fed from springs, rains and melting snows, and fluctuates much less in volume than neighboring glacier fed streams.

- \_3 -

A rough estimate of the flow at times of crossing at its mouth where it empties into the McKinley River is 4,000 cubic feet per minute. This stream is said to have an excellent site for a dam and power plant several miles above its mouth.

#### Geology and Mineralization

The predominating rocks of this area consist of paleozoic sediments deformed, tilted and penetrated by a large number of
intrusives. These sediments, (Post-Tonzona" Capps in Toklat-Tonzona
River Region U.S.G.S. Bull.792, 1925) consist of argillites, shales,
slates, and thin bedded limestones in places much silicified and
mineralized by contact metamorphism. The intrusives occur as dikes,
and sills generally trending parllel to the strike of the sediments.
They vary in composition and structure from granite to diabasic
porphry.

As we have divided the Shannon holdings into three groups it is necessary to describe three distinct types of mineralization as they must be each considered seperately.

Slippery Creek - Iron Creek Copper Group

This mineralization is distinctly a contact metamorphic mineralization in which older limestones, determined by Mr. Hopkina ar e dolováte ins as a dolomite, ( His report of Oct. 9 1929 Sample No. A, B, and C.), have been intruded by granitic rocks altering the sediments and introducing the metallic minerals. These minerals have replaced the sediments in a more or less irregular sequence with the predominating minerals depositing first. The probable order of replacement is magnetite, chalcopyrite, sphalerite, pyrite, pyrrhotite, and galena. This order does not hold true however in all outcrops examined. The two largest exposures on this mineralization at Iron Creek and at Slippery Creek show very little magnetite while at the Terminus Discovery Outcrop and the Magnetite Discovery Cut the magnetite predominates over all other metallic minerals by two to one. A hand specimen taken at the open cut Magnetite Discovery shows massive black fine grained magnetite with small seams

and fractures filled with chalcopyrite and sphalerite.

The mineralizing intrusive is a granitic dike cutting almost vertically through the sediments. Its width wad not determined but is at least several hundred feet. Its strike along the contact is about N 40° E.or parallel to the bedding planes of the limestone. Directions taken in the limestones at various points on the four claims show strikes of from North to N 50° E. with a general average of about N.30° E. and a dip of from 60° S.E. to vertical. Smaller dikes or tongues extend out to the side and penetrate the sedimentery. These are similar to the parent dike except for a considerable loss of silica and darker minerals in replacing the lime. The lime is altered greatly close to these contacts. An excellent example of these tongues is shown in photograph marked "D". This is a photograph of the Large outcrop at the head of Slippery Creek on the Green Back Claim.

The mineralization has extended into the sedimentaries for considerable distances. A measurement across the exposure on Iron Creek shows a copper - Iron mineralization of 150 feet in width while a similar measurement on the large outcrop on Slippery Creek shows 330 feet measured at N.54° E and reduces to 180 feet across the strike. Also from the float showing on the hillsides and the mineralization showing in several cuts, one is lead to expect as great a width in other places.

Very little work has been done on these claims and the principal exposures available for examination are natural ones. These surface outcrops are somewhat oxidized and leached, although not to a depth of over a few feet, and show much copper carbonate coatings as well as iron oxide. The copper mineralization although very extensive along this contact, in nomplace shows any appreciable quantity of high grade ore. All evidence points to a possible large low grade deposite of chalcopyrite, sphalerite ore.

Ninteen samples were taken on this group of claims, only

one of which represents an average width and value. Most of the open courts are small and only expose the deposite at one point. A A sample taken there would not represent any definate width or section. The samples taken are described below;

- Sample No. 1. Thirty feet above creek level on Slippery Creek

  Outcrop 30 feet N.E.of west limit of mineralization.

  Grab sample showing chalcopyrite Green Back Claim.
  - 2. Channel sample iron stained leached gouge on W.side of open cut Magnetite Discovery Cut. 10 in. wide.
  - 3. Channel sample across 8'-1" from Sl.No.2 east to edge of mineralization. massive magnetite with chalcopyrite, sphalerite, and small amounts of galena.

    Apparantly faulted on west side.
  - 4. Picked pieces from large outcrop on Slippery Creek showing chalcopyrite, sphalerite, and pyrrhotite. This represents as high grade ore as could be found.on this outcrop.
    - 5. Grab sample from Terminus Discovery Outcrop showing magnetite, chalcopyrite, and sphalerite.
  - 6. Grab sample decomposed ground up material on local fault cutting through big fron Creek Outcrop near east edge of outcrop.
  - 7. Picked pieces iron rock from Iron Creek Outcrop.
  - 8. Pieces altered limestone on Iron@reek duterop represents large portion of outcrop.
  - 9. Picked pieces from Magnetic Claim best ore .
  - 10. Brokenpieces grab sample east side Iron Creek Outcrop at creek level.
  - 11. Grab sample from Iron Creek Outcrop 30 feet above No. 1 sample No. 10.
  - 12. Grab sample Terminus Discovery across 8 feet just below discovery post.

Sample No. 13. Same as Sample marked A.

- 14. Picked Specimens not assayed
- 15. Oxidized rock from Slippery Creek Outcrop large portion outcrop this material.
- 16. Green Back Claim Slippery Creek copper carbonate and black mineral.
- 1.A. Showing minerals contained in average contact.
- B. Tested for non metallis minerals.
- C. Tested for limestone.
- 19. Grab sample of contact deposite porphry-limestone on ridge below copper peak, west side of limestone marked mineralized open cut No. 4.on map.

Assays of these samples are listed below;

Sample No.	Gold	Silver	Copper	∄ead	Zinc	Iron
1.	Trace	1.00	3,5			
2.	0.03	0.80	0.6			
3.	0.01	0.20	0.3			•
4.	0.04	4.80	6.7	0.0	23.5	12.6
5.	0,02	1.60	1.6	0.0	2.2	16.8
6.	0.01	0.10	0.0	0.0		
7.	0.01	0.20	0.4	0.0	0.1	17.2
8.	Trace	0.60	0.0			
9.	0.01	5.40	5.6	2.3	4.8	20.1
10.	0.01	0.20	0.2	0.0	0.2	15.4
11.	0.06	1.80	1.7	- 0.0	0.3	24.4
12.	0.02	0.40	1.5	0.0	0.6	5.6
15.	0.02	0.30	0.0			
17,9	Trace	Trace	0.0	1.9		11:5-

#### Native Copper Prespect

The Question Mark Claim is shown on map accompanying this report. It lies about one mile west of Shannon's Camp on Shippery Creek and at an elevation of 3600 feet. The lode is a greenstone dike striking east-west and dipping steeply to the north. This dike is from 30 to 50 feet in width and is plainly to be traced along the surface for several hundred feet by a red covering of residual hematite. The lode is bounded on the north by a dark diabasic porphry rock and on the south by a green porphyritic rock, probably diorite. The greenstone contains a number of stringers of calcite which appear barren of any other minerals where examined.

The native copper occurs finely distributed in the greenstone in minute specks and in fine irregular threads. No work has been done on this claim and there are no cuts or fresh exposures to examine and no information is available as to the extent of the mineralization throughout the greenstone. A microscopic examination will be necessary to clear up the source of the copper. No other copper minerals were observed either as sulphides or as oxidized products. It is very probable that this occurrance is a common one in which the native copper occurring in the basis igneous rock is associated with minerals of the zeolite group. This deposite may be of economic importance but considerable work must be done on it to open it up for examination before any definate value can be given it.

Two samples were taken on this lode;

Sample No. 17. Grab sample from pieces broken off of lode at outcrop 300 feet west of West fork of Slippery Or Creek. Extent of this ore not known but apparantly considerable. Lode about 50 feet wide but sample represents only 5 feet. Remainder of lode covered by iron stained capping and slide rock.

Sample No. 18. Decomposed iron oxide covering lode taken along side of cut of sample No. 17.

Sample No. Gold Silver Copper Iron Soluble

17 Trace Trace 1.8 11.6

18 0:01 0.10 Trace 8.9

#### Cinnibar-Antimony Deposite

The Merinser Claim is located a little over two miles west of Shannon's Camp on Slippery Creek and covers elevations of from 3400 feet to 3820 feet. A group of highly altered steeply, pitching sedimentary rocks strike in a direction of Nc40° E2 this formation, at the crest of a sharp ridge extending in a northerly direction on the north flank of a mountain, is an outcrop showing specks and flakes of cinnibar disseminated through the rock in small amounts. There are a number of calcite stringers cutting the rock and they also carry cinnibar in slightly greater amounts than the sedimentary rock. This mineralization shows exposed across a width of 37 feet with small amounts of cinnibar, lesser amounts of metacinnibar and an occasional globule of native mercury. The highest grade portion of this deposite appears to be along the west side where the calcite stringers appear the man more numerous. There are no contacts or walls or change in the character of the rock on either side of this deposite.

About 200 feet south east and 50 feet lower in elevation there is a granitic dike cutting across the formation with a strike of S. 45° W. and a dip of 30° to the east. This dike may have some relationship to this cinnibar deposite. Apparantly the deposite is a low grade dissemination in porus rock.

No work has been done on this deposite other than an open cut near the top of the ridge crosscutting the deposite. Five samples were taken across this width of 37 feet and one special sample;

Sample No.20 Starting at northwest side of deposite 0-62feet best ore in this sample.

211 6--12 feet.

22. 12--18 feet.

23. 18--26 feet.

24. 26--37 feet

25. Picked pieces of the best ore showing taken from northwest edge of deposite on top of ridge, several calcite stringers cutting through.

Sample Nu	nber	Mercury	Sulphur	Gold	Silver
	20.	0.07			
	21.	0.03			
	22.	0.02			
	23.	0.03			
	24.	0.03	-		
	25	0.00	Trace	Trace	Trace
Composite 20-24ind		0.03	11	Ħ	Ħ

#### Stibnite Vein

About 200 feet lower and 500 feet southwest of the cinnibar prospect, there is an antomony vein outcropping on the side of the mountain. This outcrop can be traced for several hundred feet along the mountain side by the brown stained float which shows below the vein. This vein has a strike of N.50° E, and a dip of 30° to the southeast. The vein varies in thickness of from a few inches to over 11 feet.

Several cuts have been dug along the outcrop but were sluffed in to a considerable extent. Good pure stibnite was noted piled along these cuts. One cut near the northern extremity of the outcropping vein was cleaned out and the vein the terminal extrements a width of 11 feet 6 inches. The vein here

showed about 4 inched of solid stibmite on the foot wall, 5 inches of it on the hanging wall, and several stringers scattered across the face. The gangue rock is a milky quartz with much soft brown decomposed quartz, and antimony oxide. The vein shows about 2 inches of black gouge on the foot wall. Both wall rocks are black shaley limestone.

From the amount of float scattered along the mountainside below the strike of the vein one is lead to expect a considerable extent to the stibnite along the bein but probably greatly
intermixed with the gangue.

Two samples were taken from this vein and a special piece from another reported antimony lead about five miles southwest of the Marinser Claim;

Sampla Number 26. 11'-6" across vein channel sample

- 27. Picked pieces from six outcrops representing of sorted ore from several piles along vein.
- 28. Dre from big lead reported 5 miles southwest.

Sample Number Gold Silverrantimony Lead Goppenczinoparsenic Sulphur Ihon


2618 0.18 0.10 7.2 None None Nome 0.85

27. Trace Trace 56.7 None None None 0.18 21.4 3.2


28. -----33.5 None None None 0.12

Respectfully Submitted,

Earl R. Pilgrim




About 6 miles from Camp on Shippery



Report on Shannon Property Slippary Creak Kantishna Dist.


Report. Shannon Proparty Slippary Croak KantishnaDist

Looking 5.W. From Copper Peak



D Report. Shannon Property Slippery Creak Kantishna Dist.

Report on Shannon property Slippery Creek Kantishne Dist.



E Report on Shonnon Property Slippary Crack Kantishne Dist.



-- ---

1

# United States Department of Commerce

BUREAU OF MINES

Fairbanks, Alaska Oct. 9, 1929

Mr. E. R. Pilgrim, Territorial Mning Engineer, Fairbanks, Alaska.

Dear Mr. Pilgrim;

Minerals have been identified in your samples

as follows:

Sample A contains considerable chalcopyrite also considerable sphalerite and probably pyrite. The gangue appears to be a highly altered limestone, now consisting mainly of a greenish silicate which appears to be in the pyroxene class, probably a variety of augite. No lead was seen in the sample.

Sample B consists mainly of silicates such as the augite of the previous sample with smaller amounts of epidote and similar minerals. It contains a small amount of chalcopyrite and a trace of zinc. No lead minerals were found.

Sample C is a dolomite containing a considerable amount of silicates as in sample B. A very small amount of galena is present and a trace of zinc. No copper was found.

The sample marked Shannon 16 is a dolomite containing a small amount of copper in the form of malachite. The soft dark mineral coating some of the surface is a manganese oxide, probably pyrolusite. This sample contains less of the complex silicates than the preceding samples.

The reddish mineral in sample Shannon 18 is an iron oxide, possibly hematite. It occurs as microscopic included particles. It was not possible to concentrate the red mineral by careful panning.

Very truly yours,

Paul Hopkins, // Assoc. Anal. Ohemist.

U. S. Bureau of Mines.

3762-66

## Alaska Agricultural Gollege and School of Mines

In Cooperation with

Gollege, Alaska

U S. Bureau of Mines, Department of Commerce

Oct. 18, 1929

## Mr. E. R. Pilgrim, Territorial Mining Engineer

On samples received from .....

	75	O1-	OUNCES	PER TON		PERCENTAGE OF						
Assay No. Mark on Sample		Gold Silver		Copper TEPTON	Lesd	Zino	iron (soluble	Arsenic				
01008	Shamo	n #1	Trace	1.00	<b>3.</b> 5							
01009	45	. 2	0.03	0.80	0.5							
01010	#	3	0.01	0.20	0.3							
01011	u	4	0.04	4.80	6.7	0.0	28.6	12.6				
01012	77	5	0.02	1.60	1.6	0.0	2.2	16.8				
01018	18	6	0.01	0.10	0.0	0.0						
01014	vt.	7	0.01	0.20	0.4	0.0	0.1	17.2				
01015	49	8	Trace	0.50	0.0							
01016	12	9	0.01	5.40	5.6	2.3	4.8	20.1				
01017	n	10	0.01	0.20	0-2	0.0	0.2	15.4				
01018	19	11	0 <b>.06</b>	1.80	1.7	0.0	0.3	24.4				
01019	Ħ	12	0.02	0-40	1.5	0.0	0.6	5.6				
01020	61	15	0.02	0.30	0.0							
01021	iÿ	17	Trace	Trace	1.8			11.6				
01022	77	18	0.01	0.10	Trace	<b>4 8 6 6 7 4 7 6 7</b>		8.9				
01023	78	19	Trace	Trace	0.0	1.9			race			

Assayed by,

Paul Hopkins,

Associate Anal. Chemist. U. S. Bureau of Mines.

Official

Total charges for above assays.....

Amount received from sender.....

# Alaska Agricultural Jollege and School of Mines

In Gooperation with

U. S. Bureau of Mines, Department of Commerce

College, Alaska

REPORT OF ASSAY

Oct. 26, 1929

On samples received from Mr. E. R. Pilgrim, Territorial Mining Engineer

	No.   Mark on Sample		OUNCES PER TON					PE	RCENTAGI	e of	
Assay No.			Gold	Silver	Value Per To	1	Merou	ry Sul	phur		
01027	Shannon	#20	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	49657641		, , , , , ,	. 0.0	7		<u> </u>	
01028	n.	21									
01029	79	22	0.0.00	• • +		• • • • •	. 0.0	2			
01030	11	23	******		*****		. 0.0	3			
01031	33	24	* * • • * • * • • • • • • • • • • • • •	4	** 2 5 * * *		. 0.0	3			
28010	13	Composite	Trace	Trace			0.0	3 Tr	aee		
01033	н	25	Trace	Trace			0.20	0 Ir	ace		
				_				Percenta	age of		_
				À	ntimony	Lead	Zinc	Copper	arsonic	Sulphur	Iron
01934	Ħ	26	0.18	0.10	7.2	None	None	None	0.86		
01025	11	27	Trece	Trace	56.7	Mone	None	None	0.18	21.4	3.2
01036	19	28			27 5	Jona		None	0.12		

Asseyed by,

Paul Hopkins,

Assoc. Anal. Chemist, U. S. Bureau of Mines.

Stonolulu Station, Mile 289 alaska Ry., October 24, 1929.

Mr. B. D. Stewart,

Territorial Supermisor of Immes, OFF SO 1929

Juneau, Alaska.

Dear Sir:

Mr. Earl R. Dilgrim, mining engineer, Whom you delegated to examine and report upon my mining properties in the Kantishna District, has completed his examination of the properties.

Mountain, Kantishna, the eneming of August 31st and on September 2nd We left for Slippery creek, where my claims are, and arrived at the claims on the morning of September 4th, but could easily have mode it to the Claims the eneming of September 3rd had he not laid oner

Slippery creek, the afternoon of Septemoer 3rd, on account of a rain storm, the 4th; 5th; 6th; 7th and 8th of Septemoer, we spent in going oner the property, examining the outcrops, collecting Samples and specine, measurening the outcrops, and examining the Country rock hear the deposits.

We brought in, with us, to the Railroad about Senenty, (70), pounds of Samples and Specimens, Mostly specimens, because, as Mr. Dilgrim said, "owing to the lateness of the Season, and therefore lack of Sufficient time before snowfall, and also the humber and great size of the outerops, he would not begin to have anywhere hear time to take anerage Dample's, however, to my mind, he accomplish, almost as much as though he had taken an leage Samples, because there is solunderground workings on the property and the surface appears to be considerably leached on most of the outerops, therefore an anerage Sample taken at the Surface, and especially on the

Copper minerligation, would be no criterior of the Commercial or non commercial Value of the deposits, as I see it.

Should Mr. Pilgrin's report on my properties prone fanorable are we going to let the matter rest there? Why not go right on through and help me to denelop a mine, if a mine be there, that is develop sufficient tomage, if present, to warrant the building of it from a sailword No connect the properties with the Aluska Railroad, by furnishing labor and supplies to be used in the prosecution of said develop ment, and under the direction of a Mining engineer appointed by The Supernisor of Mines, - that may seem assurd to you Mr. Stewart, but to my mind it is more logical to spend territorial monies in giving material assistance to the prospector seen fanously reported upon by as mining engineer in the employ of The Territory, than by just having an examination and

4

report made on the properties, and nothing more, Secause no single prospector, during a natural lifetime, could hope to denelop sufficient tonnage to warrant the building of a hundred miles of railroad, and the prospects Which I have, judging from surface indications, give energy promise of containing sufficient tonnage.

During a recent convention which I had with General Manager Ohlson of the Claska Railroad, he mentioned something about the U.S. Geological Survey going to take oner this work, I can see to admintage in that, I believe you inaugurated this service for the prospectors of the Territory, and in my opinion it should remain under your direction, and I take this opportunity to thank you for the service you have already caused to be performed in my behalf.

Sincerly yours.

# United States Department of Commerce

BUREAU OF MINES

Feirbanks, Alaska Oct. 9. 1929

Mr. E. R. Pilgrim, Territorial Mining Engineer, Fairbanks, Alaska.

Dear Mr. Pilgrim:

Minerals have been identified in your samples

as follows:

Sample A contains considerable chalcopyrite also considerable sphalerite and probably pyrite. The gangue appears to be a highly altered limestone, now consisting mainly of a greenish silicate which appears to be in the pyroxene class, probably a variety of augite. No lead was seen in the sample.

Sample B consists mainly of silicates such as the augite of the previous sample with smaller amounts of epidote and similar minerals. It contains a small amount of chalcopyrite and a trace of zinc. No lead minerals were found.

Sample C is a dolomite containing a considerable amount of silicates as in sample B. A very small amount of galena is present and a trace of zinc. No copper was found.

The sample marked Shannon 16 is a dolomite containing a small emount of copper in the form of malachite. The soft dark mineral coating some of the surface is a manganese oxide, probably pyrolusite. This sample contains less of the complex silicates than the preceding samples.

The reddish mineral in sample Shannon 18 is an iron oxide, possibly hematite. It occurs as microscopic included particles. It was not possible to concentrate the red mineral by careful panning.

Very truly yours,

Paul Hopkins,

Assoc. Anal. Chemist, U. S. Bureau of Mines.

3762-66

## Alaska Agriculturai Jollege and School of Mines

In Gooperation with

College, Alaska

U. S. Bureau of Mines, Department of Commerce

#### REPORT OF ASSAY

Oct. 26, 1929

On samples received from Mr. E. R. Pilgrim, Territorial Mining Engineer

		_	OUNCES PE	ER TON				PH	ERCENTAG	E OF	
Assay No.	Mark on	Sample	Gold	Silver		Ton Mercury		ry Sul	phur		
										<u>'</u>	_
01027	Shannon	#20 <b></b>					. 0.0	7			
01028	Jt	21									
01029	tt.	22					. 0.0	2			
01030	If	23					. 0.0	3			
01031	13	24									
01032	31	Composite	Trace	Trace	•		0.0	3 Tr	ace		
01033	11	25	Trace	Trace			0.20	O Tr	808		
				_			J	Percent	age of		
				ż	Intimony	Lead	Zinc	Copper	Arsenic	Sulphur	Iron
01034	fl	26	0.18	0.10	7.2	Hone	Wone	None	0.85		
01035	>+	27	Traco	Trace	56.7	None	Mone	None	0.18	21.4	3.2
01036	If.	28			33.5	Kone		None	0.12		

Assayed by,

Paul Hopkins,

Assoc. Anal. Chemist, U. S. Bureau of Mines.

Total charges for above assays...... Official

Amount received from sender......

# Alaska Agricultural Sollege and School of Mines

In Cooperation with

U.S. Bureau of Mines, Department of Commerce

College, Alaska

Oct. 18, 1929

#### REPORT OF ASSAY

			OUNCES	PER TON		PERCENTAGE OF					
Assay No.	Mark	on Sample	Gold	Silver	Mains Rentfor Copper	Lead	Zinc	Iron (soluble	Arsenic		
01008	Shanno	on #1	Trace	1.00	<b>3.</b> 5		,				
01009	33	2	0.03	0.80	0.6						
01010	- 13	3	0.01	0.20	0.3						
01011	12	4.	0.04	4.80	6.7	0.0	23.5	12.6			
01012	11	5	0.02	1.60	1.6	0.0	2.2	16.8			
01013	u	6	0.01	0.10	0.0	0.0					
01014	a	7	0.01	0.20	0.4	0.0	0.1	17.2			
01015	It	8	Trace	0.60	0.0						
01016	**	9	0.01	5.40	5.6	2.3	4.8	20.1			
01017	**	10	0.01	0.20	0.2	0.0	0.2	15.4			
01018	**	11	0.06	1.80	1.7	0.0	0.3	24.4			
01019	**	12	0.02		1.5	0.0	0.6	5.6			
01020	11	15	0.02	0.30	0.0						
01021	24	17	Trace	Trace	1.8			11.6			
01022	4	18	0.01	0.10	Trace			8.9			
01023	18	19	Trace	Trace	0.0	1.9 .			frace		

Assayed by,

Paul Hopkins,

Associate Anal. Chemist.

U. S. Bureau of Mines.

				Official
Total	charges	for	above	assays
Amou	nt recei	ived	from	sender