


- UNDIFFERENTIATED ALLUVIAL AND COLLUVIAL SEDIMENTS**
- Qu** UNDIFFERENTIATED ALLUVIAL AND COLLUVIAL SEDIMENTS
- Tertiary Volcanic Rocks**
- Ages for the Tertiary volcanic rocks based on similarity with rocks mapped as Tertiary by Csejey and others (1978).
- Tf** FELSITE/RHYODACITE FLOW (Tertiary)—Light gray to white, finely porphyritic rhyodacite with subtle planar parting. Phenocrysts compose about 7% of the rock and include quartz (rounded, embayed), potassium-feldspar (euhedral, concentrically zoned), plagioclase (large, euhedral), and biotite (fresh, small, euhedral). Matrix is very fine grained but has a bimodal size distribution. Larger matrix crystals are strongly aligned (trachytic), generally euhedral, elongate plagioclase, and square crystals of quartz (?) and (or) potassium feldspar (?). This material is mixed with an even finer matrix of granular plagioclase, quartz (?) and potassium feldspar (?). Crops out in headwater region of the east fork of Iron Creek.
- Tv** LITHIC TUFF-IGNIMBRITE, VOLCANIC BRECCIA, AND CRYSTAL TUFF (Tertiary)—Bright red-stained volcanic rocks with roughly horizontal layering between the coarse tuff breccia and smaller clast varieties. Lithic tuff-ignimbrite has a maroon tuffaceous matrix with dark gray angular clasts (up to 10 cm in diameter) composed of greenstone (Rg?) rock fragments and other brecciated volcanic material. Crystal tuffs are white to pale tan, fine grained, granular/sugary, vuggy, and may be siliceous and clay altered. Crops out in the northeastern map area.
- Tb** BASALT LAVA FLOWS (Tertiary)—Deep orangeish-red weathering, dark gray, olive(?)-phyric basalt flows with local column-jointing. Generally flat-lying flows that unconformably cap hills in the northeastern map area.
- Tertiary to Mesozoic Igneous rocks**
- Tf** FELSIC DIKES (Tertiary? to Mesozoic?)—Fine-grained felsic dikes of granitic composition. Granite dikes are found in northeast, and are generally quartz-rich, fine-grained, and equigranular. Apatite dikes intrude the large multi-phase pluton (TMzmp) south of Iron Creek and the granodiorite (TMzgd) in the west, are fine-grained, and exhibit anhedral granitic texture. Dacite dikes occur in the headwater region of the east fork of Iron Creek and are gray and white, and porphyritic to seriate textured. Dacites contain large phenocrysts of plagioclase, potassium feldspar, quartz, and biotite that are partially altered to clay sericite, epidote, and chlorite. Matrix consists of feldspars, quartz, chloritized biotite, apatite, and tiny opaque mineral(s).
- Tmzd** TRACHYTE DIKES (Tertiary? to Mesozoic?)—Light gray, porphyritic trachyte dikes. Intrude the greenstone unit (Rg) and are spatially associated with small syenite bodies within the south end of the largest granitic pluton (TMzgr). Abundant phenocrysts of square, color-zoned (pink-gray-white) potassium feldspar and white, elongate plagioclase. Hornblende, or perhaps pyroxene (?), occurs as dark green, 1- to 3-mm-long, faintly aligned phenocrysts. Locally the dikes contain xenoliths of greenstone (Rg).
- Tmgd** GABBRO (Tertiary? to Mesozoic?)—Dark brownish-black, very-fine-grained gabbro. Matrix predominantly has an equigranular, diabasic, subophitic texture, with sparse local ophitic texture. Contains fresh plagioclase (about 35% iddingsite (?) or splene (?), clinopyroxene (?), late-crystallizing opaque mineral(s), and about 1% calcite. Rare amygdalites filled with calcite and (or) quartz. Calcite occurs in thin cross-cutting veins. May be intrusive (?) into the greenstone (Rg).
- Tmgz** GRANULAR GRANITE (?) (Tertiary? to Mesozoic?)—White, fine-grained (about 1.5 mm average grain size), quartz-rich (up to 50%), sugary granular to seriate-textured pluton (of uncertain composition) with no mafic minerals. Composed of quartz and a white, anhedral, undifferentiated feldspar. Quartz occurs as anhedral grains intergrown with feldspar, and as slightly larger rounded grains up to 3 mm in diameter. Contains up to 5% orangeish-tan limonite within intergranular spaces and as fracture coatings. Commonly cut and (or) bounded by northeast-trending high-angle faults.
- Tmzt** BIOTITE GRANITE (Tertiary? to Mesozoic?)—Tan, fine- to medium-grained, generally equigranular quartz-rich (up to 40%) granite. Occurs as deeply eroded, non-resistant plutonic bodies and as small apophyses and dikes. Commonly cut and (or) bounded by northeast-trending high-angle faults. Locally contains fluorite and pegmatites. Primary minerals include quartz (large grains with undulatory extinction), plagioclase (subhedral to rarely euhedral, albite-twinned, compositionally zoned with altered cores and fresh rims), potassium feldspar (late, anhedral, often as perthite intergrowths with plagioclase), biotite (sparse, forms large crystals that are partially altered to chlorite, white mica, and (or) secondary opaques), apatite, and zircon. White mica commonly replaces feldspars. Some myrmekite.
- Tmzb** BIOTITE-HORNBLLENDE GRANODIORITE (Tertiary? to Mesozoic?)—White and black, fine-grained, equigranular granodiorite. Potassium feldspar occurs as large, concentrically zoned, subhedral crystals with or without perthitic intergrowths with plagioclase. Plagioclase occurs as large, subhedral to euhedral, albite- and Carlsbad-twinned, concentrically zoned crystals. Feldspars are generally fresh but may contain trace sericite alteration. Quartz occurs as large anhedral grains that were deposited late in the open space between feldspar crystals. Biotite occurs as large subhedral crystals, some of which are partially altered to chlorite ± epidote. Hornblende forms large, euhedral to subhedral, occasionally twinned crystals and rarely has cores of clinopyroxene. Primary opaques vary widely in size, have irregular to highly irregular (nebulous) shapes, and usually occur with mafic minerals. Apatite and zircon are common accessory minerals.
- Tmzd** GRANODIORITE AND QUARTZ DIORITE (Tertiary? to Mesozoic?)—Tan and black, fine- to medium-grained granodiorite and lesser quartz diorite. Texture varies from equigranular to seriate and poikilitic. Primary minerals include quartz, plagioclase, potassium feldspar, hornblende, biotite, apatite, and opaque mineral(s). Mafic minerals typically form about 15% of the rock.
- Tmzd** BIOTITE-HORNBLLENDE GRANITE AND TONALITE (Tertiary? to Mesozoic?)—Undifferentiated, foliated, multi-phase pluton composed of white biotite-hornblende monzogranite and biotite tonalite. Biotite-hornblende monzogranite is white, quartz-rich, hornfelsed, and contains 10% mafic minerals. Primary minerals have been partially recrystallized and altered, and include quartz, plagioclase, potassium feldspar, biotite, hornblende, and minor apatite and opaques. Biotite is extensively altered to secondary biotite, chlorite, epidote, and secondary opaques, and hornblende is partially altered to chlorite.
- Tmzt** BIOTITE-HORNBLLENDE TONALITE (Tertiary? to Mesozoic?)—White and black, fine- to medium-grained, equigranular, locally foliated biotite-hornblende tonalite or granodiorite. Primary minerals include quartz, plagioclase, potassium feldspar, hornblende and biotite, apatite, and opaque mineral(s). Secondary minerals include sericite, clay, calcite, clinzoisite and (or) epidote, and large anhedral splene crystals.
- Tmzb** BIOTITE TONALITE (Tertiary? to Mesozoic?)—White, medium-grained, porphyritic biotite tonalite. Phenocrysts include plagioclase, with or without concentric zoning, and quartz up to 1 cm in diameter in a medium gray matrix. Potassium feldspar occurs as perthite and as concentrically zoned crystals with clay-altered cores. Biotite occurs throughout the groundmass as large primary crystals surrounded by margins of smaller secondary biotite, and it also is partially altered to chlorite, sericite, epidote, secondary opaques and (or) calcite. Contains accessory apatite and large primary opaques.
- Kmp** INTERMEDIATE COMPOSITION PLUTON (Cretaceous)—Typically fine- to medium-grained, equigranular, diorite, quartz diorite, tonalite, and granodiorite. Generally highly magnetic. Exposed in a large, deeply eroded plutonic body in the southern map area. Typically contains both biotite and hornblende although biotite- and hornblende-only varieties are present locally. Porphyritic phases, mafic-rich phases, and occasional minor foliation is present near the margins. In the cirque south of Iron Creek apatite dikes intrude the pluton in an area characterized by a low aeromagnetic signature (Staff, 1998). "Ar" Ar hornblende plateau age of 75.5 ± 0.6 Ma (sample 9; table 1).
- Tmzd** BIOTITE-HORNBLLENDE DIORITE (Tertiary? to Mesozoic?)—Coarse-grained diorite with 30% mafic minerals. Pluton is metamorphosed, but relict equigranular texture is preserved. Primary minerals include plagioclase, quartz, hornblende, remnant clinopyroxene, and biotite(?). Plagioclase occurs as anhedral albite-twinned crystals intergrown with anhedral quartz. Hornblende is partially recrystallized and altered to secondary hornblende, epidote, and secondary opaques. Biotite occurs as small randomly oriented secondary grains in patches (pseudomorphs after primary biotite?) and is partially altered to chlorite and secondary opaques. Pseudomorphs after probable orthopyroxene consist of chlorite, splene, and opaque mineral(s).
- Tmzd** HORNBLLENDE DIORITE (Tertiary? to Mesozoic?)—Medium gray, fine- to medium-grained hornblende diorite composed of 25-30% hornblende and 70-75% plagioclase. Appears equigranular but contains poikilitic hornblende crystals up to 2.5 cm in diameter. Plagioclase partially altered to epidote.
- Tmzd** HORNBLLENDE (Tertiary? to Mesozoic?)—Dark green, medium grained, equigranular hornblende composed of primary oriented hornblende crystals (~30%), intergranular plagioclase (~10%), and opaque minerals (<1%, magnetite). Hornblende laths up to 1 cm.
- Tmzd** TRONDHJEMITE (Tertiary? to Mesozoic?)—White to very light green, fine- to medium-grained, equigranular trondhjemite. Primary minerals include large anhedral quartz and subhedral to anhedral plagioclase. Albite twinning and myrmekite common. In some areas, up to 95% of the rock consists of graphically intergrown quartz and plagioclase crystals. Primary mafic minerals were very minor (<1%) and have been completely altered to epidote, secondary opaques, chlorite, and (or) minor sericite. Forms large plutonic bodies in the central and southwestern map area.
- Triassic Metasedimentary and Metavolcanic Rocks**
- Rl** LIMESTONE (Triassic)—Light tan and gray, relatively fine grained limestone with a planar to mottled color distribution. Occurs in several stratigraphic horizons within a metasedimentary-metavolcanic unit (Rsv) in the eastern map area. Thickness of limestone beds varies from a meter to about 60 meters, and is locally structurally thickened, folded, and rarely brecciated. Average calcite grain diameter is <0.04 mm. Limestone is pure, but locally contains minor detrital quartz and white mica, rhomb-shaped dolomite(?) crystals, disseminated pyrite, calc-silicate alteration, and silty lites. Contains Late Triassic conodonts (B. Wardlaw, written communication, 1999).
- Rsv** METASEDIMENTARY AND METAVOLCANIC ROCKS (Triassic)—Sequence of interlayered metasedimentary and metavolcanic rocks of unknown thickness. Metasedimentary rocks include Late Triassic limestone (Rl), minor white, finely crystalline dolomite, and sparse maroon shale. Metavolcanic rocks include fissile gray tuff with white feldspar phenocrysts, dark green, very fine grained, highly foliated chloritic phyllite ± oval-shaped chlorite patches along foliation, and orange to pale maroon weathering, fissile, felsic tuff (?). Metasedimentary and metavolcanic unit locally intruded by finely crystalline, metamorphosed, ± amygdaloidal microgabbro sills. A white, fine-grained quartz-white mica schist horizon within the metasedimentary-metavolcanic unit extends for about 1 km along strike. Quartz-white mica schist contains disseminated pyrite and is locally intensely cut by quartz veins. The high silica content suggests the schist is the product of hydrothermal alteration.
- Rg** GREENSTONE (Triassic)—Dark forest green, variably magnetic, massive outcrops of basalt, microgabbro, and greenstone. Metamorphosed in lower greenschist facies and variably foliated. Textures include amygdaloidal, massive, and subophitic, to less commonly porphyritic and (or) glomeroporphyritic with 1- to 3-mm-long plagioclase phenocrysts. Primary minerals include intergrown plagioclase and clinopyroxene, and late-crystallizing opaques. Primary textures and minerals partially to completely replaced during metamorphism and deformation. Metamorphic minerals include abundant epidote, actinolite, clinzoisite, calcite, chlorite, albite, pumpellyite, secondary opaques, and minor quartz and sericite. Amygdalites commonly filled with quartz, epidote, calcite, chlorite, and pumpellyite, and rarely by pyrite, calcopyrite, and (or) hematite. The greenstone is locally cut by veins and shear zones filled with quartz, epidote, actinolite, chlorite, calcite, albite, calcopyrite, malachite, hematite, pyrite, hematite, and (or) minor magnetite. Tertiary thought to be Triassic based on spatial association with Triassic limestone (Rl). Possibly correlates with the Amphitheater Group about 150 km to the northeast (Smith, 1981).
- Rgs** GREENSTONE WITH MAFIC SILLS (Triassic)—Greenstone (Rg) intruded by volumetrically minor (about 5%) mafic sills. Sills are medium to dark green, fine-grained, and equigranular, diabasic texture is common. Sill composition ranges from diorite to gabbro. Primary minerals include plagioclase, clinopyroxene, and opaque mineral(s). Metamorphic minerals include epidote, actinolite, clinzoisite, chlorite, and secondary opaques.
- Triassic to Paleozoic Metasedimentary and Metavolcanic Rocks**
- RPzsv** METASEDIMENTARY AND MINOR METAVOLCANIC ROCKS (Triassic? to Paleozoic?)—Generally light colored metasedimentary rocks with lesser metavolcanic rocks, interbedded on the scale of millimeters to tens of centimeters. Dip of most beds is near vertical. Rock types include: 1) pale green- and orange-weathering, light greenish-gray, very finely granular siliceous metagabbro or sandstone; 2) thinly planar bedded, gray to pale pinkish-green, fine-grained, interbedded sandstone and dark gray siltstone; 3) minor, medium to light gray, tan, white, and pale maroon, thinly bedded, finely granular marble/limestone; 4) argillite; 5) chert; and 6) minor pale green- and orange-weathering, light greenish-gray, meta-volcaniclastic sandstone layers. Some granular siliceous metasedimentary rocks are hornfelsed near granite (TMzgr), and limestones locally contain 2-3 mm diameter garnets. In the northwestern map area (near hornblende [TMzb]), there is a 10-meter-thick band of white, equigranular, finely crystalline (1-3 mm average grain diameter), slightly impure (1% epidote and clinopyroxene), calcite marble of unknown age.
- RPzms** MAFIC SILLS WITH MINOR METASEDIMENTARY AND METAVOLCANIC ROCKS (Triassic? to Paleozoic?)—White and green, or dark green mafic sills and minor metasedimentary and metavolcanic rocks (RPzsv). Mafic sills are fine- to medium-grained, and range in composition from diorite to gabbro to rarely clinopyroxenite(?). Generally equigranular but occasionally diabasic textured. Most sills are composed of plagioclase, clinopyroxene, and primary opaque mineral(s). One dioritic sill contains randomly oriented hornblende laths intergrown with plagioclase.
- TPzsu** METAVOLCANIC AND METASEDIMENTARY ROCKS (Paleozoic?)—Tuffaceous and volcanoclastic rocks and minor metasedimentary rocks. Commonly thinly bedded and tightly folded. Locally contains diabase and microgabbro sills or flows (?). Rock types include: 1) tuffaceous sandstone and crystal-lithic tuff; 2) medium green, medium grained, volcanic conglomerate and tuff breccia with a tuffaceous matrix; 3) pale green, very-fine-grained fissile tuff; 4) tanish-gray to brown, fine grained, felsic sandstone, siliceous mudstone, and argillite; ± thin limestone and rare chert partings; and 5) light tanish-gray, well-sorted, clay-supported limestone breccia cemented by a tan calcareous clay matrix. Hornfelsed adjacent to intrusions. Tertiary correlated with Csejey and others (1978) Psv unit which they interpreted as a Paleozoic submarine volcanic arc.





- Maps Showing Location of Study Area**
- Map symbol key**
- Strike and dip of bedding
 - Strike of vertical bedding
 - Strike and dip of overturned bedding
 - Strike and dip of foliation
 - Strike of vertical foliation
 - Strike and dip of joint
 - Strike of vertical joint
 - Strike and dip of vein
 - Strike of vertical vein
 - Geologic unit contacts
 - Fault (generally high angle)
 - - - Fault (interpreted)
 - Fault (inferred)

The State of Alaska makes no express or implied warranties (including warranties for merchantability and fitness) with respect to the character, functions, or capabilities of the electronic data or products or their appropriateness for any user's purposes. In no event will the State of Alaska be liable for any incidental, indirect, special, consequential or other damages suffered by the user or any other person or entity whether from the use of the electronic services or products, any failure thereof or otherwise, and in no event will the State of Alaska's liability to the Requestor or anyone else exceed the fee paid for the electronic service or product.

THIS MAP IS STRICTLY PRELIMINARY. IT HAS NOT BEEN REVIEWED FOR TECHNICAL CONTENT OR FOR CONFORMITY TO THE EDITORIAL STANDARDS OF DGGS. NOR HAS IT BEEN REVIEWED BY CO-AUTHORS.

- Affiliation:
- 1 Alaska Division of Geological & Geophysical Surveys
 - 2 U.S. Geological Survey
 - 3 University of Alaska Fairbanks
 - 4 Consultant



PRELIMINARY GEOLOGIC MAP OF THE IRON CREEK AREA, TALKEETNA MOUNTAINS B-5 QUADRANGLE, ALASKA

by
M.B. Werdon¹, J.R. Riehl², J.M. Schmidt², R.J. Newberry³, and G.H. Pessel⁴
2000

References Cited

Csejey, Bela, Jr., Nelson, W.H., Jones, D.L., Silberling, N.J., Dean, R.M., Morris, M.S., Lanphere, M.A., Smith, J.G., and Silberman, J.L., 1978. Reconnaissance geologic map and geochronology, Talkeetna Mountains quadrangle, northern part of Anchorage quadrangle, and southwest corner of Healy quadrangle, Alaska. U.S. Geological Survey Open-File Report 78-558-A, 1:250,000, 1 sheet, 60 p.

Staff, Alaska Division of Geological & Geophysical Surveys, Staff, Dighem, Inc., and Staff, W.G.M., Inc., 1998. CD-ROM containing profile and gridded data and section lines of the 1997 geophysical survey data for Iron Creek area, Talkeetna Mountains Quadrangle, southcentral Alaska. Alaska Division of Geological & Geophysical Surveys Public-Data File 98-9, 1 CD-ROM.

Smith, T.E., 1981. Geology of the Clearwater Mountains, south-central Alaska. Alaska Division of Geological & Geophysical Surveys Geologic Report 60, 72 p., 2 sheets and one 1:63,360-scale map sheet.



State of Alaska
Department of Natural Resources
Division of Geological & Geophysical Surveys