

STATE OF ALASKA  
DEPARTMENT OF NATURAL RESOURCES  
DIVISION OF GEOLOGICAL AND GEOPHYSICAL SURVEYS

Steve Cowper, *Governor*

Judith M. Brady, *Commissioner*

Robert B. Forbes, *Director and State Geologist*

March 1988

This report is a preliminary publication of DGGS.  
The author is solely responsible for its content and  
will appreciate candid comments on the accuracy of  
the data as well as suggestions to improve the report.

Report of Investigations 87-19  
CHEMICAL AND BIOLOGICAL WATER QUALITY OF  
SELECTED STREAMS IN THE BELUGA COAL AREA,  
ALASKA

By  
Mary A. Maurer

STATE OF ALASKA  
Department of Natural Resources  
DIVISION OF GEOLOGICAL & GEOPHYSICAL SURVEYS

According to Alaska Statute 41, the Alaska Division of Geological and Geophysical Surveys is charged with conducting 'geological and geophysical surveys to determine the potential of Alaskan land for production of metals, minerals, fuels, and geothermal resources; the locations and supplies of ground water and construction materials; the potential geologic hazards to buildings, roads, bridges, and other installations and structures; and shall conduct such other surveys and investigations as will advance knowledge of the geology of Alaska.'

In addition, the Division of Geological and Geophysical Surveys shall collect, record, evaluate, and distribute data on the quantity, quality, and location of underground, surface, and coastal water of the state; publish or have published data on the water of the state and require that the results and findings of surveys of water quality, quantity, and location be filed; require that water-well contractors file basic water and aquifer data, including but not limited to well location, estimated elevation, well-driller's logs, pumping tests, flow measurements, and water-quality determinations; accept and spend funds for the purposes of this section, AS 41.08.017 and 41.08.035, and enter into agreements with individuals, public or private agencies, communities, private industry, and state and federal agencies; collect, record, evaluate, archive, and distribute data on seismic events and engineering geology of the state; and identify and inform public officials and industry about potential seismic hazards that might affect development in the state.

Administrative functions are performed under the direction of the Director, who maintains his office in Fairbanks. The locations of DGGS offices are listed below:

.794 University Avenue  
(Suite 200)  
Fairbanks  
(907)474-7147

.400 Willoughby Avenue  
(3rd floor)  
Juneau  
(907)465-2533

.18225 Fish Hatchery Road  
P.O. Box 772116  
Eagle River  
(907)696-0070

This report is for sale by DGGS for \$2.50. DGGS publications may be inspected at the following locations. Mail orders and inquiries should be addressed to the Fairbanks office.

.794 University Avenue  
(Suite 200)  
Fairbanks, 99709  
  
.U.S. Geological Survey  
Public Information Office  
701 C Street  
Anchorage

.400 Willoughby Avenue  
(3rd floor)  
Juneau  
  
.Information Specialist  
U.S. Geological Survey  
4230 University Drive, Room 101  
Anchorage

# CONTENTS

	<u>Page</u>
Abstract.....	1
Introduction.....	2
Study area.....	2
Methods.....	4
Chemical.....	4
Biological.....	4
Results and discussion.....	6
Chemical water quality.....	6
Field variables.....	6
Dissolved constituents.....	7
Trace metals and minor elements.....	8
Nutrients.....	10
Biological water quality.....	12
Invertebrate abundance.....	12
Invertebrate composition.....	15
Benthic ecology.....	17
Conclusions.....	18
References cited.....	18
Appendix A - Field variables and major inorganic constituents of Beluga water-quality samples.....	21
Appendix B - Minor element analysis of Beluga water-quality samples...	23
Appendix C - Nutrient analysis of Beluga water-quality samples.....	27
Appendix D-1 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, June 15, 1983.....	28
Appendix D-2 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, June 15, 1983.....	30
Appendix D-3 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, August 25, 1983.....	32
Appendix D-4 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, August 25, 1983.....	34
Appendix D-5 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, June 14, 1984.....	36
Appendix D-6 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, June 14, 1984.....	38
Appendix D-7 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, August 29, 1984.....	40
Appendix D-8 - Density, number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, August 29, 1984.....	42

	<u>Page</u>
Appendix E - Three-factor analysis of variance table, where the variable is benthic-invertebrate abundance during June and August 1983 and 1984, Middle and Lone Creeks, Beluga coal area, Alaska.....	44
Appendix F - Wet-weight biomass of benthic invertebrates collected from sites in Middle and Lone Creeks, Beluga coal area, during June and August 1983 and 1984.....	45
Appendix G-1 - Habitat parameters at benthic-invertebrate sampling sites, June 15, 1983.....	46
Appendix G-2 - Habitat parameters at benthic-invertebrate sampling sites, August 25, 1983.....	47
Appendix G-3 - Habitat parameters at benthic-invertebrate sampling sites, June 14, 1984.....	48
Appendix G-4 - Habitat parameters at benthic-invertebrate sampling sites, August 29, 1984.....	49

## FIGURES

Figure 1. Location map of benthic invertebrate and chemical water-quality sampling sites, Beluga coal area, Alaska.....	3
2. Stream discharge on each sampling data at chemical water-quality sites in five streams, Beluga coal area, Alaska.....	7
3. Trilinear diagram of water analyses for five streams in the Beluga coal area, Alaska, during 1983 and 1984.....	8
4. Seasonal variation in total aluminum, total iron, and dissolved iron in five streams, Beluga coal area, Alaska, from 1982-84.....	9
5. Seasonal variation in nitrogen and phosphorus in five streams, Beluga coal area, Alaska, from 1982-84.....	11
6. Density, wet-weight biomass, mean number of taxa and diversity of Benthic-invertebrate communities at sites in Middle and Lone Creeks, Beluga coal area, Alaska, during 1983 and 1984.....	14
7. Composition percentages of benthic invertebrates at sites in Middle and Lone Creeks, Beluga coal area, Alaska, during 1983 and 1984.....	16

## TABLE

Table 1. Mean density, mean biomass, Shannon-Weaver diversity, evenness, and mean number of taxa of benthic invertebrates in Middle and Lone Creeks by month and year.....	13
of Benthic invertebrates in Middle and Lone Creeks by month and year for June and August of 1983 and 1984....	

CHEMICAL AND BIOLOGICAL WATER QUALITY OF SELECTED STREAMS IN THE BELUGA COAL  
AREA, ALASKA

by  
Mary A. Maurer<sup>1</sup>

ABSTRACT

Chemical water quality of five streams (Chuitna River and Bishop, Capps, Middle, and Lone Creeks) in the Beluga coalfield was determined during 1983-84 to evaluate premining conditions. In addition to field measurements, inorganic constituent, trace metal, minor element, and nutrient samples were collected. Benthic invertebrates were quantitatively sampled in two of the five streams, Middle and Lone Creeks, to assess biological water quality. Results showed that the five streams have good chemical water quality with high concentrations of dissolved oxygen and low concentrations of dissolved inorganic constituents, trace metals, minor elements, and nutrients. All streams have calcium-bicarbonate water, except lower Bishop Creek, which is a mixed type (calcium-sodium bicarbonate water). Low alkalinity values in all five streams indicate poor acid-neutralizing capability. Increased stream-flow, surface runoff, and suspended sediment elevate total metal and nutrient concentrations in Bishop and Capps Creeks during June. Total iron concentrations are relatively high in all five streams. Benthic-invertebrate community structure shows that biological water quality in Middle and Lone Creeks is good. Benthic invertebrate standing crop exceeds 12,000 invertebrates per m<sup>2</sup> and number of taxa averages 19 in both streams. Although invertebrate densities vary, the composition of the taxa is similar among the sites. Chironomid midges are the most abundant taxa in both streams. High invertebrate density and numerous taxa are attributed to warm summer water temperatures, light suspended sediment loads, and groundwater-maintained winter baseflow.

---

<sup>1</sup>DGGS, P.O. Box 772116, Eagle River, Alaska 99577.

## INTRODUCTION

Surface coal mining is proposed to begin during the 1990's in the Beluga coal area. Surface-water quality protection is a primary concern in the development of these proposed coal mines because of the highly valued fishery resources of the Chuitna and Beluga Rivers and their tributaries. Planning for protection of the surface waters and their fishery resources can be enhanced through areal collection of baseline surface-water quality data prior to mining. Several studies have investigated surface-water quality in the Beluga coal area (Scully, 1981; Environmental Research and Technology, Inc. [ERT], 1984a, 1984b; Maurer and Toland, 1984). The purpose of this study is to supplement these prior studies by interpreting the second year data of the chemical water quality study and to present biological water-quality information on two streams that will be influenced by the first phase of coal mine construction.

The specific objectives of the study are to: 1) determine baseline chemical water quality in five streams within the Beluga coal area, 2) assess biological water quality by determining the benthic invertebrate community in Middle and Lone Creeks prior to mining, and 3) supplement baseline information to assess the effects of future coal mining on water quality. The emphasis of the chemical water-quality investigation is on trends in field variables, major inorganic constituents, and nutrients. Samples were collected to correspond with specific hydrologic flow conditions of early summer (June), late summer (August), early winter (December), and late winter (March). The focus of the biological water-quality investigations is on determinations of benthic invertebrate distribution and abundance. Benthic invertebrates were selected as biological indicators of water quality because they are relatively immobile, year-round inhabitants of streams, are sensitive to water chemistry and aquatic habitat changes, and are important food sources for fish (Cairns and Dickson, 1971).

## STUDY AREA

The Beluga coal area is located in southcentral Alaska on the west side of Cook Inlet, about 80 km (50 mi) west of Anchorage (fig. 1). A detailed description of the physiography, climate, and stream characteristics is given in Maurer and Toland (1984). Five nonglacial streams, Bishop Creek, Capps Creek, Middle Creek, Lone Creek, and the Chuitna River, which drain from the Beluga coalfield area, were selected to obtain areal water-quality conditions. The locations of chemical and biological water-quality sampling sites along these streams are shown on figure 1. Bishop Creek is the proposed control stream because no coal mining is planned within its watershed. All chemical water-quality sampling sites were located in the lower reaches of the streams, downstream from prospective mining. Macroinvertebrate sampling sites were located at an upper, middle, and lower reach in Middle and Lone Creeks, which drain from a proposed surface coal mine (fig. 1).




Figure 1. Location map of benthic invertebrate and chemical water-quality sampling sites, Beluga coal area, Alaska.

## METHODS

### Chemical

Stream discharge was measured on each sampling date at the chemical water-quality sites with a Marsh-McBirney current meter according to U.S. Geological Survey methods (Carter and Davidian, 1968; Buchanan and Somers, 1969). Water temperature, dissolved oxygen concentration, and specific conductance were measured in the field with a digital 4041 Hydrolab. An Orion digital pH meter was used to measure field pH. Measurements of dissolved oxygen and pH were taken in low velocity reaches within the stream to avoid streaming effects across the membrane probes. Bicarbonate alkalinity was measured in the field by titrating an untreated 200-ml sample with 0.01639N sulfuric acid to an electrometrically determined endpoint of pH 4.5 (U.S. Environmental Protection Agency [EPA], 1983).

All water samples were collected by the grab sampling technique. Samples for major inorganic constituents and dissolved trace-metal analyses were immediately filtered through a 0.45- $\mu$ m membrane filter. Total and dissolved trace-metal samples were acidified with double-distilled 70-percent nitric acid immediately after collection. Nutrient samples for total concentration analysis were untreated, while those for dissolved concentration analysis were filtered in the field through a 0.45- $\mu$ m membrane filter. All nutrient samples were frozen within eight hours of collection.

Major inorganic constituents, total trace-metal, and dissolved trace-metal samples were analyzed by Anatec Laboratories, Inc., Santa Rosa, California. All inorganic constituents were analyzed according to U.S. EPA (1983) or American Public Health Association (APHA) methods (1980). With the exception of boron, trace-metal concentrations were measured using atomic absorption spectrophotometry according to the methods of the U.S. EPA (1983). Boron concentrations were measured colorimetrically using the methods of Wolf (1974). Nutrient samples were analyzed at the Alaska Department of Fish and Game laboratory in Soldotna, Alaska. Concentrations of total phosphorus and total ammonia plus organic nitrogen were analyzed with a Technicon Auto Analyzer II. Dissolved nitrite plus nitrate and dissolved ammonia were analyzed in accordance with the methods of Stainton and others (1977). Filterable reactive phosphorus (an estimate of orthophosphate), and total filterable phosphorus were measured according to the methods of Eisenreich and others (1975).

### Biological

Three benthic invertebrate sampling sites were selected on the upper, middle, and lower reaches of Middle and Lone Creeks corresponding to synoptic survey sites 1, 2, 5, 7, 8, and 11 (fig. 1) of Maurer and Toland (1984). Sites 5 and 11 are located at chemical water quality sites (fig. 1). Samples were collected during June and August of 1983 and 1984. Statistical analyses were performed on synoptic-survey invertebrate densities using the methods of Elliott (1971) to estimate a suitable sample size per site. The results indicated that ten samples per site in Middle Creek and three samples per site in Lone Creek were required for a standard error equal to 20 percent of the



mean. Because such a sampling schedule would greatly increase the time required for sampling and analysis, an alternative of four samples per site was agreed upon. Each stream reach was separated into four equal vertical strata. A stratified random sampling technique was used; that is, one sample was randomly chosen within each stratum. Habitat variables of water depth, stream width, and water temperature were measured at each sampling point. Water velocity at the streambed was measured with a Marsh-McBirney current meter before sampling. Stream-substrate composition within the area of the samples was estimated visually by examining the relative percentages of boulder (>256 mm diam), rubble (64-256 mm diam), gravel (2-64 mm diam), and sand/silt (0.004-2.0 mm diam) (U.S. EPA, 1973).

A cylindrical aluminum substrate sampler 0.6 m high and 0.1 m<sup>2</sup> wide was used to collect benthic invertebrates. The front side of the sampler frame was covered with a net composed of 600-μm (pore diameter) NITEX (nylon mesh) to increase water flow through the sampler; 300-μm NITEX was used on the back side and trailing collection bag. Samples were collected by working the sampler into the streambed and displacing the rocks to dislodge invertebrates. Larger rocks were examined to insure that all invertebrates were removed. Invertebrates were washed into the collection bag and trapped in a detachable plastic bucket at the end of the bag. Samples were preserved with a solution of 70-percent ethyl alcohol and water, to which rose bengal bacteriological stain was added to facilitate later sorting.

All invertebrates were handpicked from sample debris and stored in 70-percent ethyl alcohol. Insects were counted and identified to the most practical taxonomic level using keys by Usinger (1956), Jensen (1966), Smith (1968), Edmunds and others (1976), Baumann and others (1977), Wiggins (1977), and Merritt and Cummins (1978). In many cases, very small specimens could be identified only to the ordinal or family taxonomic level. Non-insect invertebrates were identified to the class or ordinal level using keys published by Pennak (1978).

Invertebrate biomass was determined by measuring the wet weight ( $\pm 0.001$ g) of all invertebrates in each benthic sample. Invertebrates, the alcohol contents of the vial, and a 10-ml alcohol rinse were poured onto a tared 0.45-μm membrane filter contained in a Millipore filtering unit. A vacuum-pump was hand-operated at a pressure of 30-cm mercury for one minute to remove the excess alcohol. The invertebrates and filter were then immediately weighed on an electronic balance.

Several quantitative methods were used to analyze invertebrate samples. Insect abundance was based on density (number of invertebrates per m<sup>2</sup>). The number of taxa in each sample was determined by summing the identifiable insect families and other invertebrate groups. Invertebrate community structure was calculated using the Shannon-Weaver diversity index ( $H'$ ), a measure of the number and relative abundance of taxa in a sample, and the evenness value ( $J'$ ), a measure of the distribution of individuals among taxa in a sample (Poole, 1974). The formula for the Shannon-Weaver diversity index is  $H' = - \sum_{i=1}^s p_i \log_2 p_i$ , where  $s$  is the number of taxa and  $p_i$  is a proportion

(total number of invertebrates of the  $i^{\text{th}}$  taxa divided by the total number of invertebrates of all taxa). Evenness is expressed as  $J' = H'/H'_{\text{maximum}}$  where  $H'_{\text{maximum}} = \log_2 s$  ( $s$  = number of taxa). The diversity and evenness values for stream, year, and month were calculated on pooled samples; that is, all samples within the stream, year, or month were summed to form a single sample.

A three-factor statistical analysis of variance (Zar, 1974) was applied to invertebrate density data to determine if there were differences between streams and among sites. Prior to analysis, the density data in each sample were transformed from  $X$  to  $\log X$  to approximate a normal distribution (Elliott, 1971). The probability level used in the analysis of variance  $F$  test was = 0.05.

## RESULTS AND DISCUSSION

### Chemical Water Quality

#### Field variables

Streamflow was measured at each chemical water-quality site on each sampling date (fig. 2). The hydrographs show high flow during June in Bishop and Capps Creeks, and the Chuitna River, but little variation in streamflow in Middle and Lone Creeks. Although an attempt was made to collect data during the various flow conditions, including winter baseflow, spring runoff, summer low flow, and early winter flow, the single sampling date in June did not allow peak spring runoff to be measured in Middle and Lone Creeks. Peak spring discharge at these two streams occurs in May (ERT, 1984b). Moreover, the June measurement of suspended sediment load in Middle and Lone Creeks is significantly less than in Bishop and Capps Creeks (Maurer and Toland, 1984). These observations are important because suspended sediment has a significant effect on the chemical water quality of Bishop and Capps Creeks. Streamflow measured in the Chuitna River in June represents early summer high flow because peak spring runoff normally occurs in late May or early June (USGS, 1984; 1985).

Other variables measured in the field were water temperature, specific conductance, pH, alkalinity, and dissolved oxygen concentration (app. A). Water temperature ranged from 0°C in Bishop, Middle, and Lone Creeks, and the Chuitna River during December and March to a high of 13.8°C in the Chuitna River in August (app. A). Capps Creek showed the least variation in water temperature due to higher elevation, relatively steep gradient, and proximity to ground water sources. Specific conductance is relatively low for all the streams compared to surface waters elsewhere (Hem, 1970), averaging only 50  $\mu\text{mhos/cm}$  at the five sampling sites. Although relatively little change in specific conductance occurred seasonally among the five streams, specific conductance did vary inversely with discharge. Mean pH values show that Bishop and Capps Creeks are slightly acid, while Middle and Lone Creeks and the Chuitna River have values slightly above neutrality. The lowest pH, 5.85, was measured in Capps Creek. Bicarbonate alkalinity is similar among all sites and varied inversely with discharge. Alkalinity values, ranging from 10.5 to 46 mg/L, indicate poor ability of the streams to neutralize acids. Dissolved oxygen concentrations were generally near saturation in each stream,




Figure 2. Stream discharge on each sampling data at chemical water-quality sites in five streams, Beluga coal area, Alaska.

ranging from 92 to 100 percent saturation in the summer and from 78 to 100 percent in the winter. The lowest concentrations were measured in December 1982.

#### Dissolved constituents

The concentration of major cations and anions was consistently low. The total filterable residue (dissolved solids) concentrations varied little, ranging from 44 to 61 mg/L among the five streams (app. A). Based on the average percentage of major ion concentrations, Bishop Creek had a slightly different ionic composition than the other four streams (fig. 3).

Between 50 and 55 percent of the cations in Capps, Middle, and Lone Creeks and the Chuitna River are calcium, followed by magnesium, sodium, and potassium at 24, 22, and 4 percent, respectively. Bishop Creek, however, has equal percentages (41 percent) of calcium and sodium, a relatively low 15 percent of magnesium, and 4 percent potassium.

Bicarbonate represents about 86 percent of the anions in all five streams. Chloride and sulfate ions are relatively minor (less than 10 percent each) for all streams except Bishop Creek, where chloride is 21 percent of the total anions (fig. 3).

Based on these ionic compositions, Capps, Middle, and Lone Creeks and the Chuitna River have been classified as calcium-bicarbonate waters, while Bishop




Figure 3. Trilinear diagram of water analyses for five streams in the Beluga coal area, Alaska, during 1983 and 1984.

Creek has been classified as calcium-sodium bicarbonate water because of higher sodium and chloride content (fig. 3). The ionic composition of the middle reach of Bishop Creek does not differ appreciably from the other four streams (Scully, 1981), and the source of the sodium and chloride ions may be exposed deposits of 'very fine bonded plastic clay' which occur only along the stream's lower reach (Barnes, 1966).

Silica concentrations ranged from 9.7 to 13.6 mg/L (app. A), which are characteristic of surface waters (Hem, 1970). Significantly lower concentrations, however, were measured in August at all sites, and this may be in part due to silica utilization by aquatic algae, particularly diatoms (Reid, 1976).

#### Trace metals and minor elements

The concentrations of trace metals and minor elements measured in all five streams are generally low or below detection limits (app. B), and most elements do not vary significantly among streams nor show distinct seasonal trends. Aluminum and iron were the most abundant metals measured in all five streams, and seasonal trends are apparent in total aluminum and dissolved iron concentrations (fig. 4). Total aluminum concentrations generally were similar among streams, but concentrations were noticeably elevated in Bishop and Capps Creeks in June (2.2 mg/L and 12.0 mg/L, respectively) due to high suspended sediment load.




Figure 4. Seasonal variation in total aluminum, total iron, and dissolved iron in five streams, Beluga coal area, Alaska, from 1982-84.

Total iron concentrations varied little among streams (fig. 4), but were consistently the highest of all trace metals measured, ranging from 0.41 to 8.8 mg/L (app. B). There was little seasonal variation in total concentrations among streams, except at the Capps Creek site where a concentration of 8.8 mg/L was measured during the high suspended-sediment load period in June (fig. 4). Dissolved iron concentrations were also similar among streams, ranging from 0.02 to 1.2 mg/L (app. A), and a seasonal variation is apparent. The highest dissolved iron concentrations were measured in winter (December and March); the lowest were measured in August (fig. 4). This pattern, seen in all five streams, may be due to the accumulation of organic matter and bacteria and algal growth which facilitates the precipitation of ferric hydroxide on the stream bottom (Reid, 1976), thereby reducing the concentration of dissolved iron in August.

The Chuitna River site consistently had the least seasonal variation and lowest concentration of total and dissolved iron of all sites measured (fig. 4). Total and dissolved iron concentrations at this site averaged 0.63 mg/L and 0.26 mg/L, respectively. Although total iron concentrations in Bishop, Capps, Middle, and Lone Creeks frequently exceeded the U.S. EPA criteria for protection of fresh-water aquatic life, that is, 1.0 mg/L (EPA, 1976), dissolved iron concentrations averaged less than 1.0 mg/L in all streams (fig. 4).

Total zinc concentrations were detectable in low concentrations (<10 µg/L) in all streams, and the highest total concentration was recorded in Capps Creek, during June, at 78 µg/L. Barium and strontium were measured in low concentrations in all streams. Low concentrations of these elements are typical of many surface waters (Hem, 1970). Total manganese concentrations were detectable, but were relatively low in all streams, ranging from 0.02 mg/L to 0.28 mg/L. The highest total manganese concentrations were associated with suspended sediment in Capps Creek, and the lowest concentrations occurred in the Chuitna River, where the mean concentration was <0.03 mg/L (app. B).

### Nutrients

The concentration of dissolved nitrite plus nitrate nitrogen ranged from 0.012 to 0.541 mg/L in the five streams (app. C), being relatively high in December and March and low in August in all streams (fig. 5). The concentrations measured in December and March may be due to ground water inflow under base flow conditions. Low concentrations in August may be the result of nitrate utilization by algae and bacteria (Reid, 1976). Elevated concentrations in Bishop and Capps Creeks, 0.541 and 0.475 mg/L respectively, probably result from surface runoff associated with high streamflows in June.

Total ammonia plus organic nitrogen concentrations were similar in the five streams (fig. 5), and no seasonal trend was observed. A relatively high concentration of 0.26 mg/L was measured in Capps Creek during the June high streamflow (fig. 5). The increasing concentrations measured in all five streams from March through August 1983 is probably due to organic loading from surface runoff and to periphyton production (Reid, 1976).




Figure 5. Seasonal variation in nitrogen and phosphorus in five streams, Beluga coal area, Alaska, from 1982-84.

Dissolved ammonia nitrogen concentrations were relatively low in all five streams, ranging from 0.01 to 0.04 mg/L (app. C). The highest concentrations, 0.136 and 0.033 mg/L in Bishop Creek and Capps Creek, respectively, were measured during the increased surface runoff of June.

The measured phosphorus fractions consisted of total phosphorus, total filterable phosphorus, and filterable reactive phosphorus. The latter two correspond to dissolved phosphorus and orthophosphates, respectively (APHA, 1980). Orthophosphate is the form of phosphorus utilized by plants. Total phosphorus concentrations were similar in the five streams, ranging from 0.010 to 0.140 mg/L (fig. 5). Elevated concentrations were measured in Capps Creek in June and August (fig. 5). Although total phosphorus concentrations were not measured in Bishop Creek in June, the elevated concentration in August suggests that June concentrations in Bishop Creek were probably elevated as well. Total filterable and filterable reactive phosphorus concentrations were also similar among streams and exhibited little seasonal variation, except in Bishop and Capps Creeks (fig. 5). Total filterable phosphorus concentrations ranged from 0.005 to 0.088 mg/L and filterable reactive phosphorus concentrations ranged from 0.004 to 0.044 mg/L (app. C). The percentage of total filterable relative to total phosphorus was consistently high throughout the sampling period, excluding elevated concentrations in Bishop and Capps Creeks during June. It is therefore inferred that phosphorus is primarily in the dissolved form rather than the particulate form. Similarly, the percentage of filterable reactive to total filterable phosphorus was high in all five streams (app. C), indicating that the majority of the dissolved phosphorus is in the form of orthophosphates. These consistent concentrations of all three phosphorus fractions are probably the result of a large ground water contribution to streamflow, which, for example, is 34 percent in Lone Creek and 32 percent in Middle Creek (ERT, 1984c).

High June discharge conditions of Bishop and Capps Creeks are the probable cause of elevated concentrations of all three phosphorus fractions. However, elevated concentrations of all three fractions were measured in Bishop Creek during August under relatively low streamflow conditions. Although such elevated phosphorus concentrations cannot adequately be explained by the limited data of this study, they are most likely the result of biological processes.

## Biological Water Quality

### Invertebrate abundance

Invertebrate mean density, calculated as the number of organisms per m<sup>2</sup>, varied by less than 24 percent between Middle and Lone Creeks (table 1). The mean density was 12,085 invertebrates per m<sup>2</sup> in Middle Creek and 15,806 invertebrates per m<sup>2</sup> in Lone Creek. The mean density was approximately 26 percent higher in 1983 than in 1984 in both streams. Although June and August mean invertebrate densities were virtually constant in Middle Creek, the June density was two times greater than the August density in Lone Creek (table 1). Density decreased progressively from the upper (headwater) site to the lower site in Lone Creek (fig. 6). The pattern of invertebrate density differed somewhat in Middle Creek, with relatively high density at the upper and middle site and low density at the lower site (fig. 6).



Table 1. Mean density (no./m<sup>2</sup>), mean biomass (g/m<sup>2</sup>), Shannon-Weaver diversity, evenness, and mean number of taxa of benthic invertebrates in Middle and Lone Creeks by month and year.  $\bar{n}$  = number of samples. A 95-percent confidence interval is shown for each mean value of density and biomass. Diversity and evenness values were calculated on the basis of pooled samples.

	<u>Middle Creek</u>	<u>Lone Creek</u>
Density (no./m <sup>2</sup> )		
overall ( $\bar{n}$ = 48)	12085 ± 2389	15806 ± 4032
month ( $\bar{n}$ = 24)		
June	12330 ± 4269	21055 ± 7202
August	11841 ± 2784	10557 ± 3454
year ( $\bar{n}$ = 24)		
1983	14008 ± 4470	17944 ± 7516
1984	10162 ± 2152	13668 ± 3983
Biomass (m/m <sup>2</sup> )		
overall ( $\bar{n}$ = 48)	8.77 ± 0.91	13.40 ± 2.17
month ( $\bar{n}$ = 24)		
June	9.62 ± 1.59	17.50 ± 3.47
August	7.92 ± 0.98	9.30 ± 1.77
year ( $\bar{n}$ = 24)		
1983	9.00 ± 1.49	14.67 ± 3.74
1984	8.54 ± 1.23	12.13 ± 2.63
Diversity (H')		
overall ( $\bar{n}$ = 48)	2.88	2.57
month ( $\bar{n}$ = 24)		
June	2.13	1.71
August	2.99	3.26
year ( $\bar{n}$ = 24)		
1983	2.79	2.44
1984	2.83	2.67
Evenness (J')		
overall ( $\bar{n}$ = 48)	0.58	0.52
month ( $\bar{n}$ = 24)		
June	0.44	0.36
August	0.61	0.66
year ( $\bar{n}$ = 24)		
1983	0.56	0.50
1984	0.57	0.54
Number of Taxa		
overall ( $\bar{n}$ = 48)	19	19
month ( $\bar{n}$ = 24)		
June	17	17
August	22	22
year ( $\bar{n}$ = 24)		
1983	20	19
1984	19	20




Figure 6. Density, wet-weight biomass, mean number of taxa and diversity of benthic-invertebrate communities at sites in Middle and Lone Creeks, Beluga coal area, Alaska, during 1983 and 1984. U = upper site, M = middle site, and L = lower site. Sites 1, 2, and 5 in Middle Creek and sites 7, 8, and 11 in Lone Creek correspond to the upper, middle, and lower site in their respective stream.  $\underline{n}$  = number of samples.

The invertebrate densities of table 1 are substantially higher than those found by Scully (1981) and ERT (1984a). These differences are probably the result of different sampling methodologies. Artificial substrates and a dip net technique was used in the Scully study and a Surber sampler was used in the ERT study. The completely enclosed substrate sampler and finer net mesh size (300- $\mu$ m) used in our study resulted in higher densities. In addition, densities reported here are considerably higher than those in Maurer and Toland (1984), who sampled many microhabitats including runs, pools, and riffles. The only habitats sampled in our study were riffles and runs, which typically have higher invertebrate densities than pools (Hynes, 1970). Despite density differences between the three studies, the mean number of taxa collected was similar.

A three-factor analysis of variance (ANOVA, Zar, 1974) was performed on the invertebrate density data to determine if differences among streams (Middle vs. Lone), sites, (upper, middle, lower), and dates (June 1983, August 1983, June 1984, August 1984) were statistically significant. The results of the ANOVA analysis indicated no significant difference between streams, a highly significant difference among sites, and a significant difference among dates (app. E). This statistical test substantiates the relationships summarized on table 1 and figure 6. The only significant interaction was between stream and site; that is, mean invertebrate density of a stream was dependent on site (app. E). The interaction of stream and date approached statistical significance (calculated F value = 2.74 versus critical F value = 2.76), due to high June densities in Lone Creek.

Diversity and evenness values were slightly higher in Middle Creek. The overall diversity values, calculated from pooled samples, were 2.88 in Middle Creek and 2.57 in Lone Creek (table 1). The low to moderate numerical values in both streams indicate a fairly uneven distribution of taxa in samples (app. D). There was no difference between years but values were higher in August, due to an increase in the number of taxa. The number of taxa ranged from 11 to 27, and averaged 19 in both streams. Generally, the highest number of taxa occurred at the upper sites in both streams. The lowest number of taxa occurred at the middle site in Lone Creek, but there was no distinct trend in Middle Creek.

Invertebrate biomass was higher in Lone Creek, averaging 13.40 g/m<sup>2</sup> in Lone Creek and 8.77 g/m<sup>2</sup> in Middle Creek (table 1). Biomass in Middle Creek did not vary appreciably between the June and August sampling periods. Biomass in Lone Creek, however, was greater in June than in August by an average of 8.20 g/m<sup>2</sup> (table 1). The June increase resulted from higher biomass at the upper site (site 7) and the middle site (site 8) (app. F). Generally, there was less variability in biomass than in density (fig. 6).

#### Invertebrate composition

Five insect orders and six major groups of non-insect invertebrates were found at all sites. Diptera (true flies), predominantly chironomid midges and blackflies, were the most abundant invertebrates and represented 66 percent of the total invertebrate composition in Middle Creek and 73 percent in Lone Creek (fig. 7). Moreover, Diptera represented 80 percent of the total




Figure 7. Composition percentages (based on density) of benthic invertebrates at sites in Middle and Lone Creeks, Beluga coal area, Alaska, during 1983 and 1984.  $n$  = number of samples.

composition in June, but only 50 percent in August. As a result, the percentages of the other invertebrate groups were two to three times greater in August, the greatest increase occurring in Plecoptera. These increases are due to the appearance of early instars of nemourid and capniid stoneflies and heptageniid mayflies. The decrease in the number of Diptera is probably because of pupation and emergence of midges and blackflies during the summer.

Ephemeroptera (mayflies) was the second most abundant invertebrate group, averaging approximately 12 percent of a sample in both streams. Non-insect invertebrates constituted approximately 8 percent of the total invertebrates; Oligochaeta (aquatic earthworms) and Acarina (aquatic mites) were the most abundant taxa. Plecoptera (stoneflies) and Trichoptera (caddisflies) constituted approximately 6 percent of the total; capniid and nemourid stoneflies and glossosomatid caddisflies were the most abundant taxa in their respective groups (app. D).

The relative distribution of invertebrate taxa, based on invertebrate density by percentages, was similar among sites in Middle and Lone Creeks (fig. 7). Taxa were more evenly distributed at the lower sites of each stream, that is, site 5 in Middle Creek and site 11 in Lone Creek, because there were fewer dipteran flies present. As a result, diversity and evenness values were higher at these two sites (app. D).

## Benthic ecology

Invertebrate community structure--invertebrate abundance and composition--is the result of the inherent physical, chemical, and biological conditions in Middle and Lone Creeks. The most important physical factor is climate, which affects the aquatic-riparian habitat to such an extent that invertebrate density and taxa are characteristic of temperate, rather than subarctic streams. The climate is moderated by Cook Inlet and both streams have a southern aspect and low gradient which raises water temperature above 13°C during the summer. Streamflow is relatively stable because the ground water contribution to streamflow exceeds 30 percent in both stream (ERT, 1984c). Therefore, stream substrates probably do not freeze during winter, and, because of stable streambanks, low stream gradient, and stony substrates, erosional processes are not significant.

The chemical water quality of these streams is good. Dissolved oxygen concentrations are consistently high. Suspended sediment, trace metal, and dissolved solid concentrations are quite low, and no concentrations are high enough to inhibit the invertebrate community.

There is also an abundant and varied food supply in Middle and Lone Creeks. The majority of invertebrates present in these streams, especially mayflies and midges, feed on periphyton and organic detritus (Merritt and Cummins, 1978). Blackflies strain fine particulate organic matter from the water column, and several stonefly taxa shred coarse organic matter such as leaves and grasses. Although most caddisflies collect detritus, limnephilid caddisflies were observed scavenging salmon carcasses on the streambed. Thus, these taxa fill more than one trophic level within the invertebrate community.

Similarity in physical factors, water chemistry characteristics, and aquatic-riparian habitat in Middle and Lone Creeks produce comparable invertebrate communities. There were, however, several major habitat differences among sites (app. G). Habitats at the lower sites in both streams consisted of runs with large substrate sizes, and shading from mixed conifer-deciduous canopy. The upper sites and middle site on Lone Creek had similar habitat features: a riffle with rubble-gravel substrate and shrub-grass riparian vegetation. The middle site in Middle Creek was different from all other sites in that it had very shallow riffles, small gravel-size substrate, and riparian vegetation consisting entirely of grasses. Although only minor differences in invertebrate community structure occurred among sites, relatively higher invertebrate densities and taxa numbers at upper sites may be a result of stable ground water flow and optimal substrate size for invertebrate colonization.

Invertebrate abundance and composition are appropriate variables for determining the biological water quality of these streams. Both streams have taxa typical of unpolluted, cold-water streams with stony substrates (Hynes, 1974). Invertebrate density is relatively high but variable because substrate size is variable among sites. These relative high densities, with moderate biomass and numerous taxa, indicate a highly productive benthic invertebrate community.

## CONCLUSIONS

Chemical water quality is good and very similar in Bishop, Capps, Middle, and Lone Creeks, and the Chuitna River. These streams have high concentrations of oxygen and low concentrations of dissolved solids, trace metals, and nutrients. Lower Bishop Creek has a slightly different ionic composition than the other four streams, due to higher sodium and chloride ion concentrations. The elevated concentrations of trace metals and nutrients during June in Bishop and Capps Creeks are the result of high streamflow, surface runoff, and suspended sediment load.

Biological water quality is good in Middle and Lone Creeks. The benthic invertebrate community is characterized by relatively high density, moderate biomass, and numerous taxa. The representative taxa are typically found in well-oxygenated, clear-water streams. Invertebrate composition is dominated by chironomid midges and blackflies. Although aquatic habitat differences produce invertebrate density differences among sites, the invertebrate community structure is similar between streams.

## REFERENCES CITED

- American Public Health Association, American Water Works Association, and Water Pollution Control Federation, 1980, Standard methods for the examination of water and wastes (15th ed.): Washington, D.C., American Public Health Association, 1134 p.
- Barnes, F.F., 1966, Geology and coal resources of the Beluga-Yentna region, Alaska: U.S. Geological Survey Bulletin 1202-C, p. 1-54.
- Baumann, R.W., Gaufin, A.R., and Surdick, R.F., 1977, The stoneflies (Plecoptera) of the Rocky Mountains: Philadelphia, Pennsylvania, Memoirs of the American Entomological Society, no. 31, 208 p.
- Buchanan, T.J., and Somers, W.P., 1969, Discharge measurements at gaging stations: U.S. Geological Survey Techniques of Water-Resources Investigations, book 3, chapter A8, 65 p.
- Cairns, John Jr., and Dickson, Kenneth L., 1971, A simple method for the biological assessment of the effects of waste discharges on aquatic bottom-dwelling organisms: Journal of the Water Pollution Control Federation, v. 43, no. 5, pp. 772-775.
- Carter, R.W., and Davidian, Jacob, 1968, General procedures for gaging streams: U.S. Geological Survey Techniques of Water-Resources Investigations, book 3, chapter A6, 13 p.
- Edmunds, G.F. Jr, Jensen, S.L., and Berner, L., 1976, The mayflies of North and Central America: Minneapolis, University of Minnesota Press, 330 p.
- Eisenreich, S.J., Bannerman, R.T., and Armstrong, D.E., 1975, A simplified phosphorus analysis technique: Environmental Letters, v. 9, no. 1, p. 43-53.
- Elliott, J.M., 1971, Some methods for the statistical analysis of samples of benthic invertebrates: Freshwater Biol. Assoc. Sci. Pub. 25, 144 p.
- Environmental Research and Technology, Inc., 1984a, Diamond Chuitna project aquatic biology baseline studies report: Fort Collins, Colorado, Environmental Research and Technology, 2 v.

- Environmental Research and Technology, Inc., 1984b, Diamond Chuitna project surface water hydrology and water quality baseline studies report: Fort Collins, Colorado, Environmental Research and Technology, 4 v.
- \_\_\_\_\_ 1984c, Diamond Chuitna project groundwater hydrology baseline studies report: Fort Collins, Colorado, Environmental Research and Technology, 1 v.
- Hem, John D., 1970, Study and interpretation of the chemical characteristics of natural water: U.S. Geological Survey Water-Supply Paper 1473, 363 p.
- Hynes, H.B.N., 1970, The ecology of running waters: University of Toronto Press, 555 p.
- Hynes, H.B.N., 1974, The biology of polluted waters: University of Toronto Press, 202 p.
- Jensen, S.L., 1966, The mayflies of Idaho: Salt Lake City, University of Utah, unpublished M.S. thesis, 367 p.
- Maurer, M.A. and Toland, D.C., 1984, Water-quality data from the Beluga coal-field area, Alaska, June 1982 through March 1983: Alaska Division of Geological and Geophysical Surveys Report of Investigation 84-27, 33 p.
- Merritt, Richard W. and Cummins, Kenneth W., 1978, An introduction to the aquatic insects of North America: Iowa, Kendall/Hunt Publishing Company, 441 p.
- Pennak, R.W., 1978, Freshwater invertebrates of the United States, 2nd Edition: New York, The Ronald Press Company, 803 p.
- Poole, R.W., 1974, An introduction to quantitative ecology: New York, McGraw-Hill Book Company, Inc., 532 p.
- Reid, George K. and Wood, Richard D., 1976, Ecology of inland waters and estuaries: New York, D. Van Nostrand Company, 485 p.
- Scully, D.R., Krumhardt, A.P., and Kernodle, D.R., 1981, Hydrologic reconnaissance of the Beluga, Peters Creek, and Healy coal area, Alaska: U.S. Geological Survey Water-Resources Investigations 81-56, 71 p.
- Smith, S.D., 1968, The *Rhyacophila* of the Salmon River drainage of Idaho with special reference to larvae: Annals of the Entomological Society of America, v. 61, no. 3, p. 655-674.
- Stainton, M.P., Capel, M.J., and Armstrong, E.J., 1977, The chemical analysis of freshwater: Winnipeg, Manitoba, Freshwater Institute, Fisheries and Marine Service, miscellaneous special publication no. 25, second edition, 180 p.
- U.S. Environmental Protection Agency, 1973, Biological field and laboratory methods for measuring the quality of surface waters and effluents: U.S. Environmental Protection Agency, EPA-670/4-73-001.
- \_\_\_\_\_ 1976, Quality criteria for water: U.S. Environmental Protection Agency, EPA-440/9-76-023.
- \_\_\_\_\_ 1983, Methods for chemical analysis of water and wastes: U.S. Environmental Protection Agency, EPA-600/4-79-020.
- U.S. Geological Survey, 1984, Water resources data for Alaska, water year 1983: U.S. Geological Survey Water-Data Report AK-83-1, 364 p.
- \_\_\_\_\_ 1985, Water resources data for Alaska, water year 1984: U.S. Geological Survey Water-Data Report AK-84-1, 358 p.
- Usinger, R.L., ed., 1956, Aquatic insects of California: Berkeley, University of California Press, 508 p.
- Wiggins, G.B., 1977, Larvae of the North American caddisfly genera (Trichoptera): University of Toronto Press, 401 p.

- Wolf, B., 1974, The determination of boron in soil extracts, plant materials, composts, manures, water and nutrient solutions: Soil Science Plant Analysis, v. 2, no. 5, p. 363-374.
- Zar, J.H., 1974, Biostatistical analysis: New Jersey, Prentice-Hall, Inc., 620 p.



Appendix A. Field variables and major inorganic constituents of Beluga water-quality samples.

Time	Streamflow, instantaneous (cfs)	Specific conductance (umhos at 25°C)	pH (units)	Water temperature (°C)	Silica, dissolved SiO <sub>2</sub> (mg/l)	Oxygen, dissolved (mg/l)	Oxygen, dissolved (percent saturation)	Calcium, dissolved (mg/l as Ca)	Magnesium, dissolved (mg/l as Mg)
Bishop Creek									
12-15-82 0935	15	33	7.30	-0.1	7.5	11.3	78	4.4	1.0
03-30-83 1230	13	61	6.85	-0.1	14	12.2	86	5.9	1.3
06-14-83 0910	108	24	6.80	10.5	8.1	10.2	92	2.4	0.58
08-24-83 0940	28	60	7.25	10.1	1.4	11.8	100	5.5	1.3
12-07-83 1115	<sup>a</sup> 23	50	6.55	-0.2	14	15.2	100	4.6	0.95
03-20-84 0930	35	45	6.75	-0.1	13	13.3	91	4.4	0.94
Capps Creek									
12-15-82 1225	8.8	48	6.90	0.5	6.9	12.4	88	4.9	1.2
03-30-83 1420	5.2	57	7.25	1.1	15	13.8	99	7.3	1.7
06-14-83 1110	157	10	5.85	5.0	7.1	12.0	96	2.1	0.44
08-24-83 1105	16	39	6.95	8.2	1.4	12.5	100	5.6	1.2
12-07-83 1250	<sup>b</sup> 7.0	52	6.65	-0.4	13	15.8	100	6.0	1.4
03-20-84 1100	<sup>b</sup> 12	53	7.05	0.1	15	15.2	100	6.3	1.5
Middle Creek									
12-16-82 1225	6.9	59	7.60	0	13	12.7	88	5.0	1.7
03-31-83 1410	4.3	77	6.85	0.2	19	14.2	100	8.9	2.2
06-14-83 1350	13	51	7.35	12.3	12	9.9	94	5.1	1.4
08-24-83 1220	9.7	64	6.95	10.1	1.6	12.0	100	8.0	1.9
12-07-83 1535	9.4	55	6.85	-0.3	14	15.9	100	6.4	1.7
03-20-84 1230	13	46	6.85	-0.3	15	15.4	100	5.4	1.4
Lone Creek									
12-16-82 1100	16	58	7.20	-1.0	12	12.9	89	4.1	1.4
03-31-83 1150	9.6	66	7.10	0	16	14.1	99	7.8	1.8
06-14-83 1515	28	47	7.35	13.7	6.1	9.5	92	5.2	1.3
08-24-83 1320	20	65	7.25	11.1	0.95	11.6	100	8.0	1.7
12-07-83 1445	13	59	6.85	-0.2	16	16.3	100	6.6	1.5
03-20-84 1350	14	52	6.90	-0.3	16	15.6	100	5.7	1.4
Chuitna River									
12-16-82 0900	<sup>c</sup> 100	42	7.00	-1.0	26	12.5	86	2.9	1.1
03-31-83 1100	<sup>c</sup> 120	57	7.20	0.1	16	14.3	100	8.1	2.2
06-14-83 1700	<sup>c</sup> 470	21	7.30	11.3	7.7	10.9	100	2.5	0.66
08-24-83 1445	139	47	8.10	13.8	2.1	11.6	100	7.0	1.6
12-07-83 1400	<sup>d</sup> 100	44	6.15	-0.3	16	15.9	100	5.8	1.5
03-20-84 1450	<sup>d</sup> 100	43	7.10	-0.3	14	15.6	100	5.2	1.5

<sup>a</sup>Estimate only, ice on probe head.<sup>b</sup>U.S. Geological Survey (1985, p. 180).<sup>c</sup>U.S. Geological Survey (1984, p. 159).<sup>d</sup>U.S. Geological Survey (1985, p. 182).

## Appendix A. (con.)

	Sodium, dissolved (mg/l as Na)	Potassium, dissolved (mg/l as K)	Iron, dissolved (mg/l as Fe)	Manganese, dissolved (mg/l as Mn)	Chloride, dissolved (mg/l as Cl)	Fluoride, dissolved (mg/l as F)	Alkalinity, bicarbonate (field) (mg/l as HCO <sub>3</sub> )	Sulfate, dissolved (mg/l as SO <sub>4</sub> )	Residue, total filtrable at 180°C (mg/l)
<b>Bishop Creek</b>									
12-15-82	5.0	0.55	0.16	0.023	4.2	< 0.10	27	1.0	54
03-30-83	7.6	0.76	0.19	0.038	5.7	< 0.1	31	1.1	69
06-14-83	2.5	0.34	0.17	0.023	1.1	< 0.10	13.5	2	35
08-24-83	6.2	0.55	0.033	0.025	6.3	0.10	26	2.7	23
12-07-83	5.1	0.45	1.0	0.04	3.9	0.12	22	< 2	46
03-20-84	4.6	0.56	0.6	0.028	4.3	< 0.1	22.5	< 2	50
<b>Capps Creek</b>									
12-15-82	2.5	0.57	0.088	0.022	< 1.0	< 0.10	31	2.0	49
03-30-83	3.6	0.63	0.23	0.035	< 1.0	< 0.1	36	1.5	63
06-14-83	0.97	0.37	0.20	0.054	< 1.0	< 0.10	10.5	2.2	27
08-24-83	2.2	0.41	0.020	0.078	< 1	< 0.10	23.5	2.7	26
12-07-83	2.6	0.44	0.55	0.05	< 1	0.10	42.5	3.8	39
03-20-84	2.8	0.53	0.2	0.046	1.8	< 0.1	29.5	< 2	60
<b>Middle Creek</b>									
12-16-82	3.3	0.59	0.21	0.016	1.4	< 0.10	36.5	1.9	85
03-31-83	4.7	0.77	0.95	0.035	1.9	< 0.1	46	< 1.0	80
06-14-83	3.1	0.54	0.59	0.022	< 1.0	< 0.10	29	2	52
08-24-83	3.5	0.55	0.051	0.045	1.6	< 0.10	37.5	3.3	42
12-07-83	3.3	0.42	1.0	0.07	1.5	< 0.1	31	< 2	41
03-20-84	3.1	0.51	0.7	0.002	1.9	< 0.1	27	< 2	40
<b>Lone Creek</b>									
12-16-82	4.0	0.74	0.25	0.023	2.4	< 0.10	35.5	1.8	56
03-31-83	4.4	0.89	0.19	0.042	2.8	< 0.1	40.5	< 1.0	77
06-14-83	3.0	0.60	0.61	0.049	1.1	< 0.10	28	2.1	49
08-24-83	4.2	0.64	0.035	0.054	2.4	< 0.10	34	3.3	48
12-07-83	4.4	0.55	1.2	0.08	2.2	< 0.1	32.5	< 2	44
03-20-84	3.6	0.64	1.2	0.034	2.9	< 0.1	28	< 2	90
<b>Chuitna River</b>									
12-16-82	2.0	0.40	0.21	0.012	2.4	< 0.10	33.5	< 1.0	47
03-31-83	4.1	0.71	0.26	0.013	1.4	< 0.1	39	1.6	84
06-14-83	1.5	0.32	0.10	0.009	< 1.0	< 0.10	14	2	33
08-24-83	2.8	0.46	0.087	< 0.02	1	< 0.10	30	2.9	34
12-07-83	3.2	0.44	0.41	< 0.03	1.1	< 0.1	27	< 2	38
03-20-84	2.7	0.56	0.5	< 0.002	2.4	< 0.1	24.5	< 2	100

Appendix B. Minor-element analysis of Beluga water-quality samples.

	Time	Streamflow, instantaneous (cfs)	Aluminum, total (ug/l as Al)	Aluminum, dissolved (ug/l as Al)	Antimony, total (ug/l as Sb)	Antimony, dissolved (ug/l as Sb)	Arsenic, total (ug/l as As)	Arsenic, dissolved (ug/l as As)	Barium, total (ug/l as Ba)	Barium, dissolved (ug/l as Ba)
Bishop Creek										
	03-30-83	1230	13	230	-	< 2	-	< 2	69	-
	06-14-83	0910	108	2200	320	3	< 2	4	30	20
	08-24-83	0940	28	< 60	-	< 2	-	< 2	420	-
	12-07-83	1115	<sup>a</sup> 23	400	-	< 10	-	< 2	20	-
	03-20-84	0930	35	270	-	< 5	-	< 2	10	-
Capps Creek										
	03-30-83	1420	5.2	190	-	< 2	-	< 2	77	-
	06-14-83	1110	157	12,000	300	9	< 2	16	140	20
	08-24-83	1105	16	< 60	-	< 2	-	< 2	420	-
	12-07-83	1250	<sup>b</sup> 7.0	400	-	< 10	-	< 2	20	-
	03-20-84	1100	<sup>b</sup> 12	190	-	< 5	-	< 2	30	-
Middle Creek										
	03-31-83	1410	4.3	58	-	< 2	-	< 2	45	-
	06-14-83	1350	13	79	20	< 2	< 2	< 2	20	30
	08-24-83	1220	9.7	< 60	-	< 2	-	< 2	420	-
	12-07-83	1535	9.4	300	-	< 10	-	< 2	< 10	-
	03-20-84	1230	13	60	-	< 5	-	< 2	10	-
Lone Creek										
	03-31-83	1150	9.6	52	-	< 2	-	< 2	53	-
	06-14-83	1515	28	75	41	< 2	< 2	< 2	20	20
	08-24-83	1320	20	< 60	-	< 2	-	< 2	420	-
	12-07-83	1445	13	300	-	< 10	-	< 2	10	-
	03-20-84	1350	14	65	-	< 5	-	< 2	10	-
Chuitna River										
	03-31-83	1100	<sup>c</sup> 120	120	-	< 2	-	< 2	45	-
	06-14-83	1700	<sup>c</sup> 470	300	41	< 2	< 2	< 2	10	20
	08-24-83	1445	<sup>d</sup> 139	< 60	-	< 2	-	< 2	420	-
	12-07-83	1400	<sup>d</sup> 100	300	-	< 10	-	< 2	< 10	-
	03-20-84	1450	<sup>d</sup> 100	80	-	< 5	-	< 2	5	-

<sup>a</sup>Estimate only, ice on probe head.

<sup>b</sup>U.S. Geological Survey (1985, p. 180).

<sup>c</sup>U.S. Geological Survey (1984, p. 159).

<sup>d</sup>U.S. Geological Survey (1985, p. 182).

Appendix B. (con.)

	Beryllium, total (ug/l as Be)	Beryllium, dissolved (ug/l as Be)	Boron, total (ug/l as B)	Boron, dissolved (ug/l as B)	Cadmium, total (ug/l as Cd)	Cadmium, dissolved (ug/l as Cd)	Chromium, total (ug/l as Cr)	Chromium, dissolved (ug/l as Cr)	Copper, total (ug/l as Cu)	Copper, dissolved (ug/l as Cu)
<b>Bishop Creek</b>										
03-30-83	< 2	-	480	-	< 0.5	-	< 4	-	< 5	-
06-14-83	< 0.2	< 0.2	70	50	< 0.5	< 0.5	< 5	< 5	< 5	< 5
08-24-83	< 1	-	< 0.05	-	< 0.5	-	< 5	-	< 5	-
12-07-83	< 1	-	0.14	-	< 0.5	-	< 2	-	< 3	-
03-20-84	< 0.2	-	< 50	-	< 0.5	-	< 5	-	< 5	-
<b>Capps Creek</b>										
03-30-83	< 2	-	74	-	< 0.5	-	< 4	-	< 5	-
06-14-83	< 0.2	< 0.2	170	70	< 0.5	< 0.5	14	< 5	20	< 5
08-24-83	< 1	-	< 0.05	-	< 0.5	-	< 5	-	< 5	-
12-07-83	< 1	-	0.06	-	< 0.5	-	< 2	-	< 3	-
03-20-84	< 0.2	-	< 50	-	< 0.5	-	< 5	-	< 5	-
<b>Middle Creek</b>										
03-31-83	< 2	-	< 50	-	< 0.5	-	< 4	-	< 5	-
06-14-83	< 0.2	< 0.2	50	50	< 0.5	< 0.5	< 5	< 5	< 5	e8
08-24-83	< 1	-	< 0.05	-	< 0.5	-	< 5	-	< 5	-
12-07-83	< 1	-	0.05	-	< 0.5	-	< 2	-	< 3	-
03-20-84	< 0.2	-	< 50	-	< 0.5	-	< 5	-	< 5	-
<b>Lone Creek</b>										
03-31-83	< 2	-	< 50	-	< 0.5	-	< 4	-	< 5	-
06-14-83	< 0.2	< 0.2	50	50	< 0.5	< 0.5	< 5	< 5	< 5	< 5
08-24-83	< 1	-	< 0.05	-	< 0.5	-	< 5	-	< 5	-
12-07-83	< 1	-	0.07	-	< 0.5	-	< 2	-	4	-
03-20-84	< 0.2	-	< 50	-	< 0.5	-	< 5	-	< 5	-
<b>Chuitna River</b>										
03-31-83	< 2	-	< 50	-	< 0.5	-	< 4	-	< 5	-
06-14-83	< 0.2	< 0.2	< 50	< 50	< 0.5	< 0.5	< 5	< 5	< 5	< 5
08-24-83	< 1	-	< 0.05	-	< 0.5	-	< 5	-	< 5	-
12-07-83	< 1	-	< 0.05	-	< 0.5	-	< 2	-	< 3	-
03-20-84	< 0.2	-	< 50	-	< 0.5	-	< 5	-	< 5	-

eContamination suspected.

## Appendix B. (con.)

	Iron, total (ug/l as Fe)	Iron, dissolved (ug/l as Fe)	Lead, total (ug/l as Pb)	Lead, dissolved (ug/l as Pb)	Manganese, total (ug/l as Mn)	Manganese, dissolved (ug/l as Mn)	Mercury, total (ug/l as Hg)	Mercury, dissolved (ug/l as Hg)	Nickel, total (ug/l as Ni)	Nickel, dissolved (ug/l as Ni)
Bishop Creek										
03-30-83	1300	-	< 5	-	52	-	< 0.05	-	< 5	-
06-14-83	2000	f <sub>13</sub>	< 5	< 5	60	25	< 0.05	< 0.05	< 5	< 5
08-24-83	1200	-	7	-	99	-	< 0.05	-	< 5	-
12-07-83	1000	-	< 5	-	40	-	< 0.05	-	< 5	-
03-20-84	770	-	< 2	-	30	-	0.5	-	< 5	-
Capps Creek										
03-30-83	680	-	< 5	-	43	-	< 0.05	-	< 5	-
06-14-83	8800	f <sub>98</sub>	< 5	< 5	280	35	0.1	< 0.05	15	< 5
08-24-83	2100	-	< 2	-	190	-	< 0.05	-	< 5	-
12-07-83	550	-	< 5	-	50	-	< 0.05	-	< 5	-
03-20-84	630	-	< 2	-	53	-	< 0.5	-	< 5	-
Middle Creek										
03-31-83	1200	-	< 5	-	44	-	< 0.05	-	5.7	-
06-14-83	670	f <sub>130</sub>	13	< 5	35	35	< 0.05	< 0.05	< 5	< 5
08-24-83	1500	-	< 2	-	66	-	< 0.05	-	< 5	-
12-07-83	1000	-	< 5	-	70	-	< 0.05	-	< 5	-
03-20-84	1000	-	< 2	-	42	-	< 0.5	-	< 5	-
Lone Creek										
03-31-83	1200	-	< 5	-	57	-	< 0.05	-	< 5	-
06-14-83	1100	f <sub>40</sub>	< 5	< 5	55	55	< 0.05	< 0.05	< 5	< 5
08-24-83	1700	-	< 2	-	84	-	< 0.05	-	< 5	-
12-07-83	1200	-	< 5	-	80	-	< 0.05	-	< 5	-
03-20-84	770	-	< 2	-	60	-	0.5	-	< 5	-
Chuitna River										
03-31-83	930	-	< 5	-	22	-	< 0.05	-	< 5	-
06-14-83	500	f <sub>170</sub>	< 5	< 5	35	15	< 0.05	< 0.05	< 5	< 5
08-24-83	710	-	< 2	-	< 20	-	< 0.05	-	< 5	-
12-07-83	410	-	< 5	-	< 30	-	< 0.05	-	< 5	-
03-20-84	600	-	< 2	-	22	-	0.5	-	< 5	-

f<sub>1</sub> Low values suspected. See Appendix A for dissolved iron values.

## Appendix B. (con.)

	Selenium, total (ug/l as Se)	Selenium, dissolved (ug/l as Se)	Silver, total (ug/l as Ag)	Silver, dissolved (ug/l as Ag)	Strontium, total (ug/l as Sr)	Strontium, dissolved (ug/l as Sr)	Titanium, total (ug/l as Ti)	Titanium, dissolved (ug/l as Ti)	Vanadium, total (ug/l as V)	Vanadium, dissolved (ug/l as V)	Zinc, total (ug/l as Zn)	Zinc, dissolved (ug/l as Zn)
Bishop Creek												
03-30-83	< 4	-	< 2	-	220	-	< 50	-	< 10	-	4.8	-
06-14-83	< 2	< 2	< 2	< 2	70	65	140	< 20	< 10	< 10	9.3	15
08-24-83	< 2	-	< 2	-	150	-	< 100	-	< 10	-	5.5	-
12-07-83	< 2	-	< 2	-	10	-	< 20	-	< 10	-	4	-
03-20-84	< 2	-	< 1	-	60	-	< 20	-	< 10	-	6	-
Capps Creek												
03-30-83	< 4	-	< 2	-	330	-	< 50	-	< 10	-	2.4	-
06-14-83	< 2	< 2	< 2	< 2	170	70	840	< 20	30	< 10	78	< 2
08-24-83	< 2	-	< 2	-	190	-	< 100	-	< 10	-	15	-
12-07-83	< 2	-	< 2	-	20	-	< 20	-	< 10	-	6	-
03-20-84	< 2	-	< 1	-	70	-	50	-	< 10	-	30	-
Middle Creek												
03-31-83	< 4	-	< 2	-	260	-	< 50	-	< 10	-	3.4	-
06-14-83	< 2	< 2	< 2	< 2	80	80	< 20	< 20	< 10	< 10	2	4
08-24-83	< 2	-	< 2	-	170	-	< 100	-	< 10	-	< 2	-
12-07-83	< 2	-	< 2	-	20	-	< 20	-	< 10	-	4	-
03-20-84	< 2	-	< 1	-	40	-	< 20	-	< 10	-	1	-
Lone Creek												
03-31-83	< 4	-	< 2	-	280	-	< 50	-	< 10	-	< 2	-
06-14-83	< 2	< 2	< 2	< 2	95	95	< 20	< 20	< 10	< 10	10	4
08-24-83	< 2	-	< 2	-	170	-	< 100	-	< 10	-	< 2	-
12-07-83	< 2	-	< 2	-	10	-	< 20	-	< 10	-	4	-
03-20-84	< 2	-	< 1	-	60	-	< 20	-	< 10	-	2	-
Chuitna River												
03-31-83	< 4	-	< 2	-	260	-	< 50	-	< 10	-	< 2	-
06-14-83	< 2	< 2	< 2	< 2	60	50	< 20	< 20	< 10	< 10	< 2	e <sub>6</sub>
08-24-83	< 2	-	< 2	-	140	-	< 100	-	< 10	-	< 2	-
12-07-83	< 2	-	< 2	-	10	-	< 20	-	10	-	5	-
03-20-84	< 2	-	< 1	-	60	-	< 20	-	< 10	-	5	-

e Contamination suspected.

Appendix C. Nutrient analysis of Beluga water-quality samples.

Date	Time	Streamflow, instantaneous (cfs)	Nitrogen, ammonia + organic total (mg/l as N)	Nitrogen, NO <sub>2</sub> + NO <sub>3</sub> dissolved (mg/l as N)	Nitrogen, ammonia dissolved (mg/l as N)	Phosphorus, total (mg/l as P)	Phosphorus, total reactive (dissolved) (mg/l as P)	Phosphorus, filterable reactive (ortho,dissolved) (mg/l as P)
<b>Bishop Creek</b>								
12-15-82	0935	15	a <sub>-</sub>	0.223	0.023	0.031	0.014	0.012
03-30-83	1230	13	0.10	0.177	0.030	0.029	0.005	0.005
06-14-83	0910	108	0.15	0.541	0.136	b <sub>0.024</sub>	0.088	0.044
08-24-83	0940	28	0.14	0.056	0.005	0.060	0.046	0.029
03-20-84	0930	35	0.17	0.172	0.009	0.025	0.015	0.009
<b>Capps Creek</b>								
12-15-82	1225	8.8	a <sub>-</sub>	0.333	0.016	0.034	0.008	0.008
03-30-83	1420	5.2	0.09	0.292	0.005	0.016	0.015	0.015
06-14-83	1110	157	0.26	0.475	0.033	0.140	0.040	0.030
08-24-83	1105	16	0.13	0.123	0.007	0.056	0.005	0.004
03-20-84	1100	12	0.13	0.351	0.011	0.019	0.007	0.008
<b>Middle Creek</b>								
12-16-82	1225	6.9	a <sub>-</sub>	0.092	0.028	0.021	0.022	0.019
03-31-83	1410	4.3	0.11	0.110	0.021	a <sub>0.031</sub>	0.022	0.021
06-14-83	1350	13	a <sub>-</sub>	0.034	0.012	0.012	0.010	0.007
08-24-83	1220	9.7	0.19	0.013	0.012	0.019	0.016	0.017
03-20-84	1230	13	0.21	0.049	0.014	0.020	0.012	0.012
<b>Lone Creek</b>								
12-16-82	1100	16	0.11	0.185	0.022	0.017	0.015	0.013
03-31-83	1150	9.6	0.11	0.130	0.017	0.019	0.011	0.010
06-14-83	1515	28	0.16	0.012	0.011	0.021	0.011	0.009
08-24-83	1320	20	0.20	0.025	0.011	0.037	0.013	0.014
03-20-84	1350	14	0.21	0.102	0.019	0.015	0.010	0.010
<b>Chuitna River</b>								
12-16-82	0900	d <sub>100</sub>	a <sub>-</sub>	0.208	0.015	0.021	0.010	0.009
03-31-83	1100	d <sub>120</sub>	0.07	0.171	0.013	0.023	0.011	0.012
06-14-83	1700	d <sub>470</sub>	0.10	0.107	0.013	0.029	0.010	0.009
08-24-83	1445	139	0.14	0.023	0.006	0.010	0.005	0.008
03-20-84	1450	e <sub>100</sub>	0.15	0.130	0.005	0.018	0.011	0.011

<sup>a</sup>Missing data.

<sup>b</sup>Erroneous value suspected.

<sup>c</sup>U.S. Geological Survey (1985, p. 180).

<sup>d</sup>U.S. Geological Survey (1984, p. 159).

<sup>e</sup>U.S. Geological Survey (1985, p. 182).

Appendix D-1. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, June 15, 1983. Roman numeral represents stratum sample at site.

Taxon	Site											
	1				2				5			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Insecta</b>												
Ephemeroptera												
Unidentified												
Ephemeroptera	10	60	10	20	80	40	20	50	190	100	70	
Baetis bicaudatus	60	50	180	80	140	480	120	150	70	20	20	
Baetis tricaudatus	110	110	260	100	800	1120	910	740	30	120	80	20
Baetis sp.	130	90	200	90	130	350	80	180	500	1050	430	50
Cinygmula sp.	40	10							10			
Ephemerella doddsi		10	20	20					20	30	50	
Ephemerella infrequens/ E. inermis complex					30	10	30		10		10	
Unidentified												
Heptageniidae	430	370	620	280	140	220	60	120	170	90	180	70
Plecoptera												
Unidentified Plecoptera					10		10	20				
Unidentified												
Chloroperlidae	150	190	220	90		10	10		310	230	290	150
Isoperla sp.									10	10		
Unidentified Perlodidae						60						
Zapada cinctipes	20	20	60			20			10			20
Zapada oregonensis	110	140	120	40	30	40		20		10	20	
Trichoptera												
Unidentified Trichoptera							10			10		
Apatania sp.	10				10		60					
Brachycentrus sp.	70	20	400	60		20	30	10	60	100	80	
Glossosoma sp.	30	60	20	40					40	50	10	
Ochrotrichia sp.		90	140	10	10				220	380	300	80
Onocosmoecus sp.					10	10		30				
Unidentified Limnephilidae	50	10		30			20		150	70	160	30
Rhyacophila vepulsa			30									
Rhyacophila sp.	40	60	70	20						10		



## Appendix D-1 (con.)

Taxon	Site											
	I	II	1 III	IV	I	II	2 III	IV	I	II	5 III	IV
Diptera												
Atherix sp.	10											
Chelifera sp.	10	60	10	10	150	80	20	30			20	
Unidentified Chironomidae	3680	7810	6100	2690	6970	4460	4680	3570	2160	4130	4050	2980
Dicranota sp.	20	20	40	10	120	50	20	40	10	10	20	
Palpomyia sp.	100	50	20	10	80	10		10				
Prosimulium sp.		10	50	20	20	10					10	
Unidentified Simuliidae										20		
Simulium sp.	1440	840	31640	9220	4220	36540	1590	2630	1370	1470	3580	130
Hymenoptera												
Unidentified Hymenoptera								10				
Collembola	10					20	20			30	10	
Turbellaria	10		40	10					30	30		10
Nematoda	90	10		30	100	70	140	30	10			
Oligochaeta	50	30	40	30	830	530	20	40	10	20	10	20
Pelecypoda				40	90	10	10					
Arachnida												
Acarina	550	1020	820	370	340	350	190	650	140	330	150	70
Crustacea												
Ostracoda							10		10	10	20	90
Copepoda				10	60				10			
Total number of invertebrates/m <sup>2</sup>	7230	11140	41110	13330	14370	44510	8060	8330	5550	8330	9570	3720
Total number of taxa (based on number of insect families and other invertebrates)	19	18	17	21	16	18	16	14	19	19	17	12
Shannon-Weaver Diversity Index	2.40	1.73	1.23	1.50	2.09	1.02	1.84	2.03	2.59	2.31	2.11	1.34
Evenness	0.56	0.42	0.30	0.34	0.52	0.25	0.46	0.53	0.61	0.54	0.52	0.38

Appendix D-2. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, June 15, 1983. Roman numeral represents stratum sample at site.

Taxon	Site											
	7				8				11			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
Insecta												
Ephemeroptera												
Unidentified												
Ephemeroptera		10			10		10		40	40	70	20
<i>Baetis bicaudatus</i>	50	30	20	20	10		10			10	20	10
<i>Baetis tricaudatus</i>	160	280	200	130	260	140	50		110	60	120	130
<i>Baetis</i> sp.	40	410	250	310	140	310	50	10	50	450	340	500
<i>Ephemerella doddsi</i>	50		40	10					20	50	20	10
<i>Ephemerella infrequens</i> /												
<i>E. inermis</i> complex				20	10		10		10	10		
Unidentified												
Heptageniidae	350	380	370	780	230	190	10	10	50	80	70	20
Plecoptera												
Unidentified Plecoptera		20	30	20			20			10		
Unidentified												
Chloroperlidae	120	310	130	240	220	160	10	10	110	150	80	60
Isoperla sp.				10					50	10		
Unidentified Perlodidae			10		10							20
<i>Zapada cinctipes</i>						10				10		
<i>Zapada oregonensis</i>	20	10	20	30			10			10		
Trichoptera												
Unidentified Trichoptera				10	20				10	20		10
<i>Brachycentrus</i> sp.	60	20	30	20	10	10			30	10		120
<i>Glossosoma</i> sp.	90	30	70	40	20	30			10			10
<i>Ochrotrichia</i> sp.	30	100							130	300	930	1400
<i>Onocosmoecus</i> sp.	10			60	10		10					
Unidentified Limnephilidae	30	20	20	80	30	10				10	10	50
<i>Rhyacophila</i> sp.	10	50	20							20		

Appendix D-2 (con.)

Taxon	Site 7							Site 8						
	I	II	III	IV	I	II	III	I	II	III	IV	I	II	III
Diptera														
Chelifera sp.		20										20	20	20
Unidentified Chironomidae	9920	19830	11840	18180	11750	13380	18280	470	470	30	470	660	4130	3000
Dicranota sp.	10	50	40	50	50		10	10	10	10	10	10	20	10
Palpomyia sp.	110	20	60	350	20		10	20	20	10	20	30	20	10
Pericoma sp.		20												
Prosimulium sp.	30	130	1060	50	10	60								
Unidentified Simuliidae		20												
Simulium sp.	24470	32030	62110	8400	2670	6820	310	120	120	310	120	820	7760	6460
Molophilus sp.								10				10		13830
Collembola	10	10						10						
Turbellaria	70			10		10								10
Nematoda	80	90	20	440	10	30	10					30	30	20
Oligochaeta	300		30	20	140	1720	100	480	480	100		10	10	20
Arachnida														
Acarina	290	360	190	1180	240	50	80	10	10	80	10	50	240	180
Crustacea														260
Ostracoda	20	10	30					10	10		10	10	10	50
Copepoda														
Total number of invertebrates/m <sup>2</sup>	36330	54260	76590	30460	15880	22930	19020	1160	1160	19020	1160	2270	13460	11410
Total number of taxa (based on number of insect families and other invertebrates)	20	19	18	17	16	13	14	11	11	14	11	17	18	15
Shannon-Weaver Diversity Index	1.30	1.29	0.82	1.69	1.32	1.54	0.33	1.91	1.91	0.33	1.91	2.71	1.65	1.75
Evenness	0.30	0.30	0.20	0.41	0.33	0.42	0.09	0.55	0.55	0.09	0.55	0.66	0.40	0.45

Appendix D-3. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, August 25, 1983. Roman numeral represents stratum sample at site.

Taxon	Site											
	1				2				5			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Insecta</b>												
<b>Ephemeroptera</b>												
<i>Baetis bicaudatus</i>						10				160	30	10
<i>Baetis tricaudatus</i>												10
<i>Baetis</i> sp.	1110	3220	2640	2530	1560	3390	4070	1360	510	1000	440	360
<i>Cinygmula</i> sp.		10	10					10				
<i>Ephemerella doddsi</i>	10	10	60	20		10				40	30	10
<i>Ephemerella infrequens</i> /												
<i>E. inermis</i> complex		40		20	130	140	30	20				
Unidentified Heptageniidae	1660	2270	2770	2840	400	580	380	120	110	240	1010	1580
Unidentified Siphonuridae		40	40	10	40		10	10			10	60
<b>Plecoptera</b>												
Unidentified Capniidae	480	1580	2140	1700	910	880	1340	510	30	240	280	640
Unidentified												
Chloroperlidae	100	30	50	50					50	40	40	130
<i>Isoperla</i> sp.		10	30						10		20	30
Unidentified Perlodidae												
<i>Skwala</i> sp.		10	20	10		20	10	20				
<i>Zapada cinctipes</i>	680	1320	1180	920	900	1830	660	240	80	130	330	150
<i>Zapada oregonensis</i>		10		10	40		10			20	10	
<i>Zapada</i> sp.	180	530	230	400	10	20	10				40	40
<b>Trichoptera</b>												
<i>Apatania</i> sp.	10	20			30	30	130	210				
<i>Brachycentrus</i> sp.	280	130	170	120		10	30	40		20	100	
<i>Ecclisomyia</i> sp.					10				20	10		60
<i>Glossosoma</i> sp.	760	670	640	710	70	190	110	130	110	380	170	110
Unidentified Limnephilidae	20	20	20		10				20	30	20	20
<i>Rhyacophila vepulsa</i>	50	100	90	10								10

## Appendix D-3 (con.)

Taxon	1				Site 2				5			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Diptera</b>												
Chelifera sp.	10	10	10		10	10	60					30
Unidentified Chironomidae	5330	7020	12420	5810	7610	5970	10050	12060	350	1230	980	2740
Dicranota sp.	190	60	120	140	410	500	440	210	30	40	50	20
Hesperoconopa sp.										10	10	
Palpomyia sp.	210	120	210	350	260	140	230	40		10	10	50
Pericoma sp.	620	680	1130	1320	300	40	30	110		10	370	750
Unidentified Simuliidae	10											
Simulium sp.	10	160	60	70	20	10	10	20				
Unidentified Tipulidae					10							
Tipula sp.								10				20
<b>Hymenoptera</b>												
Unidentified Hymenoptera	10						10					
<b>Collembola</b>			10									
Turbellaria	100	280	310	110	30		10	30	30	40	150	50
Nematoda	40			40	30					10		10
Oligochaeta	90	50	70	610	1260	370	150	210	110	160	960	1530
Pelecypoda					110	10		70		40		
Gastropoda							10					
<b>Arachnida</b>												
Acarina	490	560	670	180	680	660	530	640	70	170	90	330
<b>Crustacea</b>												
Cladocera	10			10	10			10	60	40	10	10
Ostracoda	60	130	640	270	60	40	50	40	30	190	110	360
Copepoda	20	20	10	150	400	30	70	40	10	10	10	10
<b>Total number of invertebrates/m<sup>2</sup></b>	<b>12550</b>	<b>19110</b>	<b>25760</b>	<b>18420</b>	<b>15310</b>	<b>14890</b>	<b>18440</b>	<b>16160</b>	<b>1630</b>	<b>4270</b>	<b>5280</b>	<b>9130</b>
<b>Total number of taxa (based on number of insect families and other invertebrates)</b>	<b>25</b>	<b>23</b>	<b>24</b>	<b>23</b>	<b>22</b>	<b>20</b>	<b>23</b>	<b>22</b>	<b>16</b>	<b>21</b>	<b>21</b>	<b>23</b>
<b>Shannon-Weaver Diversity Index</b>	<b>2.97</b>	<b>2.92</b>	<b>2.70</b>	<b>3.12</b>	<b>2.74</b>	<b>2.64</b>	<b>2.19</b>	<b>1.64</b>	<b>3.17</b>	<b>3.11</b>	<b>3.36</b>	<b>3.12</b>
<b>Evenness</b>	<b>0.64</b>	<b>0.65</b>	<b>0.59</b>	<b>0.69</b>	<b>0.61</b>	<b>0.61</b>	<b>0.48</b>	<b>0.37</b>	<b>0.79</b>	<b>0.71</b>	<b>0.77</b>	<b>0.69</b>

Appendix D-4. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, August 25, 1983. Roman numeral represents stratum sample at site.

Taxon	7				8				11			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Insecta</b>												
<b>Ephemeroptera</b>												
<u>Baetis bicaudatus</u>			10			50	30	30		30	10	
<u>Baetis tricaudatus</u>	20		20		30				10		30	
<u>Baetis</u> sp.	2750	2680	4810	2140	3230	2800	1170	3690	260	470	590	270
<u>Epeorus</u> sp.				10								
<u>Ephemerella doddsi</u>	40		20	40	160	90	30	50		250	280	130
<u>Ephemerella infrequens</u> / <u>E. inermis</u> complex	40		10	10	10	20		60		60	20	30
Unidentified Reptageniidae	1890	2170	810	450	480	1010	320	270		290	460	220
<u>Rhithrogena</u> sp.			10						10	40	110	10
Unidentified Siphonuridae			20	20	10							10
<b>Plecoptera</b>												
Unidentified Capniidae	1100	1400	420	310	880	700	400	180	60	140	150	220
Unidentified <u>Chloroperlidae</u>	20	80	70	120	80	180	90	10	30	70	80	30
<u>Isoperla</u> sp.	120			40		50						
Unidentified Perlodidae		10	10		40		30	40		90	20	30
<u>Skwala</u> sp.	20			10		10						
<u>Taenionema</u> sp.		10		10						10	10	10
<u>Zapada cinctipes</u>	2490	900	480	660	350	400	70	160	60	740	440	350
<u>Zapada oregonensis</u>	10		40	10			10	20		10		20
<u>Zapada</u> sp.	370	110	130	190	20	20		40	10			40
<b>Trichoptera</b>												
<u>Apatania</u> sp.							20	50				
<u>Brachycentrus</u> sp.	460	130	560	210	150	110		110	30	190	70	110
<u>Ecclisomyia</u> sp.					60		10	60		60		
<u>Glossosoma</u> sp.	670	330	760	1150	110	380	50	300	10	1530	870	140
Unidentified Limnephilidae	60	10	10			20	10		20	30	10	20
<u>Onocosmoecus</u> sp.												10
<u>Psychoglypha</u> sp.	10											
<u>Rhyacophila</u> sp.						10						

## Appendix D-4 (con.)

Taxon	7				Site 8				11			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
Diptera												
Chelifera sp.	20					10				30	10	
Unidentified Chironomidae	9020	14480	3620	3700	3250	1380	340	530	130	350	680	700
Dicranota sp.	210	210	120	130	290	160	190	260	20	20	60	20
Palpomyia sp.	90	30	30	90	60	30	10	20				20
Pericoma sp.	1800	2930	2910	1900	1250	480	140	30	60	760	420	50
Unidentified Simuliidae									10			
Simulium sp.		30	50	10	40	40		20			40	20
Unidentified Tipulidae					20							
Tipula sp.	10											
Hymenoptera												
Unidentified Hymenoptera	10					10						
Collembola	10				10			10				
Turbellaria	230	40	30	70	20	20	90	20	40	40	610	170
Nematoda		20	20	120	20	10	220	90	10			
Oligochaeta	40	270	170	140	750	80	370	920	60	120	1070	390
Pelecypoda								10		10		
Arachnida												
Acarina	1100	550	430	450	410	150	70	40	30	220	80	100
Crustacea												
Cladocera	60	40	40	30	10				20	80		
Ostracoda	130	130	80	40	80	50	20	20	20	40	40	10
Copepoda	20			30	70				10	50	30	10
Total number of invertebrates/m <sup>2</sup>	22820	26560	15690	12090	11890	8270	3690	7040	910	5730	6190	3140
Total number of taxa (based on number of insect families and other invertebrates)	23	21	22	23	24	23	18	22	19	22	21	22
Shannon-Weaver Diversity Index	2.93	2.35	2.88	3.11	3.10	3.11	3.29	2.61	3.51	3.50	3.55	3.60
Evenness	0.65	0.53	0.65	0.69	0.68	0.69	0.79	0.59	0.83	0.78	0.81	0.81

Appendix D-5. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, June 14, 1984. Roman numeral represents stratum sample at site.

Taxon	Site											
	1				2				5			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Insecta</b>												
Ephemeroptera												
Unidentified												
Ephemeroptera	50	50	20	130	80		50	30	180	330	120	60
Baetis bicaudatus	70	30	80	100	10	30	40	10	320	1050	260	20
Baetis tricaudatus	90	70	150	70	120	110	200	50	110	140	150	130
Baetis sp.		10		60	40		100	30	730	130	10	150
Ephemerella doddsi	10		50						20	130	10	
Ephemerella infrequens/ E. inermis complex					30							
Unidentified Heptageniidae	460	230	830	460			230	10	330	340	310	160
Plecoptera												
Unidentified Plecoptera			10									
Unidentified												
Chloroperlidae	80	70	320	190		10	20		130	100	110	190
Isoperla sp.	10											
Unidentified Perlodidae			10	10	30	10	60	10				
Zapada cinctipes	30		10	10								
Zapada oregonensis	60	10	20	50								
Trichoptera												
Unidentified Trichoptera			100		70	50	150	20		10		70
Apatania sp.					20	90		10				
Brachycentrus sp.	70	50	60	40						40	10	
Glossosoma sp.	80	70	70	80					300	270	80	10
Ochrotrichia sp.			10		10				340	570	420	760
Onocosmoecus sp.					40	20	20					
Psychoglypha sp.								100				
Unidentified Limnephilidae	20	20		40	150	80		670	90	60	30	90
Rhyacophila vepulsa	30		10									
Rhyacophila sp.										10		



## Appendix D-5 (con.)

Taxon	1				Site 2				5			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
Diptera												
Chelifera sp.					360	30	50		10	20		
Unidentified Chironomidae	2640	1620	3920	4520	12630	6670	11090	5470	3260	4010	3650	7270
Dicranota sp.	50	10	90	60	190	160	60	10	30			30
Dixa sp.											10	
Palpomyia sp.	290	40	60	240	240	30	30		10		10	250
Prosimulium sp.			20	10								
Simulium sp.	3260	1910	8560	5210	710	280	5820	330	180	120	20	10
Unidentified Tipulidae						10						
Hymenoptera												
Unidentified Hymenoptera	30							10	10			
Turbellaria			30	40	10		30		60	30	10	40
Nematoda	10	20		20	240	110	50	60			10	
Oligochaeta	420	70	30	570	340	870	140	640	130	20	140	790
Pelecypoda					130	20		80				
Gastropoda								10				
Arachnida												
Acarina	110	100	50	340	270	140	320	30	70	240	250	680
Crustacea												
Cladocera		50		40				10	10	30		
Ostracoda	10	20	10	40	220	20	10	50	120	40		100
Copepoda		10		20	30	10			40			30
Total number of invertebrates/m <sup>2</sup>	7880	4460	14520	12350	15970	8750	18470	7640	6480	7690	5610	10840
Total number of taxa (based on number of insect families and other invertebrates)	19	17	18	19	17	15	15	15	19	17	16	15
Shannon-Weaver Diversity Index	2.35	2.19	1.70	2.17	1.42	1.42	1.46	1.53	2.48	2.30	1.90	1.86
Evenness	0.55	0.54	0.41	0.51	0.35	0.36	0.37	0.39	0.58	0.56	0.48	0.48

Appendix D-6. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, June 14, 1984. Roman numeral represents stratum sample at site.

Taxon	7				Site 8				11			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Insecta</b>												
Ephemeroptera												
Unidentified Ephemeroptera	40	10		10	10	30	30	30	280	520	120	180
<u>Baetis bicaudatus</u>	250	320	160		360	670	80	190	200	1500	1360	320
<u>Baetis tricaudatus</u>	40	130	140	70	30	20	80		260	240	150	190
<u>Baetis</u> sp.	30	50	20	130	30	30	130		310	1290	120	190
<u>Cinygmula</u> sp.								60				
<u>Epeorus</u> sp.	10		30									
<u>Ephemerella doddsi</u>		40	10	10	20				30	70	30	50
<u>Ephemerella infrequens</u> /												
<u>E. inermis</u> complex		10		10				10	10			10
Unidentified Heptageniidae	740	790	410	520	40	90	150	40	210	260	210	100
Plecoptera												
Unidentified Plecoptera						10			10			
Unidentified Capniidae										10		10
Unidentified Chloroperlidae	150	480	260	120	470	250	390	130	180	510	300	100
<u>Isoperla</u> sp.	20		50			20	10	10	20	10		
Unidentified Perlodidae	30	20		40	40	40	20	30	20			30
<u>Zapada oregonensis</u>	40	20	30			10	10	10				
Trichoptera												
Unidentified Trichoptera	10			20		10	130	50			10	10
<u>Brachycentrus</u> sp.	40	30	30	10		10		40	70	70	30	80
<u>Glossosoma</u> sp.	260	350	60	20	90	70	70	260	10	40		10
<u>Ochrotrichia</u> sp.					10		10		250	950	660	240
<u>Onocosmoecus</u> sp.	20						20	60				
<u>Psychoglypha</u> sp.								20				
Unidentified Limnephilidae			10	30		10	160	420	30		10	10
<u>Rhyacophila</u> sp.									10			

## Appendix D-6 (con.)

Taxon					Site							
	I	II	7 III	IV	I	II	8 III	IV	I	II	11 III	IV
Diptera												
Chelifera sp.		10					10	10	20			
Unidentified Chironomidae	4790	7710	5640	8380	7850	8800	13250	8410	3310	4680	3320	7040
Dicranota sp.	60	20	40		40		20	30	30	60	40	10
Palpomyia sp.	170	50	60	70	10		10	10	10	20	10	40
Prosimulium sp.			20	10		20						
Simulium sp.	12770	8680	34050	3970	2570	16120	570	330	1610	910	1420	1440
Hymenoptera												
Unidentified Hymenoptera		10								10		
Collembola	10											
Turbellaria	70	10	10		60	10	50	20	50			20
Nematoda	240	90	70	40	140	180	60	30	20		30	30
Oligochaeta	260	30	10	40	990	420	500	520	90	310		30
Pelecypoda		10		30			20					
Arachnida												
Acarina	180	110	70	230	20	20	250	130	280	180	90	190
Crustacea												
Cladocera									110	160	10	40
Ostracoda		20	10	20		10	20	10	30	80	60	80
Copepoda					20				80	40	20	10
Total number of invertebrates/m <sup>2</sup>	20230	19000	41190	13780	12800	26850	16050	10860	7540	11920	8000	10460
Total number of taxa (based on number of insect families and other invertebrates)	17	20	18	16	16	15	19	19	22	19	16	21
Shannon-Weaver Diversity Index	1.72	1.83	0.90	1.58	1.82	1.41	1.19	1.47	2.59	2.55	2.38	1.75
Evenness	0.42	0.42	0.22	0.39	0.45	0.36	0.28	0.35	0.58	0.60	0.59	0.40

Appendix D-7. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Middle Creek, Beluga coal area, August 29, 1984. Roman numeral represents stratum sample at site.

Taxon	Site											
	1				2				5			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Insecta</b>												
<b>Ephemeroptera</b>												
Baetis sp.	570	710	320	70	430	630	340	230	280	500	560	210
Cinygma sp.											10	10
Ephemerella doddsi	30			20					70	200	80	20
Ephemerella infrequens/ E. inermis complex				20	10		10		10		40	20
Unidentified Heptageniidae	250	390	200	100	180	480	550	120	560	1230	750	1010
Rhithrogena sp.									60	20	10	10
Unidentified Siphonuridae				10			20		10	10	40	10
<b>Plecoptera</b>												
Unidentified Plecoptera					10							
Unidentified Capniidae	380	1450	420	790	540	600	480	490	120	190	180	770
Unidentified Chloroperlidae	90	70	10	90			20		50	80	10	100
Isoperla sp.									10	20	10	
Unidentified Perlodidae									10			
Skwala sp.					40	30	10					
Zapada cinctipes	90	60	110	40	310	530	310	10	30	320	870	170
Zapada oregonensis			20							40		
Zapada sp.	40	40	40		10	20				20	40	20
<b>Trichoptera</b>												
Unidentified Trichoptera		20	10	10	40	10	20			10		
Apatania sp.	10	10		10	350	240	80	90				
Brachycentrus sp.	10	20	20			20	10		190	310	140	
Ecclisomyia sp.	60			70	20				30	40	100	10
Glossosoma sp.	800	620	480	280	100	40	40	30	1560	3620	1030	740
Arctopsyche sp.									10	10	10	
Psychoglypha sp.										10	10	
Unidentified Limnephilidae	30	30	30	20	30			10	10	30	140	
Rhyacophila vepulsa		10	10								50	10
Rhyacophila sp.			10						10	10		

## Appendix D-7 (con.)

Taxon	Site											
	I	II	III	IV	I	II	III	IV	I	II	III	IV
Diptera												
Chelifera sp.			10	10	230	40	20	20	30	120	10	
Unidentified Chironomidae	2300	5170	1880	2430	9900	15980	8770	7230	1980	2890	1480	1140
Dicranota sp.	30	70		10	190	280	250	170	80	120	20	10
Hesperoconopa sp.		10								30	10	
Palpomyia sp.	440	220	130	290	1740	1610	430	90	20	30	10	280
Pericoma sp.	100	260	230	120	1120	1560	970	140	440	2020	200	400
Prosimulium sp.										10	10	
Simulium sp.	130	60	40	20				10	10	10		
Tipula sp.						30						
Hymenoptera												
Unidentified Hymenoptera		10			10			20	10			
Collembola								20	30			
Turbellaria	50	120	20	10				20	60	60	240	30
Nematoda	50	10			320	20	30			10		
Oligochaeta	260	240	50	80	270	140	250	40	90	130	180	320
Pelecypoda					620		50	30				
Gastropoda					20		10	10				
Arachnida												
Acarina	760	670	450	30	550	790	390	580	240	610	70	40
Crustacea												
Cladocera		10		10	40		10	40	130	80	20	100
Ostracoda	70	80	20	20	480	130	110	350	390	280	20	60
Copepoda	180	380	100	100	910	220	50	20	130	10		40
Total number of invertebrates/m <sup>2</sup>	6730	10740	4610	4660	18470	23400	13230	9770	6660	13080	6350	5530
Total number of taxa (based on number of insect families and other invertebrates)	20	22	19	21	22	18	24	22	27	26	24	20
Shannon-Weaver Diversity Index	3.28	2.77	2.99	2.54	2.72	1.95	2.11	1.71	3.29	3.16	3.39	3.32
Evenness	0.76	0.62	0.70	0.58	0.61	0.47	0.46	0.38	0.69	0.67	0.74	0.77

Appendix D-8. Density (numbers/m<sup>2</sup>), number of taxa, diversity, and evenness of benthic invertebrates collected in Lone Creek, Beluga coal area, August 29, 1984. Roman numeral represents stratum sample at site.

Taxon	7				Site 8				11			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
<b>Insecta</b>												
<b>Ephemeroptera</b>												
Baetis sp.	900	790	360	280	470	660	590	150	130	350	420	270
Cinygma sp.												10
Ephemerella doddsi	20	20	40	40		10	20		40	70	100	10
Ephemerella infrequens/ E. inermis complex		10	10		10			30	30	20		20
Unidentified Heptageniidae	380	560	510	470	180	790	30	80	540	190	230	190
Rhithrogena sp.							10		40	40	50	10
Unidentified Siphonuridae		20		10	10				10		20	
<b>Plecoptera</b>												
Unidentified Capniidae	790	1360	730	630	1000	2230	90	210	920	40	110	580
Unidentified Chloroperiidae	20	160	120	170	100	140	30		200	70	90	70
Isoperla sp.	20	10				10	20	10		30	30	
Pteronarcissa badia									10	10	10	
Skwala sp.		10		10				30				
Zapada cinctipes	890	400	110	170	60	330	10	40	270	540	130	60
Zapada oregonensis		10	20								10	
Zapada sp.	120			30	20	20	10	20				
<b>Trichoptera</b>												
Apatania sp.	20	20	50	100		20	930	310				
Brachycentrus sp.	120	40	40	170	20	50	40	10	20	40	80	
Ecclisomyia sp.				10				20				
Glossosoma sp.	270	440	470	660	200	840	1180	630	380	140	420	30
Psychoglypha sp.			10									
Unidentified Limnephilidae	20	30	10				10		10			10

## Appendix D-8 (con.)

Taxon	7				Site 8				11			
	I	II	III	IV	I	II	III	IV	I	II	III	IV
Diptera												
Chelifera sp.	10	10	10					20	50	40	20	
Unidentified Chironomidae	4950	21550	10640	15560	1240	3540	830	1900	1890	380	1440	770
Dicranota sp.	220	230	170	440	240	250	70	50	380	80	260	10
Palpomyia sp.	680	390	330	490	200	180			20		10	120
Pericoma sp.	1390	2180	2650	1610	120	1060	10	30	2310	150	760	210
Simulium sp.	60	120	60			30			20	20	40	
Collembola				10					10		10	10
Turbellaria	60	40	120	40	40	60			80	20	10	90
Nematoda	1160	70	20	20		10	270	190			20	20
Oligochaeta	320	40	40	60	290	510	210	180	330	40	300	840
Pelecypoda				10		10		10	10	10		
Arachnida												
Acarina	880	1020	500	800	130	470	150	400	530	140	370	90
Crustacea												
Cladocera	70	30	40	50	20	110	30	10	70	80	120	80
Ostracoda	330	300	60	500	90	210		10	460	190	160	170
Copepoda	40	710	40	40	20	60	20		120	30	10	40
Total number of invertebrates/m <sup>2</sup>	13740	30570	17160	22380	4460	11600	4560	4340	8880	2720	5230	3710
Total number of taxa (based on number of insect families and other invertebrates)	23	24	22	24	19	23	18	19	25	22	25	20
Shannon-Weaver Diversity Index	3.28	1.91	2.14	1.98	3.25	3.25	2.99	2.84	3.38	3.74	3.56	3.36
Evenness	0.72	0.42	0.48	0.43	0.76	0.73	0.72	0.67	0.73	0.84	0.77	0.78

Appendix E. Three-factor analysis of variance table, where the variable is benthic-invertebrate abundance (in numbers/m<sup>2</sup>) during June and August 1983 and 1984, Middle and Lone Creeks, Beluga coal area, Alaska. The number of samples used in this analysis = 96.

Source of variation	Degrees of freedom	Sum of squares	Mean sum of squares	Calculated F	Critical F <sup>a</sup>	Conclusion
stream	1	0.038	0.038	0.587	F <sub>0.05</sub> [1,72] = 4.00	Accept H <sub>0</sub>
site	2	3.023	1.511	23.182	F <sub>0.05</sub> [2,72] = 3.15	Reject H <sub>0</sub>
date	3	0.607	0.202	3.106	F <sub>0.05</sub> [3,72] = 2.76	Reject H <sub>0</sub>
stream X site	2	1.299	0.649	9.963	F <sub>0.05</sub> [2,72] = 3.15	Reject H <sub>0</sub>
stream X date	3	0.536	0.179	2.742	F <sub>0.05</sub> [3,72] = 2.76	Accept H <sub>0</sub>
site X date	6	0.718	0.120	1.836	F <sub>0.05</sub> [6,72] = 2.25	Accept H <sub>0</sub>
stream X site X date	6	0.361	0.060	0.923	F <sub>0.05</sub> [6,72] = 2.25	Accept H <sub>0</sub>

<sup>a</sup>There are no critical values for v<sub>2</sub> = 72, so the values for the next lowest degrees of freedom, v<sub>2</sub> = 60, were used.



Appendix F. Wet-weight biomass (g/m<sup>2</sup>) of benthic invertebrates collected from sites in Middle and Lone Creeks, Beluga coal area, during June and August 1983 and 1984. Biomass is summed for each stream, site and month. Roman numeral represents stratum sample at site.

		Wet-weight Biomass (g/m <sup>2</sup> )						
		Middle Creek			Lone Creek			
Month	Sample	1	2	5	7	8	11	
June 1983	I	6.73	12.45	6.94	18.71	14.12	5.36	
	II	8.17	16.61	5.28	21.21	31.05	11.45	
	III	18.75	6.87	11.99	40.88	19.46	7.99	
	IV	<u>9.78</u>	<u>8.78</u>	<u>4.13</u>	<u>24.69</u>	<u>6.04</u>	<u>18.73</u>	
	Σ	43.43	44.71	28.34	116.48	105.49	43.53	219.69
August 1983	I	9.40	7.41	3.13	14.01	10.64	3.08	
	II	9.41	10.73	6.31	8.79	11.01	21.78	
	III	9.90	8.81	7.08	14.77	7.35	11.13	
	IV	<u>7.78</u>	<u>11.32</u>	<u>8.25</u>	<u>8.64</u>	<u>15.25</u>	<u>5.90</u>	
	Σ	36.49	38.27	24.77	99.53	46.21	41.89	132.35
June 1984	I	9.89	13.41	7.77	17.11	17.60	9.99	
	II	5.62	11.57	14.47	20.49	26.09	13.17	
	III	12.77	11.25	6.47	23.67	16.81	10.05	
	IV	<u>7.51</u>	<u>7.83</u>	<u>5.83</u>	<u>10.95</u>	<u>21.61</u>	<u>12.74</u>	
	Σ	35.79	44.06	34.54	114.39	72.22	45.95	200.28
August 1984	I	10.11	6.84	8.46	6.07	5.40	9.11	
	II	8.41	8.88	12.72	8.25	12.08	4.86	
	III	7.18	5.51	7.73	8.20	10.51	6.96	
	IV	<u>6.24</u>	<u>3.79</u>	<u>4.59</u>	<u>9.03</u>	<u>6.65</u>	<u>3.78</u>	
	Σ	31.94	25.02	33.50	90.46	31.55	24.71	90.90
	Σ	147.65	152.06	121.15	420.86	255.47	156.08	643.22

Appendix G-1. Habitat parameters at benthic-invertebrate sampling sites, June 15, 1983. Roman numeral represents stratum at site.

Middle Creek													

<sup>a</sup>Missing data.

Appendix G-2. Habitat parameters at benthic-invertebrate sampling sites, August 25, 1983. Roman numeral represents stratum at site.

<u>Middle Creek</u>												
	1				Site 2				5			
Time	1318				1210				1046			
Water temperature (°C)	8.8				8.7				7.8			
Stream width (m)	2.4				3.3				3.7			
Riparian habitat (%)												
Conifers	-				-				10			
Deciduous trees	-				-				40			
Shrubs/brush	-				-				30			
Grasses	100				100				20			
Benthos Collection Point	I	II	III	IV	I	II	III	IV	I	II	III	IV
Stream substrate composition (%)												
Boulder	-	-	-	-	-	-	-	-	-	80	50	50
Rubble	-	30	30	-	-	-	-	-	70	-	30	25
Gravel	80	70	70	100	100	100	100	100	30	20	20	25
Sand/silt	20	-	-	-	-	-	-	-	-	-	-	-
Water depth (m)	0.06	0.06	0.06	0.03	0.03	0.03	0.08	0.21	0.24	0.21	0.08	0.08
Water velocity (m/sec)	0.15	0.26	0.29	0.12	0.17	0.19	0.41	0.40	0.41	0.27	0.18	0.10

<u>Lone Creek</u>												
	7				Site 8				11			
Time	1455				1616				0910			
Water temperature (°C)	11.0				11.3				8.2			
Stream width (m)	4.3				5.2				4.9			
Riparian habitat (%)												
Conifers	-				-				10			
Deciduous trees	10				10				20			
Shrubs/brush	70				70				70			
Grass	20				20				-			
Benthos Collection Point	I	II	III	IV	I	II	III	IV	I	II	III	IV
Stream substrate composition (%)												
Boulder	50	-	30	-	-	-	-	-	-	-	-	-
Rubble	40	40	20	40	-	20	40	70	-	80	100	90
Gravel	-	60	50	60	90	70	60	30	100	20	-	-
Sand/silt	10	-	-	-	10	10	-	-	-	-	-	-
Water depth (m)	0.09	0.09	0.11	0.09	0.09	0.03	0.06	0.09	0.10	0.18	0.18	0.13
Water velocity (m/sec)	0.38	0.24	0.41	0.48	0.59	0.48	0.44	0.76	0.59	0.87	0.48	0.24

Appendix G-3. Habitat parameters at benthic-invertebrate sampling sites, June 14, 1984. Roman numeral represents stratum at site.

<u>Middle Creek</u>												
	1				Site 2				5			
Time	1145				1050				0944			
Water temperature (°C)	11.1				10.6				9.9			
Stream width (m)	2.4				3.0				4.9			
Riparian habitat (%)												
Conifers	-				-				10			
Deciduous trees	-				-				50			
Shrubs/brush	50				-				30			
Grasses	50				100				10			
Benthos Collection Point	I	II	III	IV	I	II	III	IV	I	II	III	IV
Stream substrate composition (%)												
Boulder	-	-	-	-	-	-	-	-	-	-	90	-
Rubble	40	60	30	50	-	-	100	40	80	90	10	70
Gravel	60	40	70	50	50	50	-	-	20	10	-	30
Sand/silt	-	-	-	-	50	50	-	60	-	-	-	-
Water depth (m)	0.08	0.08	0.08	0.06	0.15	0.14	0.08	0.05	0.27	0.15	0.14	0.09
Water velocity (m/sec)	0.34	0.30	0.40	0.49	0.34	0.12	0.37	0.24	0.43	0.55	0.21	0.12

<u>Lone Creek</u>												
	7				Site 8				11			
Time	1230				1330				0840			
Water temperature (°C)	12.0				12.4				10.2			
Stream width (m)	4.9				3.7				6.1			
Riparian habitat (%)												
Conifers	-				-				-			
Deciduous trees	-				20				80			
Shrubs/brush	70				70				20			
Grass	30				10				-			
Benthos Collection Point	I	II	III	IV	I	II	III	IV	I	II	III	IV
Stream substrate composition (%)												
Boulder	-	-	-	-	-	-	-	-	-	-	-	-
Rubble	80	90	70	-	50	50	-	40	40	90	100	90
Gravel	20	10	30	80	40	40	100	40	40	10	-	10
Sand/silt	-	-	-	20	10	10	-	20	20	-	-	-
Water depth (m)	0.08	0.11	0.09	0.18	0.15	0.18	0.18	0.14	0.18	0.18	0.24	0.12
Water velocity (m/sec)	0.34	0.21	0.55	0.34	0.52	0.58	0.43	0.34	0.43	0.52	0.61	0.43

Appendix G-4. Habitat parameters at benthic-invertebrate sampling sites, August 29, 1984. Roman numeral represents stratum at site.

<u>Middle Creek</u>												
1					Site 2				5			
Time	1450				1410				1300			
Water temperature (°C)	7.2				7.0				6.6			
Stream width (m)	2.4				3.7				4.9			
Benthos Collection Point	I	II	III	IV	I	II	III	IV	I	II	III	IV
Stream substrate composition (%)												
Boulder	-	-	-	-	-	-	-	-	-	80	100	-
Rubble	40	40	60	80	-	-	-	-	95	10	-	50
Gravel	60	50	40	20	90	100	100	80	5	10	-	40
Sand/silt	-	10	-	-	10	-	-	20	-	-	-	10
Water depth (m)	0.15	0.17	0.08	0.06	0.06	0.06	0.08	0.24	0.18	0.17	0.12	0.06
Water velocity (m/sec)	0.37	0.06	0.37	0.06	0.21	0.37	0.40	0.18	0.30	0.27	0.09	0.06

<u>Lane Creek</u>												
7					Site 8				11			
Time	0930				1040				1140			
Water temperature (°C)	5.5				6.4				6.4			
Stream width (m)	5.5				3.7				6.1			
Benthos Collection Point	I	II	III	IV	I	II	III	IV	I	II	III	IV
Stream substrate composition (%)												
Boulder	-	-	-	-	-	-	-	-	-	-	-	-
Rubble	90	100	80	80	50	60	40	-	70	100	90	70
Gravel	10	-	20	10	50	40	60	60	30	-	10	30
Sand/silt	-	-	-	10	-	-	-	40	-	-	-	-
Water depth (m)	0.09	0.09	0.09	0.14	0.06	0.15	0.14	0.12	0.08	0.09	0.18	0.06
Water velocity (m/sec)	0.24	0.15	0.43	0.24	0.30	0.37	0.37	0.18	0.34	0.61	0.64	0.15