

BRISTOL BAY–ALASKA PENINSULA BIBLIOGRAPHY OF SELECTED REFERENCES

- Addicott, W.O., 1971, Tertiary marine mollusks of Alaska; An annotated bibliography: U.S. Geological Survey Bulletin 1343, 30 p.
- Alaska Department of Natural Resources, 1971, Aeromagnetic map, Northwest Nushagak Bay Quadrangle: BBNC files, data set index no. 461, 1 sheet, scale 1:1,250,000.
- 1971, Aeromagnetic survey, Goodnews Bay Quadrangle: BBNC files, data set index no. 488, 1 sheet, scale 1:250,000.
- Alaska Department of Natural Resources and Bristol Bay Native Corporation, 2003, Memorandum of Understanding Between the Alaska Department of Natural Resources and Bristol Bay Native Corporation to Facilitate Oil and Gas Lease Sales on State and BBNC Land in the Bristol Bay Region, 3 p.
- Alaska Division of Geological & Geophysical Surveys staff, 1986, Summary of existing data and potential for commercial hydrocarbon accumulations, Bristol Bay, Alaska: Alaska Division of Geological & Geophysical Surveys Public-Data File 86-13, 20 p.
- Alaska Division of Oil & Gas, 2004, General information and digital well data—Alaska Peninsula areawide lease sale and Bristol Bay Basin exploration license area: Information and data Compilation, v. 1.
- 2004, Seismic data—Alaska Peninsula areawide lease sale and Bristol Bay basin exploration license area: Information and data compilation, v. 3.
- 2004, Scanned electric well logs—Alaska Peninsula areawide lease sale and Bristol Bay basin exploration license area: Information and data compilation, v. 2.
- 2004, Alaska Peninsula areawide oil and gas lease sale and Bristol Bay basin exploration license area, information and data compilation: Alaska Division of Oil & Gas, Alaska Peninsula oil and gas resource series, 3 volume CD-ROM.
- Alaska Oil and Gas Conservation Commission (AOGCC), 2004, Alaska exploratory wells list: <http://www.state.ak.us/local/akpages/ADMIN/ogc/homeogc.shtml>
- Albanese, M.D., and Goff, K.M., 1987, Petrified wood occurrences in Alaska: Alaska Division of Geological & Geophysical Surveys Public-Data File 87-26, 11 p.
- Allaway, W.H., Jr., 1982, Sedimentology and petrography of the Shelikof Formation, Alaska Peninsula: San Jose, CA, unpublished M.S. thesis, San Jose State University, 81 p.
- Allaway, W.H., Jr., and Miller, J.W., 1984, Newly recognized sedimentary environments in the Shelikof Formation, in Coonrad, W.L., and Elliot, R.L., eds., The United States Geological Survey in Alaska; Accomplishments during 1981: U.S. Geological Survey Circular 868, p. 34–37.
- Allaway, W.H., Jr., Detterman, R.L., Miller, J.W., and Magoon, L.B., 1984, Stratigraphic clarification of the Shelikof Formation, Alaska Peninsula, in Stratigraphic notes, 1983: U.S. Geological Survey Bulletin 1537-A, p. A21-A27.
- Allison, R.C., 1978, Late Oligocene through Pleistocene molluscan faunas in the Gulf of Alaska region: The Veliger, v. 21, p. 171–188.
- Allison, R.C., and Addicott, W.O., 1973, The *Mytilus-middendorffi* group (*Bivalvia*) of the North American Miocene [abs.]: Geological Society of America Abstracts with Programs, v. 5, p. 2–3.
- 1976, The North Pacific Miocene record of *Mytilus (Plicatomytilus)*, a new subgenus of *Bivalvia*: U.S. Geological Survey Professional Paper 962, 22 p.
- Amoco, 1979, Geologic map of the Alaska Peninsula: Amoco Production Company Geologic Report FR-02-79, (unpublished report of 1977 field season, available through Alaska Geologic Materials Center), scale 1 in = 16,000 ft (1:192,000).
- 1979, High sensitivity magnetometer survey of the Alaska Peninsula, upper half: BBNC files, data set index no. 703, 1 sheet, scale 1:96,000.
- 1979, High sensitivity magnetometer survey of the Alaska Peninsula, lower half: BBNC files, data set index no. 704, 1 sheet, scale 1:96,000.
- 1979, High sensitivity magnetometer survey of the Alaska Peninsula: BBNC files, data set index no. 771, 1 sheet, scale 1:96,000.
- 1980, Bouguer gravity map, Bristol Bay: BBNC files, data set index no. 705, 1 sheet, scale 1:250,000.
- 1982, Seismic map of shallow unconformity, east and southeast of Egegik Bay, Alaska Peninsula: BBNC files, data set index no. 688, 1 sheet, scale 1:48,000.

- Amoco, 1982, Seismic map of deep unconformity, east and southeast of Egegik Bay, Alaska Peninsula: BBNC files, data set index no. 687, 1 sheet, scale 1:48,000.
- 1982, ANM shotpoints; Bristol Bay, southern peninsula, Egegik Bay: BBNC files, data set index no. 689, 1 sheet, various scales.
- 1982, Alaska Peninsula, Bouguer gravity, density 2.67: BBNC files, data set index no. 702, 2 sheets, scale 1:250,000.
- 1983, Geologic map of the Staniukovich Mountain area: Amoco Production Company Geologic Report FW-18-81R, (unpublished report of 1981 field season, available through Alaska Geologic Materials Center), scale 1:63,360.
- 1983, Top of unconformity (time–structure); Bristol Bay, southern peninsula, Ugashik: BBNC files, data set index no. 686, 1 sheet, scale 1:48,000.
- 1983, Deep unconformity; Bristol Bay, southern peninsula, Egegik Bay: BBNC files, data set index no. 687, 1 sheet, scale 1:48,000.
- 1983, Shallow unconformity; Bristol Bay, southern peninsula, Egegik Bay: BBNC files, data set index no. 688, 1 sheet, scale 1:48,000.
- 1983, BBNC southern peninsula, seismic base for gravity and magnetics, base map 3: BBNC files, data set index no. 727, 1 sheet, scale 1:250,000.
- 1985, Post drilling appraisal of the Amoco Becharof State No. 1 well: BBNC files, data set index no. 623, 35 p.
- Atwood, W.W., 1911, Geology and mineral resources of parts of the Alaska Peninsula: U.S. Geological Survey Bulletin 467, 137 p.
- Barbacka, Maria, Pálffy, J., and Smith, P.L., 2006, Hettangian (Early Jurassic) plant fossils from Puale Bay (Peninsular terrane, Alaska): Review of Palaeobotany and Palynology, v. 142, p. 33–46.
- Barnes, I., and McCoy, G.A., 1979, Possible role of mantle-derived CO₂ in causing two “phreatic” explosions in Alaska: Geology, v. 7, p. 434–435.
- Bascle, R., Evans, D., Seidlitz, A., and Borkowske, J., 1987, Alaska Peninsula/Becharof National Wildlife Refuge oil and gas assessment: Bureau of Land Management, 83 p., http://www.blm.gov/pgdata/etc/medialib/blm/ak/aktest/energy/egy_pubs.Par.81720.File.dat/ak_peninsula_becharof_oga.pdf
- Beeman, W.R., Obuch, R.C., and Brewton, J.D., 1996, Digital map data, text, and graphical images in support of the 1995 National Assessment of United States Oil and Gas Resources: U.S. Geological Survey Digital Data Series DDS-35, 1 disk, <http://pubs.usgs.gov/dds/dds-035/DDS-35.html>
- Beikman, H.M., 1980, Geologic map of Alaska: U.S. Geological Survey Professional Paper 171, 1 plate, scale 1:2,500,000.
- Bergman, S.C., Murphy, John, and Kelley, Shari, this volume.
- Bishop, W.C., 1985, History, geology, energy, and mineral resources of Bristol Bay area: Bristol Bay Native Corporation, Internal Report, 145 p.
- Blackwell, D.D., and Richards, M., editors, 2004, Geothermal Map of North America: Tulsa, OK, American Association of Petroleum Geologists, 1 sheet, scale 1:6,500,000; <http://www.smu.edu/geothermal/2004NAMap/2004NAMap.htm>.
- Blasko, D.P., 1976, Oil and gas seeps in Alaska: Alaska Peninsula, western Gulf of Alaska: U.S. Department of the Interior, Bureau of Mines Report of Investigations no. 8122, 78 p.
- Blodgett, R.B., this volume.
- Blodgett, R.B., and Clautice, K.H., 2005, Oil and gas seeps of the Puale Bay–Becharof Lake–Wide Bay region, northern Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2005-6, 13 p., 1 sheet.
- Blodgett, R.B., and Frýda, J., 2001, On the occurrence of *Spinidelphinulopsis whaleni* (Late Triassic [early Norian] Gastropoda) in the Cornwallis Limestone, Kuiu Island, southeastern Alaska (Alexander terrane) and its paleobiogeographic significance: Bulletin of the Czech Geological Survey, v. 76, no. 4, p. 235–242.
- 2001, Upper Triassic gastropod biogeography of western North America [abs.]: Geological Society of America Abstracts with Programs, v. 33, no. 3, p. A-53.
- Blodgett, R.B., and Sralla, Bryan, 2008, A major unconformity between Permian and Triassic strata at Cape Kekurnoi, Alaska Peninsula; Old and new observations on stratigraphy and hydrocarbon potential, in Haeussler, P.J., and Galloway, J.P., eds., Studies by the U.S. Geological Survey in Alaska, 2006: U.S. Geological Survey Professional Paper 1739-E, 13 p. [<http://pubs.usgs.gov/pp/pp1739/e/>].

- Blodgett, R.B., Finzel, E.S., Reifenhohl, R.R., Clautice, K.H., Ridgway, K.D., and Gillis, R.J., 2008, Jurassic through Pliocene age megafossil samples collected in 2005 by the Alaska Division of Geological & Geophysical Surveys from the Bristol Bay–Port Moller area, Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2008-2, 12 p.
- Blodgett, R.B., Frýda, J., and Stanley, G.D., Jr., 2001, *Delphinulopsidae*, a new neritopsoid gastropod family from the Upper Triassic (upper Carnian or lower Norian) of the Wallowa terrane, northeastern Oregon: *Journal of Czech Geological Society*, v. 46, no. 3–4, p. 307–318.
- 2003, Late Triassic gastropod biogeography of southern Alaskan accreted terranes [abs.]: *Geological Society of America Abstracts with Programs*, v. 35, no. 4, p. 59.
- 2005, Upper Triassic gastropod fauna from southern Alaska and its implications for terrane accretion [abs.]: *Geological Society of America Abstracts with Programs*, v. 37, no. 7, p. 82.
- 2006, Late Triassic gastropod faunas of western North America—A useful tool in terrane analysis [abs.]: *Geological Society of America Abstracts with Programs*, v. 38, no. 5, p. 81.
- Blodgett, R.B., Weems, R.E., and Wilson, F.H., 1995, Upper Jurassic reptiles from the Naknek Formation, Alaska Peninsula; A glimpse into Alaska's own "Jurassic Park": *Geological Society of America, Abstracts with Programs*, v. 27, no. 5, p. 6.
- Blodgett, R.B., Wilson, F.H., Stanley, G.D., Jr., McRoberts, C.A., and Sandy, M.R., 2000, Upper Triassic stratigraphy and fauna of the Taylor Mountains D-2 and D-3 quadrangles (SW part of the Farewell terrane), southwest Alaska [abs.]: *Geological Society of America Abstracts with Programs*, v. 32, no. 6, p. A-4.
- Bolger, G.W., and Reifenhohl, R.R., this volume.
- Bond, G.C., Lewis, S.D., Taber, J., Steckler, M.S., and Kominz, M.A., 1988, Evidence for formation of a flexural back arc basin by compression and crustal thickening in the central Alaska Peninsula: *Geology*, v. 16, p. 1,147–1,150.
- Boreham, C.J., and Powell, T.G., 1993, Petroleum source rock potential of coal and associated sediments—qualitative and quantitative aspects, in Law, B.E., and Rice, D.D., eds., *Hydrocarbons from coal: American Association of Petroleum Geologists Studies in Geology* 38, p. 133–157.
- Bristol Bay Native Corporation, 2003, Regional Land Status Map: Bristol Bay Native Corporation, Bristol Environmental & Engineering Services Corporation, 1 sheet.
- 2004, Bristol Bay Native Corporation Oil and Gas Files: 5 p.
- Brockway, Ronald, Alexander, B., Day, P., Lyle, W.M., Hiles, R., Decker, W., Polski, W., and Reed, B.L., 1975, Bristol Bay region, stratigraphic correlation section, southwest Alaska: Anchorage, The Alaska Geological Society, 1 sheet.
- Burk, C.A., 1965, *Geology of the Alaska Peninsula—Island Arc and Continental Margin*: The Geological Society of America Memoir 99, 250 p., 3 sheets.
- Capps, S.R., 1921, Preliminary report on petroleum in Alaska: *U.S. Geological Survey Bulletin* 719, 83 p.
- 1923, The Cold Bay district, in *U.S. Geological Survey Staff, Mineral resources of Alaska, report on progress of investigations in 1921*: *U.S. Geological Survey Bulletin* 739, p. 77–116, 1 plate, scale 1:250,000.
- 1937, Kodiak and vicinity, Alaska: *U.S. Geological Survey Bulletin* 868-B, p. 93–134, 2 plates, scale 1:250,000.
- Carlson, W.D., 1990, Mechanisms and kinetics of apatite fission track annealing: *American Mineralogist*, v. 75, p. 1,120–1,139.
- Carlson, W.D., Donelick, R.A., and Ketcham, R.A., 1999, Variability of apatite fission track annealing kinetics I: Experimental results: *American Mineralogist*, v. 84, p. 1,213–1,223.
- Case, J.E., Detterman, R.L., Wilson, F.H., Chuchel, B.A., and Yount, M.E., 1988, Maps showing aeromagnetic survey and geologic interpretation of parts of the Ugashik and Karluk Quadrangles, Alaska: *U.S. Geological Survey, Miscellaneous Field Studies Map* 1539-D, 12 p., 2 plates, scale 1:250,000.
- Citron, G.P., Kay, R.W., Mahlburg Kay, S., Snee, L.W., and Sutter, J.F., 1980, Tectonic significance of early Oligocene plutonism on Adak Island, central Aleutian Islands, Alaska: *Geology*, v. 8, p. 375–379.
- Clayton, J.L., 1993, Composition of crude oils generated from coals and coaly organic matter in shales, in Law, B.E., and Rice, D.D., eds., *Hydrocarbons from coal: American Association of Petroleum Geologists Studies in Geology* 38, p. 185–201.
- Clegg, P., and Holdsworth, R.E., 2005, Complex deformation as a result of strain partitioning in transpression zones; an example from the Leinster Terrane, SE Ireland: *Journal of the Geological Society*, v. 16.2, p. 187–202.
- Comer, C.D., and Martin, G.C., 1994, Petroleum potential of Mesozoic strata of the Cook Inlet outer continental shelf planning area, Alaska, *Third Symposium on Studies Related to Continental Margins—A Summary of Year Five and Year Six Activities*: Austin, TX, p. 172–180.

- Connelly, William, and Moore, J.C., 1979, Geologic map of the northwest side of the Kodiak Islands, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map 1057, 2 plates, scales 1:250,000 and 1:63,360. [See USGS Open-File Report 77-382]
- Conwell, C.N., and Triplehorn, D.M., 1978, Herendeen Bay–Chignik coals, southern Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Special Report 8, 15 p., 2 sheets.
- Cronin, M.A., Brannon, Ernie, and Moulton, Larry, 1999, Natural oil seeps in the Puale Bay area of the Becharof National Wildlife Refuge, and the Blue Creek area of the Alaska Peninsula National Wildlife Refuge, Alaska—A report to Refuge Manager, Becharof National Wildlife Refuge (dated 27 September 1999).
- Crowley, K.D., 1985, Thermal significance of fission track length distributions: *Nuclear Tracks*, v. 10, p. 311–322.
- Csejtey, B., Jr., and St. Aubin, D.R., 1981, Evidence for northwestward thrusting of the Talkeetna Superterrane, and its regional significance, *in* Albert, N.R.D., and Hudson, T., eds., 1981, *The U.S. Geological Survey in Alaska; Accomplishments during 1979: U.S. Geological Survey Circular 823-B*, p. B49–B51.
- Dall, W.H., 1882, Notes on Alaska Cenozoic deposits: *American Journal of Science*, 3rd Series, v. 24, p. 67–68.
- Decker, P.L., this volume.
- Decker, J., Bergman, S.C., Blodgett, R.B., Box, S.E., Bundtzen, T.K., Clough, J.G., Coonrad, W.L., Gilbert, W.G., Miller, M.L., Murphy, J.M., Robinson, M.S., and Wallace, W.K., 1994, Geology of southwestern Alaska, *in* Plafker, G., and Berg, H.C., eds., *The geology of Alaska: Boulder, CO, Geological Society of America, The Geology of North America*, v. G1, p. 285–310.
- Decker, P.L., Finzel, E.S., Ridgway, K.D., Reifentstuhl, R.R., and Blodgett, R.B., 2005, Preliminary summary of the 2005 field season; Port Moller, Herendeen Bay, and Dillingham areas, Bristol Bay Basin, Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2005-7, 55 p., 2 sheets.
- Decker, P.L., Reifentstuhl, R.R., Finzel, E.S., and Helmold, K.P., 2006, Play concepts for reopening the Bristol Bay basin; Tertiary and Mesozoic petroleum systems of the Alaska Peninsula, (abs): *American Association of Petroleum Geologists Bulletin*, v. 90, Program Abstracts (digital).
- Decker, P.L., Reifentstuhl, R.R., and Gillis, R.J., this volume.
- Demaison, G., and Huizinga, B.J., 1994, Genetic classification of petroleum systems using three factors; Charge, migration and entrapment, *in* Magoon, L.B., and Dow, W.G., eds., *The petroleum system—from source to trap: American Association of Petroleum Geologists Memoir 60*, p.73–89.
- DeMets, Charles, Gordon, R.G., Argus, D.F., and Stein, Seth, 1994, Effects of recent revisions to the geomagnetic reversal time scale on estimates of current plate motions: *Geophysical Research Letters*, v. 21, p. 2,191–2,194.
- Detterman, R.L., 1978, Interpretation of depositional environments in the Chignik Formation, Alaska Peninsula, *in* Johnson, K.M., ed., *The United States Geological Survey in Alaska; Accomplishments during 1977: U.S. Geological Survey Circular 772-B*, p. B62–B63.
- 1990, Stratigraphic correlation and interpretation of exploratory wells, Alaska Peninsula: U.S. Geological Survey Open-File Report 90-279, 51 p.
- Detterman, R.L., and Hartssock, J.K., 1966, Geology of the Iniskin–Tuxedni region, Alaska: U.S. Geological Survey Professional Paper 512, 78 p., 6 plates, scale 1:63,360.
- Detterman, R.L., and Miller, R.L., 1985, Kaguyak Formation—An Upper Cretaceous flysch deposit, *in* Bartsch-Winkler, Susan, and Reed, K.M., eds., *United States Geological Survey in Alaska: Accomplishments during 1983: U.S. Geological Survey Circular 945*, p. 49–51.
- Detterman, R.L., and Reed, B.L., 1980, Stratigraphy, structure, and economic geology of the Iliamna Quadrangle, Alaska: U.S. Geological Survey Bulletin 1368-B, 86 p., 1 plate, scale 1:250,000.
- Detterman, R.L., Case, J.E., Church, Stanley E., Frisken, J.G., Wilson, F.H., and Yount, M.E., 1990, The Alaska Mineral Resource Assessment Program (AMRAP); background information to accompany folio of geologic and resource maps of the Ugashik, Bristol Bay, and western part of Karluk quadrangles, Alaska, U.S. Geological Survey Circular 1046, 14 p.
- Detterman, R.L., Case, J.E., Miller, J.W., Wilson, F.H., and Yount, M.E., 1996, Stratigraphic framework of the Alaska Peninsula: U.S. Geological Survey Bulletin 1969-A, 74 p.
- Detterman, R.L., Case, J.E., Wilson, F.H., and Yount, M.E., 1987, Geologic map of the Ugashik, Bristol Bay, and western part of Karluk quadrangles, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map 1685, 1 plate, scale 1:250,000.

- Detterman, R.L., Case, J.E., Wilson, F.H., Yount, M.E., and Allaway, W.H., Jr., 1983, Generalized geologic map of the Ugashik, Bristol Bay, and part of Karluk quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map 1539-A, 1 plate, scale 1:250,000.
- Detterman, R.L., Hudson, Travis, Plafker, George, Tysdal, R.G., and Hoare, J.M., 1976, Reconnaissance geologic map along Bruin Bay and Lake Clark faults in Kenai and Tyonek quadrangles, Alaska: U.S. Geological Survey Open-File Map 76-477, 4 p., 1 plate, scale 1:250,000.
- Detterman, R.L., Miller, J.W., and Case, J.E., 1985, Megafossil locality map, checklists, and pre-Quaternary stratigraphic sections of Ugashik, Bristol Bay, and part of Karluk quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map 1539-B, 3 plates, scale 1:250,000.
- Detterman, R.L., Miller, T.P., Yount, M.E., and Wilson, F.H., 1979, Generalized geologic map of the Chignik and Sutwik Island quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map 1053-A, 1 plate, scale 1:250,000.
- Detterman, R.L., Miller, T.P., Yount, M.E., and Wilson, F.H., 1981, Geologic map of the Chignik and Sutwik Island quadrangles, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map 1229, 1 plate, scale 1:250,000.
- 1981, Quaternary geologic map of the Chignik and Sutwik Island quadrangles, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map 1292, 1 plate, scale 1:250,000.
- Detterman, R.L., Wilson, F.H., Yount, M.E., and Miller, T.P., 1987, Quaternary geologic map of the Ugashik, Bristol Bay, and western part of the Karluk quadrangles, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map 1801, 1 plate, scale 1:250,000.
- Detterman, R.L., Yount, M.E., and Case, J.E., 1981, Megafossil sample locality map, checklists, and stratigraphic sections of the Chignik and Sutwik Island quadrangles, Alaska: U.S. Geological Survey Miscellaneous field Studies Map MF-1053-N, 2 plates, scale 1:250,000.
- Dickinson, W.R., 1970, Interpreting detrital modes of graywacke and arkose, *Journal of Sedimentary Petrology*, v. 40, p. 695–707.
- Dickinson, W.R. and Suczek, C.A., 1979, Plate tectonics and sandstone compositions: *American Association of Petroleum Geologists*, v. 63, p. 2,164–2,182.
- Dickson, J.A.D., 1965, A modified staining technique for carbonates in thin section: *Nature*, v. 205, p.587.
- 1966, Carbonate identification and genesis as revealed by staining: *Journal of Sedimentary Petrology*, v. 36, p. 491–505.
- Dodson, M.H., and McClelland-Brown, E., 1985, Isotopic and paleomagnetic evidence for rates of cooling, uplift, and erosion, *in* Snelling, N.J., ed., *The chronology of the geological record*: Geological Society of London, Memoir 10, p. 315–325.
- Donelick, R.A., Ketcham, R.A., and Carlson, W.D., 1999, Variability of apatite fission-track annealing kinetics; II, Crystallographic orientation effects: *American Mineralogist*, v. 84, p. 1,224–1,234.
- Donelick, R.A., O'Sullivan, P.B., and Ketcham, R.A., 2005, Apatite fission-track analysis: *Reviews in Mineralogy and Geochemistry*, v. 58, no. 1, p. 49–94.
- Dow, W.G., 1977, Kerogen studies and geological interpretations: *Journal of Geochemical Exploration*, v. 7, p. 79–99.
- 1977, Petroleum source beds on continental slopes and rises, *in* *Geology of continental margins*: American Association of Petroleum Geologists Short Course Note Series No. 6, p. D1–D37.
- 1983, Geochemical analysis of North Aleutian Shelf Cost No. 1 well, Alaska: U.S. Minerals Management Service public well files – Robertson Research Report No. 823/135, 315 p.
- Dubois, G.D., Wilson, F.H., and Shew, N.B., 1987, Potassium-argon age determinations from the Port Moller and Stepovak Bay quadrangles, Alaska Peninsula: U.S. Geological Survey Open-File Report 87-191, 1 plate, scale 1:250,000.
- Dumoulin, J.A., Whalen, M.T., and Harris, A.G., 2008, Lithofacies, age, and sequence stratigraphy of the Carboniferous Lisburne Group in the Skimo Creek area, central Brooks Range, *in* Haeussler, Peter J., and Galloway, John P., eds., *Studies by the U.S. Geological Survey in Alaska, 2006*: U.S. Geological Survey Professional Paper 1739-B, 64 p. [<http://pubs.usgs.gov/pp/pp1739/b/>].
- Dunham, R.J., 1962, Classification of carbonate rocks according to texture, *in* Ham, W.E., ed., *Classification of carbonate rocks*: American Association of Petroleum Geologists Memoir 1, p. 108–121.
- Eakins, G.R., 1970, A petrified forest on Unga Island, Alaska, Alaska Division of Mines and Geology, Special Report 3, 19 p.

- Eckdale, A.A., Bromley, R.G., and Pemberton, S.G., 1984, Ichnology trace fossils in sedimentology and stratigraphy: Society of Economic Paleontologists and Mineralogists Short Course 15, 317 p.
- Elder, W.P., and Miller, J.W., 1991, Maps showing fossil localities and checklists of Jurassic and Cretaceous macrofauna of western Alaska: U.S. Geological Survey Open-File Report 91-629, 71 p., 3 plates, scale 1:500,000.
- Embry, A.F., and Klovan, J.E., 1972, Absolute water depth limits of late Devonian paleoecological zones: Geological Research, v. 61, p. 672–686.
- Epstein, A.G., Epstein, J.B., and Harris, L.D., 1977, Conodont color alteration—An index to organic metamorphism: U.S. Geological Survey Professional Paper 995, 27 p.
- Espitalié, J., Deroo, G., and Marquis, F., 1985, La pyrolyse Rock-Eval et ses applications: Revue de l'Institut Français du Pétrole, v. 40, p. 563–579 and 755–784.
- Espitalié, J., Laporte, J.L., Madec, M., Marquis, F., Leplat, P., Paulet, J., Boutefeu, A., 1977, Methode rapide de caracterisation des roches meres, de leur potential petrolier et de leur degre d'evolution: Revue de l'Institut Français du Pétrole, v. 32, p. 23–42.
- Espitalié, J., Madec, M., and Tissot, B., 1977, Source rock characterization method for petroleum exploration: Proceedings of the 9th Annual Offshore Technology Conference, v. 3, p. 439–448.
- Fairchild, D.T., 1977, Paleoenvironments of the Chignik Formation, Alaska Peninsula: Fairbanks, Alaska, University of Alaska Fairbanks, M.S. thesis, 168 p.
- Finzel, E.S., Reifenhohl, R.R., Decker, P.L., and Ridgway, K.D., 2005, Sedimentology, stratigraphy, and hydrocarbon reservoir-source rock potential, using surface and subsurface data, of Tertiary and Mesozoic strata, Bristol Bay Basin and Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2005-4, 67 p.
- Finzel, E.S., Ridgway, K.D., Reifenhohl, R.R., Blodgett, R.B., White, J., and Decker, P.L., in press, Stratigraphic framework and estuarine depositional environments of the Miocene Bear Lake Formation, Bristol Bay basin, Alaska: Onshore equivalents to potential reservoir strata in a frontier gas-rich basin: American Association of Petroleum Geologists Bulletin.
- Fiorillo, A.R., Website showing hadrosaur footprint from Cretaceous rocks at Aniakchak crater: <http://www.uaf.edu/seagrant/NewsMedia/01ASJ/09.21.01dinosaur.html> (accessed August 20, 2008).
- Fischer, P.H., 1872, Sur quelques fossiles de l'Alaska, rapportés par. M.A. Pinart: Comptes Rendus, v. 75, p. 1,784–1,786.
- 1875, Sur quelques fossiles de l'Alaska, in Pinard, A.L., Voyages à la côte nord-ouest de l'Amérique exécutés dans les années 1870–1872: Paris, E. Leroux, pt. 1, p. 33–36.
- Fleet, A.J., and Scott, A.C., 1994, Coal and coal-bearing strata as oil-prone source rocks; An overview, in Scott, A.C., and Fleet, A.J., eds., Coal and coal-bearing strata as oil-prone source rocks?: Geological Society Special Publication No. 77, p. 1–8.
- Fleischer, R.L., Price, P.B., and Walker, R.M., 1975, Nuclear tracks in solids: Los Angeles, University of California Press, 605 p.
- Flett, T.O., 1988, Geothermal gradient, in Turner, R.F., ed., Geological and operational summary, North Aleutian Shelf COST No. 1 well, Bering Sea, Alaska: U.S. Minerals Management Service, OCS Report MMS 88-0089, p. 180–183.
- 1988, Organic geochemistry, in Turner, R.F., ed., Geological and operational summary, North Aleutian Shelf COST No. 1 well, Bering Sea, Alaska: U.S. Minerals Management Service, OCS Report MMS 88-0089, p. 184–202.
- Flügel, E., 2004, Microfacies of carbonate rocks; Analysis, interpretation, and application: Berlin-Heidelberg, Springer-Verlag, 976 p.
- Franks, S.G., and Hite, D.M., 1980, Controls on zeolite cementation in Upper Jurassic sandstones, lower Cook Inlet, Alaska: AAPG Bulletin, v. 64, no. 5, p. 708–709.
- Frech, Fritz, 1908, Die zircumpacifische Trias; Lethaea geognostica, Teil 2, Das Mesozoicum: Band 1, Trias, v. 4, p. 488–509.
- Fryda, J., and Blodgett, R. B., 2001, *Chulitmacula*, a new paleobiogeographically distinctive gastropod genus from Upper Triassic strata in accreted terranes of southern Alaska: Journal of Czech Geological Society, v. 46, no. 3–4, p. 299–306.
- Gabb, W.M., 1864, Description of Triassic fossils of California and the adjacent territories: California Geological Survey, Paleontology, v. 1, p. 17–35, plates 3–6.
- Gallagher, K., Brown, R.W., and Johnson, C., 1998, Fission track analysis and its applications to geological problems: Annual Reviews of Earth and Planetary Sciences, v. 26, p. 519–572.

- Galloway, W.E., 1974, Deposition and diagenetic alteration of sandstone in northeast Pacific arc-related basins; Implications for graywacke diagenesis: Geological Society of America Bulletin, v. 85, no. 3, p. 379–390.
- Gillis, R.J., Reifenhstahl, R.R., and Decker, P.L., 2008, Implications of new apatite and zircon fission-track thermochronology for Mesozoic and Tertiary basin margin exhumation, upper Alaska Peninsula [abs.], in Garver, J.I., and Montario, M.J., eds., Extended Abstract Volume: FT 2008 – The 11th International Conference on Thermochronometry, Anchorage Alaska, Sept. 15–19, 2008, p. 95, http://www.union.edu/ft2008/Abstract_volume.html.
- Gillis, R.J., Reifenhstahl, R.R., Decker, P.L., Strauch, A.L., and Helmold, K.P., 2007, Onshore Alaska Peninsula petroleum system assessment: preliminary summary of outcrop studies of the Port Moller area, 2006 [abs]: American Association of Petroleum Geologists Bulletin, v. 91, Annual Meeting Program Abstracts (digital), http://www.searchanddiscovery.net/documents/2007/07018annual_abs_ingbch/abstracts/lbGillis.htm.
- Gleadow, A.J.W., and Duddy, I.R., 1981, A natural long term annealing experiment for apatite: Nuclear Tracks, v. 5, p. 169–174.
- Gleadow, A.J.W., Duddy, I.R., and Lovering, J.F., 1983. Fission track analysis; A new tool for the evaluation of thermal histories and hydrocarbon potential: Australian Petroleum Association Journal, v. 23, p. 93–102.
- Gluyas, J.G., 1984, Early carbonate diagenesis within Phanerozoic shales and sandstones of the northwest European shelf: Clay Minerals, v. 19, p. 309–321.
- Gradstein, F.M., Ogg, J.G., Smith, A.G., Agterberg, F.P., Bleeker, W., Cooper, R.A., Davydov, V., Gibbard, P., Hinnov, L.A., House, M.R., Lourens, L., Luterbacher, H.P., McArthur, J., Melchin, M.J., Robb, L.J., Shergold, J., Villeneuve, M., Wardlaw, B.R., Ali, J., Brinkhuis, H., Hilgen, F.J., Hooker, J., Howarth, R.J., Knoll, A.H., Laskar, J., Monechi, S., Plumb, K.A., Powell, J., Raffi, I., Röhl, U., Sadler, P., Sanfilippo, A., Schmitz, B., Shackleton, N.J., Shields, G.A., Strauss, H., Van Dam, J., van Kolfschoten, T., Veizer, J., and Wilson, D., 2004, A geologic time scale 2004: Cambridge University Press, 589 p.
- Grant-Mackie, J.A., and Silberling, N.J., 1990, New data on the Upper Triassic bivalve *Monotis* in North America and the new subgenus *Pacimonotis*: Journal of Paleontology, v. 64, no. 2, p. 240–254.
- Green, P.F., Duddy, I.R., Gleadow, A.J.W., and Lovering, J.F., 1989, Apatite fission track analysis as a paleotemperature indicator for hydrocarbon exploration, in Naeser, N.D., and McCulloch, T.H., eds., Thermal history of sedimentary basins; methods and case histories: New York, Springer-Verlag, p. 181–195.
- Green, P.F., Duddy, I.R., Laslett, G.M., Hegarty, K.A., Gleadow, A.J.W., and Lovering, J.F., 1989, Thermal annealing of fission tracks in apatite; 4, Quantitative modeling techniques and extension to geological timescales: Chemical Geology (Isotope Geoscience Section), v. 79, no. 2, p. 155–182.
- Gulf Oil, 1962, Total magnetic contours, Port Moller area: BBNC files, data set index no. 695, 1 sheet, scale 1:8,000.
- 1962, Gravity meter survey, Port Moller–Port Heiden area, Alaska: BBNC files, data set index no.693, 1 sheet, scale 1:8,000.
- 1964, Port Moller project; Fireweed and Sandy River area, Deep Phantom Horizon #2: BBNC files, data set index no. 694, 1 sheet, scale 1:8,000.
- 1969, Port Moller project, Fireweed and Sandy River area, Deep Phantom Horizon # 2: Interpretive seismic map, BBNC files, data set index no. 694, 1 sheet, scale 1:8,000.
- Hanna, G.D., Leach, C.E., Tallant, R.L., Bryan, J.J., and Scott, E.W., 1937, Preliminary geologic report on the Bear Creek Anticline, Alaska—1937: San Francisco, Tide Water Associated Oil Co.–Standard Oil Co. of California–Union Oil Co. of California internal report, 20 p.
- Hanson, B.M., 1957, Middle Permian limestone on Pacific side of Alaska Peninsula: American Association of Petroleum Geologists Bulletin, v. 41, no. 10, p. 2,376–2,378.
- Hazzard, J.C., Borax, E., Bryan, J.J., and Shoemaker, R.W., 1950, Cretaceous rocks in the Kamishak Bay area, Cook Inlet, Alaska: Science, v. 112, no. 2904, p. 226–227.
- Helmold, K.P., and Brizzolara, D.W., 2005, Reservoir quality of Tertiary sandstones from Bristol Bay basin, Alaska Peninsula: preliminary report: Alaska Division of Oil and Gas internal report; http://www.dog.dnr.state.ak.us/oil/products/publications/akpeninsula/reservoir_quality_report.pdf
- Helmold, K.P., Brizzolara, D.W., and Reifenhstahl, R.R., this volume.
- Hill, M.D., 1979, Volcanic and plutonic rocks of the Kodiak–Shumagin shelf, Alaska; Subduction deposits and near-trench magmatism: Santa Cruz, CA, University of California Santa Cruz, Ph.D. Dissertation, 259 p.
- Hite, David, 2004, Bristol Bay Native Corporation—Hydrocarbon potential of Native lands in the Bristol Bay area, southern Alaska: BBNC files, Data Set Index No. 774.

- Hite, David, 2004, Hydrocarbon potential of the Alaska Peninsula and Bristol Bay basin, southern Alaska, - Emphasis on Bristol Bay Native Corporation lands, 8 p.
- Hoare, H.J., and Coonrad, W.L., 1961, Geologic map of the Hagemeister Island Quadrangle, Alaska: U.S. Geological Survey Miscellaneous Geologic Investigations, Map 321, 1 plate, scale 1:250,000.
- 1961, Geologic map of the Goodnews Quadrangle, Alaska: U.S. Geological Survey Miscellaneous Geologic Investigations, Map 339, 1 plate, scale 1:250,000.
- 1978, Geologic Map of the Goodnews and Hagemeister Island Quadrangles region, southwestern Alaska: U.S. Geological Survey Open-File Report 78-9B, 2 plates, scale 1:250,000.
- Hollick, A., 1930, The Upper Cretaceous floras of Alaska: U.S. Geological Survey Professional Paper 182, 185 p.
- Hoose, P.J., and Whitney, John, 1980, Map showing selected geologic features of the outer continental shelf, Shelikof Strait, Alaska: U.S. Geological Survey Open-File Report 80-2035, 1 plate, scale 1:250,000.
- Hunt, J.M., 1991, Generation of gas and oil from coal and other terrestrial organic matter: *Organic Geochemistry*, v. 17, p. 673–680.
- Imlay, R.W., 1953, Callovian (Jurassic) ammonites from the United States and Alaska, Part 2. Alaska Peninsula and Cook Inlet regions: U.S. Geological Survey Professional Paper 249-B, p. 41–108.
- 1975, Stratigraphic distribution and zonation of Jurassic (Callovian) ammonites in southern Alaska: U.S. Geological Survey Professional Paper 836, 28 p.
- 1981, Early Jurassic ammonites from Alaska: U.S. Geological Survey Professional Paper 1148, 49 p.
- 1981, Early Jurassic ammonites from Alaska: U.S. Geological Survey Professional Paper 1190, 40 p.
- 1984, Early and middle Bajocian (Middle Jurassic) ammonites from southern Alaska: U.S. Geological Survey Professional Paper 1322, 38 p.
- Imlay, R.W., and Detterman, R.L., 1977, Some Lower and Middle Jurassic beds in Puale Bay–Alinchak Bay area, Alaska Peninsula: *Bulletin of the American Association of Petroleum Geologists*, v. 61, no. 4, p. 607–611.
- Iriondo, Alexander, Kunk, M.J., and Wilson, F.H., 2003, ⁴⁰Ar/³⁹Ar geochronology of igneous rocks in the Taylor Mountains and Dillingham quadrangles in southwestern Alaska: U.S. Geological Survey Open-File Report 03-0421, 32 p.
- Jeffords, R.M., 1957, Permian fossils from an island in Puale Bay, Alaska: Houston, Humble Oil & Refining Co. Geologic Research Report 57-24, 6 p.
- Jones, D.L., 1973, Structural elements and biostratigraphic framework of Lower Cretaceous rocks in southern Alaska, *in* *The Boreal Lower Cretaceous: Liverpool, England*, Seel House Press, p. 1–18.
- Jones, D.L., and Detterman, R.L., 1966, Cretaceous stratigraphy of the Kamishak Hills, Alaska Peninsula, *in* *Geological Survey research 1966, Chapter D*: U.S. Geological Survey Professional Paper 550-D, p. D53–D58.
- Jones, D.L., and Miller, J.W., 1976, Preliminary geologic map of the Alaska Peninsula showing post-Callovian Mesozoic fossil localities: U.S. Geological Survey Open-File Report 76-76, 2 plates, scale 1:500,000.
- Jones, D.L., and Silberling, N.J., 1979, Mesozoic stratigraphy—the key to tectonic analysis of southern and central Alaska: U.S. Geological Survey Open-File Report 79-1200, 37 p.
- Jones, D.L., Silberling, N.J., and Hillhouse, John, 1977, Wrangellia—A displaced terrane in northwestern North America: *Canadian Journal of Earth Science*, v. 14, p. 2565–2577.
- Jones, R.R., and Tanner, P.W.G., 1995, Strain partitioning in transpression zones: *Journal of Structural Geology*, v. 17, p. 793–802.
- Katz, B.J., 1983, Limitations of ‘Rock-Eval’ pyrolysis for typing organic matter: *Organic Geochemistry*, v. 4, p. 195–199.
- Keller, A.S., and Reiser, H.N., 1959, Geology of the Mt. Katmai area, Alaska: U.S. Geological Survey Bulletin 1058-G, p. 261–298, 1 plate, scale 1:250,000.
- Kelley, J.S., 1980, Environments of deposition and petrography of Lower Jurassic volcanoclastic rocks, southwestern Kenai Peninsula, Alaska: Davis, University of California, Ph.D. thesis, 304 p.
- Kelley, J.S., and Denman, J.M., 1972, Geological literature on the Alaska Peninsula and adjacent area: Alaska Division of Geological & Geophysical Surveys Special Report 20, 64 p.
- Kellum, L.B., 1945, Jurassic stratigraphy of Alaska and petroleum exploration in northwest America: *Transactions of the New York Academy of Science*, ser. 2, v. 7, no. 8, p. 201–209.
- Kellum, L.B., Daviess, S.N., and Swinney, C.M., 1945, Geology and oil possibilities of the southeastern part of the Wide Bay anticline, Alaska, U.S. Geological Survey Open-File Report, 17 p.

- Kennedy, G.C., and Waldron, H.H., 1955, Geology of Pavlof volcano and vicinity, Alaska: Bulletin 1028-A, p. 1–19, scale 1:100,000.
- Kirschner, C.L., 1988, Map showing sedimentary basins of offshore and continental shelf areas, Alaska: U.S. Geological Survey Miscellaneous Investigations Series 1873, 1 plate, scale 1:2,500,000.
- Knappen, R.S., 1929, Geology and mineral resources of the Aniakchak district: U.S. Geological Survey Bulletin 797, p. 161–223.
- Kummel, Bernhard, 1953, American Triassic coiled nautiloids: U.S. Geological Survey Professional Paper 250, 104 p., 19 plates.
- Langford, F.F., and Blanc-Valleron, M.-M., 1990, Interpreting Rock-Eval pyrolysis data using graphs of pyrolyzable hydrocarbons vs. total organic carbon: American Association of Petroleum Geologists Bulletin, v. 74, p. 799–804.
- Laniz, R.V., Stevens, R.E., and Norman, M.B., 1964, Staining of plagioclase feldspar and other minerals: U.S. Geological Survey Professional Paper 501-B, p. B152–B153.
- Lankford, S.M., and Magoon, L.B., 1977, Petrography of the Upper Jurassic through Oligocene sandstones in the Cape Douglas–Kamishak Hills area, lower Cook Inlet in Johnson, K.M., ed., The United States Geological Survey in Alaska, Accomplishments during 1977, U.S. Geological Survey Circular 772-B, p. B60–B62.
- Larson, J.A., 1988, Paleontology and Biostratigraphy, in Turner, R.F., ed., Geological and operational summary, North Aleutian Shelf COST No. 1 well, Bering Sea, Alaska: U.S. Minerals Management Service, OCS Report MMS 88-0089, p. 159–179.
- Laslett, G.M., Green, P.F., Duddy, I.R., and Gleadow, A.J.W., 1987, Thermal modeling of fission tracks in apatite: 2. A quantitative analysis: Chemical Geology, v. 65, p. 1–13.
- Levinson, S.A., and Jeffords, R.M., 1956, Megafossils and spores and pollen from Puale Bay, Alinchak Bay, and Cape Kekurnoi, Alaska: Houston, Humble Oil & Refining Co. Geologic Research Report, 10 p., 4 plates.
- Lindholm, R.C., and Finkelman, R.B., 1972, Calcite staining; Semiquantitative determination of ferrous iron: Journal of Sedimentary Petrology, v. 42, p. 239–242.
- Loveland, A.M., Reifenhuhl, R.R., Gillis, R.J., and Decker, P.L., 2007, Outcrop sample results from mercury injection capillary pressure analyses, Bristol Bay, Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Raw Data File 2007-3, 11 p.
- Lyle, W.M., and Dobey, P.M., 1973, Geologic and mineral evaluation of the Aniakchak River drainage, Alaska Peninsula, for wild and scenic river study: Alaska Division of Geological & Geophysical Surveys Open-File Report 26, 21 p., 1 sheet, scale 1:63,360.
- 1974, Geologic evaluation of the Herendeen Bay area, Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Open-File Report 48, 22 p., 3 sheets.
- Lyle, W.M., Morehouse, J.A., Palmer, I.F., Jr., and Bolm, J.G., 1979, Tertiary formations and associated Mesozoic rocks in the Alaska Peninsula area, Alaska, and their petroleum reservoir and source-rock potential: Alaska Division of Geological & Geophysical Surveys Geologic Report 62, 69 p., 19 sheets.
- Mackevett, E.M., Jr., and Plafker, G., 1974, The Border Ranges Fault in southcentral Alaska: U.S. Geological Survey Journal of Research, v. 2, no. 3, p. 323–329.
- MacNeil, F.S., 1970, New Pliocene *Chlamys* (*Swiftopecten*) and *Beringius* from the Alaska Peninsula: Nautilus, v. 84, p. 69–74.
- 1973, Marine fossils from the Unga Conglomerate Member of the Bear Lake Formation, Cape Aliaksin, Alaska Peninsula, Alaska: Science Reports of the Tohoku University, Series 2, Geology Tohoku Daigaku Rika Hokoku, Dai 2: Shu Chishitsugaku v. 6, no. 6, p. 117–123.
- Magoon, L.B., 1994, Tuxedni–Hemlock(!) petroleum system in Cook Inlet, Alaska, U.S.A., in Magoon, L.B., and Dow, W.G., eds., The petroleum system—from source to trap: American Association of Petroleum Geologists Memoir 60, p. 359–370.
- Magoon, L.B., and Anders, D.E., 1990, Oil-source rock correlation using carbon isotope data and biological marker compounds, Cook Inlet-Alaska Peninsula [abs.]: American Association of Petroleum Geologists Bulletin, v. 74, p. 711.
- Magoon, L.B., and Anders, D.E., 1992, Oil-to-source rock correlation using carbon-isotopic data and biological marker compounds, Cook Inlet–Alaska Peninsula, Alaska, in Moldowan, J.M., Albrecht, P., and Philip, R.P., eds., Biological Markers in Sediments and Petroleum: Prentice Hall, Englewood Cliffs, New Jersey, p. 241–274.
- Magoon, L.B., and Claypool, G.E., 1981, Petroleum geology of Cook Inlet basin; an exploration model: American Association of Petroleum Geologists Bulletin, v. 65, p. 1,043–1,061.

- Magoon, L.B., Adkison, W.L., Chmelik, F.B., Dolton, G.L., Fischer, M.A., Hampton, M.A., Sable, E.G., and Smith, R.A., 1976, Hydrocarbon potential, geologic hazards, and infrastructure for exploration and development of the lower Cook Inlet, Alaska: U.S. Geological Survey Open-File Report 76-449, 132 p.
- Magoon, L.B., Egbert, R.M., and Petering, George, 1978, Upper Jurassic and Cretaceous rocks of the Kamishak Hills–Douglas River area, lower Cook Inlet, in Johnson, K.M., ed., The United States Geological Survey in Alaska: Accomplishments during 1977: U.S. Geological Survey Circular 772-B, p. B57–B59.
- Magoon, L.B., Griesbach, F.B., and Egbert, R.M., 1980, Nonmarine Upper Cretaceous rocks, Cook Inlet, Alaska: AAPG Bulletin, v.64, p. 1259–1266.
- Magoon, L.B., Molenaar, C.M., Bruns, T.R., Fisher, M.A., and Valin, Z.C., 1995, Southern Alaska Province (003), in Gautier, D.L., Dolton, G.L., Takahashi, K.I., and Varnes, K.L., eds., 1995 National Assessment of United States Oil and Gas Resources on CD-ROM: U.S. Geological Survey Digital Data Series 30 (CD-ROM), 21 p.
- Mancini, E.A., and Deeter, T.M., 1977, Alaska Peninsula Late Cretaceous fore-arc deposition [abs.]: American Association of Petroleum Geologists Bulletin, v. 61, no. 5, p. 811.
- Mancini, E.A., Deeter, T.M., and Wingate, F.H., 1978, Upper Cretaceous arc–trench gap sedimentation on the Alaska Peninsula: Geology, v. 6, p. 437–439.
- Marincovich, Louie, Jr., 1981, *Tyrannoberingius rex*, a new genus and species of Miocene gastropod from Alaska: Journal of Paleontology, v. 55, p. 176–179.
- 1981, Fossils from the Bear Lake Formation and fossils from the Meshik Formation, in Detterman, R.L., Yount, E., and Case, J., eds., Megafossil localities, checklists, and stratigraphic sections, Chignik and Sutwik Island Quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map 103-N, 2 plates.
- 1983, Molluscan paleontology, paleoecology, and North Pacific correlations of the Miocene Tachilni Formation, Alaska Peninsula, Alaska: Bulletin of American Paleontology, v. 84, no. 317, p. 59–155, plates 12–23.
- 1984, Asiatic mollusks in Miocene faunas of the Alaska Peninsula: U.S. Geological Survey Professional Paper 1375, p. 179–180.
- 1988, Miocene mollusks from the lower part of the Bear Lake Formation on Ukolnoi Island, Alaska Peninsula, Alaska: Natural History Museum of Los Angeles County, Contributions to Sciences, no. 397, 20 p.
- Marincovich, Louie, Jr., and Kase, Tomoki, 1986, An occurrence of *Turritella (Hataiella) sagai* in Alaska; implications for the age of the Bear Lake Formation: Tokyo, Japan, Bulletin of the National Science Museum, Series C, Geology and Paleontology, v. 12, no. 2, p. 61–66.
- Marincovich, Louie, Jr., and Powell, C.L., II, 1989, Preliminary Tertiary molluscan biostratigraphy of the Alaska Peninsula, southwestern Alaska: U.S. Geological Survey Open-File Report 89-674, 2 plates.
- Marincovich, Louie, Jr., Barinov, K.B., and Oleinik, A.E., 2002, The *Astarte* (Bivalvia: Astartidae) that document the earliest opening of Bering Strait: Journal of Paleontology, v. 76, p. 239–245.
- Martin, G.C., 1915, The western part of Kenai Peninsula, in Martin, G.C., Johnson, B.L., and Grant, U.S., Geology and mineral resources of Kenai Peninsula, Alaska: U.S. Geological Survey Bulletin 587, p. 41–112.
- 1916, Triassic rocks of Alaska: Geological Society of America Bulletin, v. 27, p. 685–718
- 1921, Preliminary report on petroleum in Alaska: U.S. Geological Survey Bulletin 719, 83 p., 5 plates, scale 1:125,000.
- 1924, Correlation and paleogeography of the Cretaceous of Alaska: Pan-Pacific Scientific Congress, Australia, 1923, Proceedings, v. 2, p. 1,345–1,359.
- 1926, The Mesozoic stratigraphy of Alaska. U.S. Geological Survey Bulletin 776, 493 p.
- Martin, G.C., and Katz, F.J., 1912, A geologic reconnaissance of the Iliamna region, Alaska: U.S. Geological Survey Bulletin 485, 138 p.
- McClay, K., and Bonora, M., 2001, Analog models of restraining stepovers in strike-slip fault systems: American Association of Petroleum Geologists Bulletin, v. 85, p. 233–260.
- McGee, D.L., 1973, Gulf of Alaska petroleum seeps: Alaska Division of Geological & Geophysical Surveys Open-File Report 32, 9 p.
- McGimsey, R.G., Neal, C.A., and Girina, O., 2003, 1998 volcanic activity in Alaska and Kamchatka: summary of events and response of the Alaska Volcano Observatory: U.S. Geological Survey Open-File Report 03-423, 35 p.
- McLean, H., 1977, Organic geochemistry, lithology, and paleontology of Tertiary and Mesozoic rocks from wells on the Alaska Peninsula: U.S. Geological Survey Open-File Report 77-813, 63 p.
- 1979, Sandstone petrology: Upper Jurassic Naknek Formation of the Alaska Peninsula and coeval rocks on the Bering Shelf: Journal of Sedimentary Petrology, v. 49, no. 4, p. 1263–1268.

- McLean, Hugh, Engelhardt, C.L., and Howell, D.G., 1978, Reconnaissance geologic map of the Cold Bay and False Pass quadrangles, Alaska: U.S. Geological Survey Open-File Report 78-323, 1 plate, scale 1:250,000.
- McRoberts, C.A., 1997, Late Triassic (Norian–Rhaetian) bivalves from the Antimonio Formation, northwestern Sonora, Mexico: *Revista Mexicana de Ciencias Geológicas*, v. 14, no. 2, p. 167–177.
- McRoberts, C.A., and Blodgett, R.B., 2002, Upper Triassic (Norian) mollusks from the Taylor Mountains Quadrangle, southwest Alaska, *in* Wilson, F.H., and Galloway, J.P., eds., *Geologic Studies in Alaska by the U.S. Geological Survey, 2000: U.S. Geological Survey Professional Paper 1662*, p. 55–75.
- Melim, L.A., Swart, P.K., and Maliva, R.G., 1995, Meteoric like fabrics forming in marine waters; Implications for the use of petrography to identify diagenetic environments: *Geology*, v. 23, p. 755–758.
- Melim, L.A., Westphal, H., Swart, P.K., Eberli, G.P., and Munnecke, A., 2002, Questioning carbonate diagenetic paradigms; Evidence from the Neogene of the Bahamas: *Marine Geology*, v. 185, p. 24–54.
- Merritt, R.D., 1986, Coal resources of the Miocene Unga Conglomerate Member, Bear Lake Formation, Unga Island, Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Public-Data File 86-69, 33 p.
- Merritt, R.D., and McGee, D.L., 1986, Depositional environments and resource potential of Cretaceous coal-bearing strata at Chignik and Herendeen Bay, Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Public-Data File 86-72, 44 p.
- Merritt, R.D., McGee, D.L., Clough, J.G., and Belowich, M.A., 1987, Coal Atlas of the Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys unpublished report.
- Mertie, J.B., Jr., 1938, The Nushagak district, Alaska: U.S. Geological Survey Bulletin 903, 96 p., 1 plate, scale 1:250,000.
- Meyer, J.F., Jr., 2007, Total intensity magnetic anomaly for the Bristol Bay region and Alaska Peninsula: Alaska Department of Natural Resources, Division of Oil & Gas, Alaska Peninsula Oil and Gas Field Map Series, plate 3 of 3, scale 1:500,000.
- Meyer, J.F., Jr., Hansen, J.J., Brizzolara, D. W., Pritchard, M.E., Boggess, P.L., Beaty, C.J., Dirks, K.L., Gumpert, J.D., Bolin, K.A., Gibler, K.I., and Phillipson, E.M., 2004, Bristol Bay region and Alaska Peninsula oil and gas programs: Alaska Department of Natural Resources, Division of Oil & Gas, Alaska Peninsula Resource Series, plate 1 of 4.
- 2004, Regional geology of the Bristol Bay region and Alaska Peninsula: Alaska Department of Natural Resources, Division of Oil & Gas, Alaska Peninsula Resource Series, plate 2 of 4.
- 2004, Oil and gas well data for the Alaska Peninsula and Bristol Bay region: Alaska Department of Natural Resources, Division of Oil & Gas, Alaska Peninsula Resource Series, plate 3 of 4.
- 2004, Seismic data availability for the Bristol Bay region and the Alaska Peninsula: Alaska Department of Natural Resources, Division of Oil & Gas, Alaska Peninsula Resource Series, plate 4 of 4.
- 2004, Alaska aeromagnetic compilation, Ugashik quadrangle: Alaska Department of Natural Resources, Division of Oil & Gas, Alaska Peninsula Resource Series, plate 10 of 19, scale 1:250,000.
- 2004, Oil and gas resource map series legend: Alaska Department of Natural Resources, Division of Oil & Gas, Alaska Peninsula Resource Series, plate 19 of 19, Scale 1:250,000.
- 2004, Alaska Peninsula areawide oil and gas lease sale: information and data compilation industry packet: Alaska Division of Oil & Gas Alaska Peninsula Resource Series (digital, updated 2007).
- Mickey, M.B., Haga, H. Boettcher, R.S., and Kling, S.A., 2005, Northwestern Alaska Peninsula–Bristol Bay basin biostratigraphy study: Micropaleo Consultants Job No. 25-104, 287 p., 4 oversize plates, 40 oversize figures, Alaska Division of Oil & Gas online publication <http://www.dog.dnr.state.ak.us/oil/products/publications/akpeninsula/biostrat.htm>
- Micropaleo Consultants, Inc., 2005, North Aleutian shelf public biostratigraphic data set; 11-well study with cross sections correlating the COST well with Alaska Peninsula wells, 286 p., 18 sheets.
- Miller, D.J., Payne, T.G., and Gryc, G., 1959, Geology of possible petroleum provinces in Alaska: U.S. Geological Survey Bulletin 1094, 131 p.
- Miller, J.W., and Jones, D.L., 1981, A field guide to some common megafossils from post-Callovian Mesozoic rocks of the Alaska Peninsula: U.S. Geological Survey Open-File 81-745, 15 p., 1 plate.
- Mobil Oil, 1959, Kuskokwim district, Bristol Bay project: Interpretative seismic map, BBNC files, data set index no. 698, 1 sheet, scale 1:8,000.
- 1963, Exhibit 4, structure contour map, seismic phantom horizon of probably late Miocene: BBNC files, data set index no. 697, 1 sheet, scale 1:8,000.

- Mojsisovics, Edmund, von, 1886, Arktische Triasfaunen: Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg, 7th ser., v. 33, no. 6, 159 p., 20 plates.
- Molenaar, C.M., 1995, Alaska Peninsula, *in* Magoon, L.B., Molenaar, C.M., Bruns, T.R., Fisher, M.A., Valin, Z.C., Southern Alaska Province (003), *in* Gautier, D.L., Dolton, G.L., Takahashi, K.I., and Varnes, K.L., eds., 1995 National assessment of United States oil and gas resources—Results, methodology, and supporting data: U.S. Geological Survey Digital Data Series 30, CD-ROM, p. 2–4.
- 1996, Thermal-maturity patterns and geothermal gradients on the Alaska Peninsula, *in* Johnsson, M.J., and Howell, D.G., eds., Thermal evolution of sedimentary basins in Alaska: U.S. Geological Survey Bulletin 2142, p. 11–19.
- Möller, N.K., and Kvingan, K., 1988, The genesis of nodular limestones in the Ordovician and Silurian of the Oslo Region (Norway): *Sedimentology*, v. 35, p. 405–420.
- Moore, D.M., and Reynolds, R.C., Jr., 1989, X-ray diffraction and the identification and analysis of clay minerals: Oxford University Press, 332 p.
- Moore, G.W., 1967, Preliminary geologic map of Kodiak Island and vicinity, Alaska: U.S. Geological Survey Open-File Report 67-161, 1 plate, scale 1:250,000.
- 1967, Preliminary geologic map of the Kodiak Islands and vicinity: U.S. Geological Survey Open File Report 271, 1 plate, scale 1:250,000.
- Motyka, R.J., Liss, S.A., Nye, C.J., and Moorman, M.A., 1993, Geothermal resources of the Aleutian Arc, Alaska: Alaska Division of Geological & Geophysical Surveys Professional Report 114, 17 p.
- Murphy J.M., and Clough, J.G., 1999, Low-temperature thermal history using fission track dating in three wells in southern Alaska offshore basins; Lower Cook Inlet, Shelikof Strait, and Stevenson Trough: *Marine Georesources and Geotechnology*, v. 17, p. 271–281.
- Naeser, C.W., 1979, Fission track dating and geological annealing of fission tracks, *in* Jager, E. and Hunziker, J.C., eds., *Lectures in Isotope Geology*: New York, Springer-Verlag, p. 154–169.
- 1981, The fading of fission tracks in the geologic environment; Data from deep drill holes: *Nuclear Tracks*, v. 5, p. 248–250.
- Naeser, C.W., and Faul, H., 1969, Fission track annealing in apatite and sphene: *Journal of Geophysical Research*, v. 74, p. 705–710.
- Newton, C.R., 1983, Paleozoogeographic affinities of Norian bivalves from the Wrangellian, Peninsular, and Alexander terranes, northwestern North America, *in* Stevens, C.H., ed., *Pre-Jurassic rocks in western North American suspect terranes*: Society of Economic Paleontologists and Mineralogists Symposium, Sacramento, CA, 1983, p. 37–48.
- 1983, Norian (Late Triassic) mollusks of Cordilleran allochthonous terranes: *Paleoecology and paleozoogeography*: Santa Cruz, University of California, Ph.D. thesis, 173 p.
- 1983, Paleozoogeographic affinities of Norian bivalves from the Wrangellian, Peninsular, and Alexander terranes, northwestern North America, *in* Stevens, C.H., ed., *Pre-Jurassic rocks in western North American suspect terranes*: Sacramento, CA, Society of Economic Paleontologists and Mineralogists Symposium, p. 37–48.
- 1989, Triassic–Jurassic boundary section at Puale Bay, Alaska Peninsula; Comparative diversity patterns of skeletal faunas and trace fossils: *American Association of Petroleum Geologists Bulletin*, v. 74, p. 730.
- 1990, Triassic–Jurassic boundary section at Puale Bay, Alaska Peninsula; Comparative diversity patterns of skeletal faunas and trace fossils [abs.]: *American Association of Petroleum Geologists Bulletin*, v. 74, no. 5, p. 730.
- Nilsen, T., 1985, Sedimentology of tidally deposited Miocene Bear Lake Formation, Alaska Peninsula: *American Association of Petroleum Geologists, Bulletin*, v. 69, no. 4, p. 673.
- Nilsen, T.H., 1984, Miocene back-arc tidal deposits of the Bear Lake Formation, Alaska Peninsula, *in* Reed, K.M., and Bartsch-Winkler, Susan, eds., *The United States Geological Survey in Alaska; Accomplishments during 1982*: U.S. Geological Survey Circular 939, p. 85–88.
- Palache, Charles, 1904, Geology about Chichagof Cove, Stepovak Bay; with notes on Popof and Unga Islands (Alaska): *Harriman Alaska Expedition*, v. 4, p. 69–88.
- Pálfy, József, 1997, Calibration of the Jurassic time scale: Vancouver, University of British Columbia, Ph.D. dissertation, 180 p.
- Pálfy, József, Smith, P.L., Mortensen, J.K., and Friedman, R.M., 1999, Integrated ammonite biochronology and U-Pb geochronometry from a basal Jurassic section in Alaska: *Geological Society of America Bulletin*, v. 111, no. 10, p. 1,537–1,549.

- Pálffy, József., Smith, P.L., Mortensen, J.K., and Friedman, R.M., 1999, Integrated ammonite biochronology and U-Pb geochronometry from a basal Jurassic section in Alaska: *Geological Society of America Bulletin*, v. 111, no. 10, p. 1,537–1,549.
- Parker, J., and Newman, R., 1995, Alaska federal offshore; descriptions of geologic plays; 1995 national resource assessment: U.S. Minerals Management Service, North Aleutian Basin Assessment Province <http://www.mms.gov/alaska/re/asmtdata/naleutia/naleplay.htm>
- Paul, T.A., and Fitzgerald, P.G., 1992, Transmission electron microscopic investigation of fission tracks in fluorapatite: *American Mineralogist*, v. 77, p. 336–344.
- Peters, K.E., 1986, Guidelines for evaluating petroleum source rock using programmed pyrolysis: *American Association of Petroleum Geologists Bulletin*, v. 70, p. 318–329.
- Peters, K.E., and Cassa, M.R., 1994, Applied source rock geochemistry, in Magoon, L.B., and Dow, W.G., eds., *The petroleum system—from source to trap*: American Association of Petroleum Geologists Memoir 60, p. 93–117.
- Phillips Petroleum, 1974, Seismic program maps: BBNC files, data set index no. 422, 1 sheet, scale 1:250,000.
- 1974, Aeromagnetic interpretation, Chignik area: BBNC files, data set index no. 661, several sheets, multiple scales.
- 1974, BBNC program map, seismic; Nugashak, Dillingham, Naknek, Ugashik: BBNC files, data set index no. 668, several sheets, various scales.
- 1974, BBNC program map, seismic; Chignik, Sutwik Island, Stepovak Bay: BBNC files, data set index no. 669, several sheets, various scales.
- 1974, Chignik Bay and Mitrofinia shot points, BBNC files, data set index no. 712, 1 sheet, scale 1:8,000.
- 1974, Chignik Bay seismic interpretation: BBNC files, data set Index No. 711, 1 sheet, scale 1:8,000.
- 1974, Total magnetic intensity, Nushagak Basin, Alaska: BBNC files, data set index no. 744, 1 sheet, scale 1:16,000.
- 1975, Petrology report, Alaska Peninsula–Nushagak area for Bristol Bay Native Corporation: BBNC files, data set index no. 423, 73 p.
- 1975, Bouguer gravity maps, Mitrofinia area, Alaska: BBNC files, data set Index No. 560, 2 sheets, scale 1:63,360.
- Pinart, A.L., 1873, Voyage à la côte nord-ouest de l'Amérique d'Ounalashka à Kadiak: Paris, *Bulletin de la Société de Géographie*, série VI, v. 6, p. 561–580.
- Powell, T.G., and Boreham, C.J., 1994, Terrestrially sourced oils; Where do they exist and what are our limits of knowledge? – A geochemical perspective, in Scott, A.C., and Fleet, A.J., eds., *Coal and coal-bearing strata as oil-prone source rocks?: Geological Society Special Publication No. 77*, p. 11–29.
- Price, P.B., and Walker, R.M., 1963, Fossil tracks of charged particles in gneiss and the age of minerals: *Journal of Geophysical Research*, v. 68, p. 4,847–4,862.
- Reed, B.L., and Lanphere, M.A., 1973, Alaska–Aleutian Range batholith: Geochronology, chemistry, and relation to circum-Pacific plutonism: *Geological Society of America Bulletin* 84: p. 2,583–2,610.
- Reed, B.L., Miesch, A.T., and Lanphere, M.A., 1983, Plutonic rocks of Jurassic age in the Alaska–Aleutian Range batholith: chemical variations and polarity: *Geological Society of American Bulletin*, v. 94, p. 1232–1240.
- Reifenstuhel, R.R., and McCarthy, Paul, 2004, Petroleum reservoir characterization, source rock potential, fossil fuel resources, and basin analyses, Bristol Bay Basin, Alaska: AETDL Quarterly Report, October, 2004, 1 p.
- Reifenstuhel, R.R., Bailey, R.D., and Finzel, E.S., 2005, Bristol Bay and Alaska Peninsula 2004: Fieldwork and sample analyses compilation report: Alaska Division of Geological & Geophysical Surveys Preliminary Interpretive Report 2005-1, 20 p.
- Reifenstuhel, R.R., Decker, P.L., Gillis, R.J., Helmold, K.P., and Strauch, A., 2007, Hydrocarbon systems, basin analyses, and 2006 field and subsurface data; Bristol Bay, Alaska Peninsula, Frontier Basin [abs.]: *American Association of Petroleum Geologists Bulletin*, v. 91, Program Abstracts (digital).
- Reifenstuhel, R.R., Shafer, D.C., Ryherd, T.J., Brizzolara, D.W., and Blodgett, R.B., 2004, Summary of May 25–June 4, 2004, field notes and samples, Puale Bay and Wide Bay areas, Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Raw Data File 2004-3, 16 p.
- Riehle, J.R., and Detterman, R.L., 1993, Quaternary geologic map of the Mount Katmai quadrangle and adjacent parts of the Naknek and Afognak quadrangles, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map 2032, 1 plate, scale 1:250,000.

- Riehle, J.R., Detterman, R.L., Yount, M.E., and Miller, J.W., 1993, Geologic map of the Mt. Katmai quadrangle and portions of the Naknek and Afognak quadrangles, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-2204, 1 plate, scale 1:250,000. [see USGS Open-File Report 87-593]
- Riehle, J.R., Wilson, F.H., Shew, Nora, and White, W.H., 1999, Geology of Unga Island and the northwestern part of Popof Island, in Riehle, J.R., ed., A geological and geophysical study of the Gold–Silver vein systems of Unga Island, southwestern Alaska: U.S. Geological Survey Open-File Report 99-136, CD-ROM.
- Saltus, R.W., Connard, G.G., and Hill, P.L., 1999, Alaska aeromagnetic compilation—digital grids and survey data: U.S. Geological Survey Open-File Report 99-502, CD-ROM.
- Saltus, R.W., Haeussler, P.J., Bracken, R.E., Doucette, J.P., and Jachens, R.C., 2001, Anchorage Urban Region Aeromagnetism (AURA) Project – preliminary geophysical results: U.S. Geological Survey Open-File Report 01-0085, 21 p.
- Saltus, R.W., Hudson, T.L., Wilson, F.H., 2003, Geophysical identification and geological implications of the southern Alaska magnetic trough, U.S. Geological Survey Open-File Report 03-200, 1 plate (poster).
- Sandy, M.R., Blodgett, R.B., and Frýda, J., 2001, Paleobiogeographic signatures for Upper Triassic brachiopods and gastropods from Kuiu Island and adjacent Keku Strait, SE Alaska [abs.]: Geological Society of America Abstracts with Programs, v. 33, no. 3, p. A-53.
- Scholl, D.W., Vallier, T.L., and Stevenson, A.J., 1986, Terrane accretion, production, and continental growth; A perspective based on the origin and tectonic fate of the Aleutian–Bering Sea region: *Geology*, v. 14, p. 43–47.
- 1987, Geologic evolution and petroleum geology of the Aleutian Ridge, in Scholl, D.W., Grantz, Arthur, and Vedder, J.G., eds., *Geology and resource potential of the continental margin of western North America and adjacent ocean basins, Beaufort Sea to Baja California*: Houston, TX, Circum-Pacific Council For Energy and Mineral Resources, Houston, TX, p. 73–102.
- Schreurs, G., and Colletta, B., 2002, Analog modelling of continental transpression, in Schellart, P., and Passchier, C., eds., *Analogue modelling of large-scale tectonic processes: Journal of the Virtual Explorer*, v. 7., p. 67–78.
- Sherwood, K.W., and Craig, J.D., 2001, Prospects for development of Alaska natural gas—A review: U.S. Minerals Management Service, Resource Evaluation Office, 135 p.
- Sherwood, K.W., Comer, D., and Larson, J., 2003, Review of geologic model for North Aleutian Basin: U.S. Minerals Management Service, “Handout,” 25 p.
- Sherwood, K.W., Craig, J.D., and Cooke, L.W., 1996, Endowments of undiscovered conventionally recoverable and economically recoverable oil and gas in the Alaska federal offshore: U.S. Minerals Management Service, Alaska Outer Continental Shelf Region, OCS Report MMS 96-0033, 6 p.
- Sherwood, K.W., Larson, J., Comer, C.D., Craig, J.D., and Reitmeier, C., 2006, North Aleutian basin OCS planning area assessment of undiscovered technically-recoverable oil and gas: U.S. Minerals Management Service Alaska OCS Region report, 138 p., 4 plates, www.mms.gov/alaska/re/reports/rereport.htm
- Silberling, N.J., 1985, Biogeographic significance of the Upper Triassic bivalve *Monotis* in circum-Pacific accreted terranes, in Howell, D.G., ed., *Tectonostratigraphic terranes of the Circum-Pacific region*: Circum-Pacific Council for Energy and Mineral Resources Earth Sciences Series No. 1, p. 63–70.
- Silberling, N.J., Grant-Mackie, J.A., and Nichols, K.M., 1997, The Late Triassic bivalve *Monotis* in accreted terranes of Alaska: U.S. Geological Survey Bulletin 2151, 21 p.
- Smith, J.P., 1927, Upper Triassic marine invertebrate faunas of North America: U.S. Geological Survey Professional Paper 141, 262 p.
- Smith, W.R., 1924, The Upper Cretaceous invertebrate fauna of Alaska: Baltimore, MD, The Johns Hopkins University, unpublished Ph.D. dissertation, 178 p.
- 1925, The Cold Bay–Katmai district, in *Mineral resources of Alaska; report on progress of investigations in 1923*: U.S. Geological Survey Bulletin 773, p. 183–207.
- 1926, Geology and oil development of the Cold Bay district, in *Mineral Resources of Alaska; Report on progress of investigations in 1924*: U.S. Geological Survey Bulletin 783, p. 63–88.
- Smith, W.R., and Baker, A.A., 1924, The Cold Bay–Chignik district, in *Mineral resources of Alaska; Report on progress of investigations in 1922*: U.S. Geological Survey Bulletin 755, p. 151–218.
- Sneider, R.M., 1997, Petrophysical properties of seals, in Surdam, R.C., ed., *Seals, traps and the petroleum system*: Tulsa, OK, American Association of Petroleum Geologists Memoir 67, 317 p.
- Spicer, R.A., 1987, Late Cretaceous floras and terrestrial environment of northern Alaska, in Tailleux, Irv, and Weimer, Paul, eds., *Arctic North Slope Geology, Volume 1: Pacific Section*, SEPM, and The Alaska Geological Society, p. 497–512. [Mentions floras from Chignik area, Nelson Island, Nunivak Island, Yukon River, North Slope; see fig. 2 for location]

- Spurr, J.E., 1900, A reconnaissance in southwestern Alaska in 1898: U.S. Geological Survey 20th Annual Report, Part 7, p. 31–264.
- Sralla, Bryan, 2007, Structural geometry, kinematics, and hydrocarbon exploration potential of Sapsuk Lake anticline, Herendeen Bay region, Alaska Peninsula [abs]: *Alaska Geology*, v. 38, no. 1, p. 1–2.
- Sralla, Bryan, and Blodgett, R.B., 2007, Reservoir potential of Late Triassic Kamishak Formation; Puale Bay, Alaska Peninsula [abs.]: American Association of Petroleum Geologists Annual Convention & Exhibition, Long Beach, CA, 2007, Abstracts, v. 91 (digital), p. 132.
- Stanley, G.D., Jr., 1979, Paleogeology, structure, and distribution of Triassic coral buildups in western North America: *University of Kansas Paleontological Contributions*, v. 65, p. 1–58.
- Stanton, T.W., and Martin, G.C., 1905, Mesozoic section on Cook Inlet and Alaska Peninsula: *Geological Society of America Bulletin*, v. 16, p. 391–410.
- Strauch, A.L., Gillis, R.J., Reifenhohl, R.R., and Decker, P.L., 2006, 2006 Bristol Bay, Alaska Peninsula field summary and outcrop sample results from porosity & permeability and mercury injection capillary pressure analyses: Alaska Division of Geological & Geophysical Surveys Raw Data File 2006-1, 65 p.
- Streckeisen, A., 1976, To each plutonic rock its proper name: *Earth Science Review*, v. 12, p. 1–33.
- Swanson, B.F., 1981, A simple correlation between permeabilities and mercury capillary pressures: *Journal of Petroleum Technology*, p. 2,498–2,504.
- Symonds, R.B., Ritchie, B.E., McGimsey, R.G., Ort, M.H., Poreda, R.J., Evans, W.C., and Janik, C.J., 1997, Investigations of gas seeps and springs in the vicinity of The Gas Rocks, south shore Becharof Lake, Alaska: U.S. Geological Survey Open-File Report 97-0127, 27 p.
- Tagami, T., and O'Sullivan, P.B., 2005, Fundamentals of fission-track thermochronology: *Reviews in Mineralogy and Geochemistry*, v. 58, p. 19–47.
- Teller, Friedrich, 1886, Die Pelecypod-Fauna von Werchojansk in Ostsiberien, in Mojsisovics, Edmund, von, ed., *Arktische Triasfaunen: Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg*, 7th ser., v. 33, no. 6, p. 103–137.
- Thomas, C.P., Doughty, T.C., Faulder, D.D., and Hite, D.M., 2004, South-central Alaska natural gas study: U.S. Department of Energy, National Energy Technology Laboratory–Arctic Energy Office, www.netl.doe.gov, 207 p.
- Tissot, B.P., and Welte, D.H., 1984, *Petroleum formation and occurrence*, 2nd ed.: Berlin, Springer-Verlag, 699 p.
- Triplehorn, D.M., 1976, Large kaolinite crystals in the Chignik Formation (Upper Cretaceous), Herendeen Bay, in *Short Notes on Alaskan Geology 1976*: Alaska Division of Geological & Geophysical Surveys Geologic Report 51F, p. 23–25.
- Troutman, S.M., and Stanley, R.G., 2003, Maps showing sedimentary basins, surface thermal maturity, and indications of petroleum in the Central Alaska Province, U.S. Geological Survey Miscellaneous Field Studies Map 2428, pamphlet, 2 plates, scale 1:2,500,000.
- Tucker, M.E., and Wright, V.P., 1990, *Carbonate Sedimentology*: Oxford, England, Blackwell Publishing, 482 p.
- Turner, R.F., ed., 1988, Geological and operational summary, North Aleutian Shelf COST No. 1 well, Bering Sea, Alaska: U.S. Minerals Management Service, OCS Report MMS 88-0089, 256 p., 2 plates.
- U.S. Geological Survey, 1963, Aeromagnetic survey of parts of the Ugashik and Karluk quadrangles: BBNC files, data set index no. 526, 1 plate, scale 1:125,000.
- 1963, Aeromagnetic map of parts of the Dillingham quadrangle: BBNC files, data set index no. 727, 1 plate, scale 1:125,000.
- 1971, Aeromagnetic survey, Goodnews area, Bethel A-1 through A-5 quadrangles: BBNC files, data set index no. 528, 1 plate, scale 1:63,360.
- U.S. Geological Survey, 1971, Aeromagnetic survey, Goodnews area, Hagemester C-3 through C-7 and D-1 through D-6 quadrangles: BBNC files, data set index no. 529, 11 sheets, plate 1:63,360.
- 1971, Aeromagnetic survey, Goodnews area, Goodnews Bay A-1 through A-8, B-1 through B-8, and C-1 through C-8 quadrangles: BBNC files, data set index no. 530, 24 plates, scale 1:63,360.
- 1971, Aeromagnetic survey, Goodnews area, Nushagak Bay B-4, C-4, and D-4 through D-8 quadrangles: BBNC files, data set index no. 531, 5 plates, scale 1:63,360.
- 1975, Aeromagnetic map of part of the Naknek quadrangle, Alaska: BBNC files, data set index no. 505, 1 plate, scale 1:250,000.
- 1978, Aeromagnetic map of Chignik and Sutwik Island quadrangles, Alaska: U.S. Geological Survey Open-File Report 78-262, 1 plate, scale 1:250,000.

- U.S. Geological Survey, 1978, Aeromagnetic maps of the Chignik and Sutwik Island quadrangles: BBNC files, data set index no. 556, 11 plates, scale 1:63,360.
- 1978, Aeromagnetic map of the Chignik and Sutwik Island quadrangles: BBNC files, data set index no. 557, 1 plate, scale 1:250,000.
- 1995, 1995 national assessment of the United States oil and gas resources: U.S. Geological Survey National Oil and Gas Resource Assessment Team, U.S. Geological Survey Circular 1118, 20 p.
- U.S. Minerals Management Service, 1985, Final Environmental Impact Statement, North Aleutian Basin Sale No. 92, OCS EIS/EA MMS 85-0052, v. 1, 583 p., http://www.mms.gov/alaska/ref/EIS%20EA/N_Aleutian_Basin_FEIS_92/mms85_0052Volume1.pdf
- 2000, Undiscovered Oil and Gas Resources, Alaska Federal Offshore, December 2000 Update, U.S. Minerals Management Service, 5 p., <http://www.mms.gov/alaska/re/uogr/uogr.pdf>.
- 2006, North Aleutian Basin planning area (Alaska); Province summary, 2006 oil and gas assessment, (<http://www.mms.gov/alaska/re/reports/2006Asmt/NAGA/naga.HTM>), 24 p.
- Union Oil, 1974, Plate 2B, Seismic project map, Nushagak Project: BBNC files, data set index no. 670, 1 sheet, scale 1:63,360.
- Vacquier, V., 1963, General principles, in Vacquier, V., Steenland, N. C., Henderson, R.G., and Zietz, I., Interpretation of aeromagnetic maps: Geological Society of America Memoir 47, 151 p., 14 plates.
- Van Kooten, G.K., Short, J.W., and Kolak, J.J., 2002, Low-maturity Kulthieth Formation coal; A possible source of the polycyclic aromatic hydrocarbons in benthic sediments of the northern Gulf of Alaska: Environmental Forensics, v. 3, p. 227–241.
- Veach, N.J., 1973, Aeromagnetic map, northeastern part of Hagemeister Island Quadrangle: Alaska Division of Geological & Geophysical Surveys Alaska Open-File Report 16, 5 p., 1 sheet, scale 1:250,000.
- von Huene, Roland, Moore, J.C., and Moore, G.W., 1979, Cross section of Alaska Peninsula–Kodiak Island–Aleutian Trench: Geological Society of America Map and Chart Series, 1979, v. MC-28A, 2 sheets, scale 1:250,000.
- Waldron, H.H., 1961, Geologic reconnaissance of Frosty Peak Volcano and vicinity, Alaska: U.S. Geological Survey Bulletin 1028-T, p. 677–708, 1 plate, scale 1:250,000.
- Walker, K.T., McGeary, S.E., and Klemperer, S.L., 2003, Tectonic evolution of the Bristol Bay basin, southeast Bering Sea: Constraints from seismic reflection and potential field data: Tectonics, v. 22, no. 5, p. 19.
- Wang, Jason, 1987, Sedimentology of Norian (Late Triassic) cherts and carbonates from the Peninsular Terrane, Puale Bay, Alaska Peninsula: Syracuse, NY, Syracuse University, unpublished M.S. thesis, 124 p.
- Wang, Jason, Newton, C.R., and Dunne, L., 1988, Late Triassic transition from biogenic to arc sedimentation on the Peninsular terrane, Puale Bay, Alaska Peninsula: Geological Society of America Bulletin, v. 100, no. 9, p. 1,466–1,478.
- Waples, D., 1981, Organic geochemistry for exploration geologists: Minneapolis, Minnesota, Burgess Publishing Company, 151 p.
- Weems, R.E., and Blodgett, R.B., 1996, The Pliosaurid *Megalneusaurus*; A newly recognized occurrence in the Upper Jurassic Naknek Formation of the Alaska Peninsula, in Moore, T.E., and Dumoulin, J.A., eds., Geologic studies in Alaska by the U.S. Geological Survey, 1994: U.S. Geological Survey Bulletin 2152, p. 169–175.
- Westermann, G.E.G., 1964, The ammonite fauna of the Kialagvik Formation at Wide Bay, Alaska Peninsula, Part 1, Lower Bajocian (Aalenian): Bulletins of American Paleontology, v. 47, no. 216, p. 327–503.
- 1969, The ammonite fauna of the Kialagvik Formation at Wide Bay, Alaska Peninsula, Part 2, Sonninia Sowerbyi Zone (Bajocian): Bulletins of American Paleontology, v. 57, no. 255, p. 4–226.
- 1978, *Alaskinia* nom. nov. for *Alaskoceras* Westermann, 1969 non Miller and Kummel, 1945; Jurassic Ammonitina: Journal of Paleontology, v. 52, p. 604.
- Wetzel, A., and Werner, F., 1980, Morphology and ecological significance of Zoophycos in deep-sea sediments off northwest Africa: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 32, p. 185–212.
- Whalen, M., this volume.
- Whalen, M.T., Eberli, G.P., van Buchem, F.S.P., Mountjoy, E.W., and Homewood, P.W., 2000, Bypass margins, basin-restricted wedges and platform-to-basin correlation, upper Devonian, Canadian Rocky Mountains; Implications for sequence stratigraphy of carbonate platform systems: Journal of Sedimentary Research, v. 70, p. 913–936.
- White, C.A., 1889, Mesozoic molluscs from the southern coast of the Alaskan Peninsula: U.S. Geological Survey Bulletin 51, p. 64–70.

- Whitney, John, Hoose, P.J., Smith, L.M., and Lybeck, Lynn, 1980, Geologic cross sections of the Outer Continental Shelf of Shelikof Strait, Alaska: U.S. Geological Survey Open-File Report 80-2036, 1 plate, scale 1:250,000.
- Wilson, F.H., 1977, Some plutonic rocks of southwestern Alaska, a data compilation: U.S. Geological Survey Open-File Report 77-501, 7 p., 4 plates, scale 1:1,000,000.
- 1985, The Meshik arc—An Eocene to earliest Miocene magmatic arc on the Alaska Peninsula: Alaska Division of Geological & Geophysical Surveys Professional Report 88, 14 p.
- 1989, Geologic setting, petrology, and age of Pliocene to Holocene volcanoes of the Stepovak Bay area, western Alaska Peninsula, *in* Dover, J.H., and Galloway, J.P., eds., *Geologic studies in Alaska by the U.S. Geological Survey*, 1988: U.S. Geological Survey Bulletin 1903, p. 84–95.
- Wilson, F.H., and Coonrad, W.L., 2005, The Togiak–Tikchik Complex of southwestern Alaska, a replacement for the Gemuk Group: Stratigraphic nomenclature that has outlived its time: U.S. Geological Survey Scientific Investigations Report 2005-5019, 12 p.
- Wilson, F.H., and Shew, N., 1992, Map and tables showing geochronology and whole-rock geochemistry of selected samples, Ugashik and part of Karluk quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1539-E, 12 p., 1 plate, scale 1:250,000.
- Wilson, F.H., and Weber, F.R., 2001, Quaternary geology, Cold Bay and False Pass quadrangles, Alaska Peninsula, *in* Gough, L.P., and Wilson, F.H., eds., *Geologic studies in Alaska by the U.S. Geological Survey*, 1999: U.S. Geological Survey Professional Paper 1633, p. 51–71.
- Wilson, F.H., Case, J.E., and Detterman, R.L., 1985, Preliminary description of a Miocene zone of structural complexity in the Port Moller and Stepovak Bay quadrangles, Alaska, *in* Bartsch-Winkler, Susan, and Reed, K.M., eds., *The United States Geological Survey in Alaska, Accomplishments during 1983: U.S. Geological Survey Circular*, 945, p. 55–56.
- Wilson, F.H., Detterman, R.L., and Case, J.E., 1985, The Alaska Peninsula terrane; A definition: U.S. Geological Survey Open-File Report 85-450, 17 p.
- Wilson, F.H., Detterman, R.L., and DuBois, Gregory, 1999, Digital data for the geologic framework of the Alaska Peninsula, Southwest Alaska, and the Alaska Peninsula Terrane: U.S. Geological Survey Open-File Report 99-317, 41 p., 1 plate, scale 1:500,000, <http://wrgis.wr.usgs.gov/open-file/of99-317>.
- Wilson, F.H., Detterman, R.L., Miller, J.W., and Case, J.E., 1995, Geologic map of the Port Moller, Stepovak Bay, and Simeonof Island quadrangles, Alaska Peninsula, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-2272, 2 plates, scale 1:250,000.
- Wilson, F.H., Gajewski, S.Z., and Angeloni, L.M., 1986, Geological literature of the Alaska Peninsula to 1985: U.S. Geological Survey Open-File Report 86-176, 113 p.
- Wilson, F.H., Gaum, W.C., and Herzon, P.L., 1981, Maps and tables showing geochronology and whole-rock geochemistry, Chignik and Sutwik Island quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1053-M, 3 plates, scale 1:250,000.
- Wilson, F.H., Hudson, T.L., Grybeck, Donald, Stoesser, D.B., Preller, C.C., Bickerstaff, Damon, Labay, Keith, and Miller, M.L., 2003, Preliminary geologic map of the northeast Dillingham quadrangle (D-1, D-2, C-1, and C-2), Alaska: U.S. Geological Survey Open-File Report 03-105, 13 p., 1 plate, scale 1:100,000.
- Wilson, F.H., Shew, N., DuBois, G.D., and Bie, S.W., 1994, Sample locality map and analytical data for potassium-argon ages in the Port Moller, Stepovak Bay, and Simeonof Island quadrangles, Alaska Peninsula: U.S. Geological Survey Miscellaneous Field Studies Map MF-2155-E, 18 p., 1 plate, scale 1:250,000.
- Wilson, F.H., Weber, F.R., Dochat, T.M., Miller, T.P., and Detterman, R.L., 1997, Revised geologic map of the Cold Bay and False Pass quadrangles, Alaska Peninsula: U.S. Geological Survey Open-File Report 97-866, 34 p., 1 plate, scale 1:250,000.
- Wisehart, R.M., 1971, Paleoenvironmental analysis of the Bear Lake Formation (upper and middle Miocene), Alaska Peninsula, Alaska: Los Angeles, CA, University of California, M.S. thesis, 112 p.
- Worrall, D.M., 1991, Tectonic history of the Bering Sea and the evolution of Tertiary strike-slip basins of the Bering Shelf: Geological Society of America Special Paper 257, 120 p., 1 oversize sheet, 4 plates, scale 1:2,500,000.
- Yount, M.E., Wilson, F.H., and Miller, J.W., 1985, Newly discovered Holocene volcanic vents, Port Moller and Stepovak Bay quadrangles, Alaska Peninsula, *in* Bartsch-Winkler, Susan, and Reed, K.M., eds., *The United States Geological Survey in Alaska, Accomplishments during 1983: U.S. Geological Survey Circular*, 945, p. 60–62.
- Zhang, Ning, Blodgett, R.B., and Stanley, G.D., 2003, Paleontological database applications for Alaskan terrane studies [abs.]: Geological Society of America Abstracts with Programs, v. 35, no. 4, p. 24.