


INDEX MAP




PANORAMIC VIEW OF THE MENDENHALL GLACIER TERMINUS, APRIL 1966




AREAL VIEW OF MENDENHALL VALLEY, MENDENHALL GLACIER
IN THE BACKGROUND, SEPTEMBER 1966



QUALITY OF SURFACE AND GROUND WATERS MAP



OCCURRENCE OF WATER AND ICE MAP




SURFICIAL GEOLOGIC MAP AND SECTIONS

EXPLANATION
GEOLOGIC FORMATIONS AND THEIR
WATER-BEARING PROPERTIES


- Qal Alluvial deposits
Sand and gravel, well sorted and sorted by stream action, better than average permeability, will yield moderate supplies of water
- Qc Clay
Deposited marginally to Lake Lake as much as 60 feet thick. Not water bearing
- Qm Colluvial deposits
Landslide and talus deposits, chiefly very coarse boulders and rock fragments, poorly sorted. Not a source of water
- Qf Flood-plain and outwash deposits
Silt and sand, locally gravel and sand, yield moderate quantities of water having iron content more than 2 ppm and high silt content. Relatively impermeable
- Qab Abandoned outwash channel deposits
Sand and gravel, locally boulders, predominantly younger than A. D. 1750, richest aquifers in valley. May produce large quantities of water having relatively low iron content
- Qfa Alluvial fan and delta deposits
Sand and gravel, locally silt and clay, yield moderate quantities of water
- Qr Glacier recessional deposits
Silty sand and gravel yield moderate quantities of water. Include some outwash or alluvial deposits and contain permeable sands in abandoned stream channels
- Qem End-moraine deposits
Cobbles, gravel, and silty sand, predominantly coarse and poorly sorted, yield moderate quantities of water
- Qla Lateral glacier margin deposits
Silty gravel and sand, locally coarse boulders, probably yield moderate quantities of water. No known wells
- Qm Alluvial, estuarine, and marine deposits
Gravelly sand, silt, and clay, which generally are not water bearing, to younger (Recent) deposits of clay, silt, and sand, which yield moderate quantities of water
- UNCONFORMITY
- Mu Metamorphic rocks of igneous and sedimentary origin
Gneiss, granite, argillite, slate, quartz, diorite, and gneiss. Average yield, 1 gpm or less
- Contact
Dashed where approximately located; short dashed where inferred
- Gravel pit
- Well
- Test hole
- Wood sample
- TP-1
- Test pit

QUATERNARY


MESOCZOIC




MARINE PHASE (10,000-4500 B. C.)



ESTUARINE AND NONMARINE PHASE (4500 B. C.-A. D. 1750)



NONMARINE PHASE (A. D. 1750 TO PRESENT)



PALEOGEOGRAPHIC MAPS

WATER RESOURCES AND SURFICIAL GEOLOGY OF THE MENDENHALL VALLEY, ALASKA

By
William W. Barnwell and Charles W. Boning
1968