


MAP SHOWING LOCATION OF GLACIERS, GLACIER-DAMMED LAKES
GLACIER-SHEATHED VOLCANOES AND AREAS SUBJECT TO
OUTBURST FLOODING IN SOUTHEASTERN ALASKA

OUTBURST FLOODS AND NON-OUTBURST FLOODS
ON FOUR RIVERS



1964 OUTBURST FLOOD ON SNOW AND
KENAI RIVERS




Table 1.—Information on selected glacier dammed lakes and the areas flooded by outbursts. (Lake area includes the part of the ice dam which shows crevasses evidence of floating on the lake. The areas flooded may include the entire flood plain of the affected river listed. Unless indicated otherwise, the lakes are in Alaska and source maps are those of the U.S. Geological Survey.)																			
Basin	Lake No. on map	Name of lake or depression	Maximum area (km ²)		Damming glacier	Area flooded	Comments, hazards, and recommendations	Topographic maps and data sources in addition to aerial photographs											
Salmon River	1	Summit Lake (British Columbia)	4.2	1.6	Salmon	Salmon River	Outburst floods began abruptly in 1961. Future outburst floods may occur at 1- to 2-year intervals. Flood damage from this lake was reported by Alaska Department of Highways (1970). ***The drainage of the lake this year caused extensive damage to the Hyder road from mile 6 to 9 with various minor roadway washouts from 1 to 6 miles. ***Due to the magnitude of the flood and extensive damage caused by the flooding, Governor Miller declared Hyder a disaster area. Extreme flood hazards in Salmon River valley. Monitored by Canada.	Map, Iskut River 104B, Canada; Map, Salmon Glacier, British Columbia, Special Map by Canadian Army Survey Establishment, 1959 Alaska Department of Highways (1970, p.10), Doell (1963, p.435), Field (1958a, p.2a.1.17), Fisher (1969), Gilbert (1969), Mathews 1965, p.46, and 1971) U.S. Geological Survey gaging station 15-80											
Siskine River	2	Flood Lake (British Columbia)	2.9	1.1	Flood	Flood River	Little data on flood history. Lake drains under Glacier, probably annually. Lake is shown much smaller on 1909 map. Extreme flood hazards exist in Flood River and moderate flood hazards in Siskine River lowlands. Collecting data on future floods is recommended.	Map, Telegraph Creek 104G, Canada; Map, International Boundary Commission, Sheet 5, 1909 Dawson (1889, p.538), Field (1958a, p.2a.1.45), Kerr (1928, p.16A), Muir (1915, p.101), Patterson (1908, p.558) Seidmore (1899, p.9), Stone (1963a)											
Thomas Bay	3	Unamed	.8	.3	Patterson	Patterson River	Little data on flood history. Drains annually most years during spring or summer. Extreme flood hazard in Patterson River lowlands. Collecting data on future floods is recommended.	Map, Petersburg (D-2), Alaska Stone (1963a)											
	4	Witches Cauldron	2.0	.8	Baird	Baird Glacier outwash closed basin in the future. In such situations large lakes generally form subglacially; there is little evidence of a subglacial lake at present. Extreme flood hazard on Baird Glacier flood plain due to many dumping lakes in basin.	Map, Seward (A-2), Alaska Field (1958a, p.2a.1.50)												
Taku Inlet	5	Tulequah Lake (British Columbia)	4.0	1.5	Tulequah	Tulequah River, Taku River	This lake and a smaller lake up glacier dump most years. In 1920, Tulequah Lake covered 6 square kilometers (2.3 square miles). A midwinter outburst occurred in January 1926. Extreme flood hazard in Tulequah River and moderate flood hazard in Taku River lowlands. Monitoring is recommended.	Map, Tulequah 104K, Canada; International Boundary Commission (1952, p.98-99) Field (1958a, p.2a.1.70), Kerr (1934, 1936), Mandy (1936, p.820), Marcus (1960), Miller (1952, p.74-80, 1963, p.116-200, 1970, p.20), Stone (1955, 1963a)											
	6	Dead Branch	3.4	1.3	Norris	Grizzly Bar	Central crevassing indicates the presence of a subglacial lake which occasionally causes the ice to flow. Continued recession could form a large lake here. Glory Lake, near terminus of Norris Glacier, dumped vigorously until melting of the ice dam around 1930. These floods prevented the growth of vegetation on Grizzly Bar. Vegetation now becoming established indicates that the Dead Branch subglacial lake has not yet caused major outburst flooding. Moderate flood hazard on Grizzly Bar.	Map, Juneau (B-1), Alaska											
Katikash River	7	Unamed	1.0	.4	Metale	Katikash River	May dump annually, generally drains in late August. Collecting data on future floods is recommended.	Map, Skagway (A-1), Alaska											
Glacier Bay	8	Unamed	5.2	2.0	Carroll	Carroll Glacier outwash plain	In 1968 and in 1969 a large lake was formed between Carroll and Pataua Glaciers by the Carroll Glacier surge of 1968. The lake drained under the Carroll Glacier in September each year. In the near future this lake will probably shift to dumping under Plateau Glacier due to the latter's retreat; virtual disappearance of Plateau Glacier's ice dam by about 1990 will drain the lake basin. Extreme flood hazard on Carroll Glacier outwash plain. Monitoring as long as a large lake forms is recommended.	Map, Mt. Fairweather (D-2), Alaska Map, Skagway (A-5), Alaska											
Lituya Bay	9	Desolation Valley	4.1	1.6	Lituya	Lituya Glacier outwash plain	Former subglacial lake recently exposed by glacier's recession. Extreme flood hazard on outwash plain at terminus of Lituya Glacier.	Map, Mt. Fairweather (C-5), Alaska											
Aleak River	10	Recent Lake Aleak (Yukon Territory)	30.7	12.7	Lowell	Aleak River	Extremely hazardous Recent Lake Aleak will reform only if glacier surges strongly. Monitoring glacier surges is recommended.	Map, Dezadeash 115A, Canada Kendle (1953, p.21-22, map 1019A), McConnell (1904, p.3A-4A), Tarr and Martin (1914, p.194)											
	11	Unamed (British Columbia)	16.7	6.7	Tweedamir	Aleak River	Hazardous lake may form if glacier surges moderately. A lake was apparently formed by a surge which occurred around 1945. Monitoring glacier surges is recommended.	Map, Tashishini River 114P, Canada Map, Mt. St. Elias, 1:250,000, Alaska											
	12	Unamed (British Columbia)	6.2	2.4	Konamox	Melt Creek, Aleak River	Large lake now forming by retreat of Melt Creek Glacier. As lake increases in size major floods may result. Moderate flood hazard on Melt Creek and Aleak River flood plains.	Map, Tashishini River 114P, Canada											
Yakutat Bay	13	Russell Flood at present time	260	100	Hubbard	Would drain directly to Dunsmuir channel by Bering River under or along margin of glacier	Hubbard Glacier has advanced intermittently since mapped in 1895. The glacier will close off the entrance to Russell Flood in about 20 years if the present average rate of advance continues. No present flood hazard but extreme danger to boats near glacier margin and in tidal currents at mouth of flood.	Map, Mt. St. Elias, 1:250,000, Alaska Map, International Boundary Commission, Sheet 13, 1895 Muir (1915, p.108-109, pl. 36, map 3)											
Bering River	14	Berg Lake	28	11	Bering	Bering River, Bering Lake	The lake is presently spilling over a bedrock saddle. Retreat of Bering Glacier has greatly increased the size of this lake and recently created an extreme flood hazard on Bering River lowlands. Monitoring is recommended.	Map, Bering Glacier, 1:250,000, Alaska Ellsworth and Davenport (1915, p.36, pl. 2), Field (1958b, p.2a.3.4-8), Martin (1905, p.17, 1908, p.46-48, pl. 2), Post (1967, table 5), Stone (1963a)											
Copper River	15	Iceberg Lake	1.8	.7	Tazlina	Tazlina Glacier outwash plain, Tazlina Lake, Tazlina River, Copper River	Two lakes drain subglacially. In 1962 these lakes dumped at the same time resulting in the highest measured flood on the Tazlina River. Strandlines above Iceberg Lake indicate that the lake has been about 100 meters (300 feet) higher in recent decades. Extreme flood hazard in Tazlina lowlands, moderate flood hazard in Copper River valley. Monitoring is recommended. (See lake No. 16)	Map, Valdez (C-7 and C-8), Alaska Post (1967, table 5), Raple, Sater and Field (1965a, p.18-19, 29-30), Stone (1963a) U.S. Geological Survey gaging stations 15-2020 and 2120											
	16	Unamed, south	2.6	1.0	Nelchina	Nelchina River, Tazlina Lake, Tazlina River, Copper River	Two lakes drain subglacially, probably at 2- to 4-year intervals. Extreme flood hazard in Nelchina River and moderate flood hazard in Tazlina River lowlands. If combined with simultaneous floods from lakes No. 15 extremely hazardous floods could occur on the Tazlina and Copper River lowlands. Monitoring is recommended.	Map, Valdez (C-8), Alaska Post (1967, table 5), Raple, Sater and Field (1965a, p.18-19, 29-30), Stone (1963a) U.S. Geological Survey gaging stations 15-2020 and 2120											
	17	Lower Skolai Lake	1.0	.4	Nizina	Nizina River, Chinika River, Copper River	A lake 1 kilometer (0.6 mile) long, which drains along glacier margin, has formed infrequently in recent years. Cape (1916) reported "The glacier close the subglacial outlet of this lake, which then rises rapidly until the hydraulic pressure is sufficient to reopen a channel beneath the ice. Once opened, the lake waters pour out with a rush, flooding Nizina Valley below and leaving icebergs stranded high on the sides of the deserted lake basin." Moffitt (1930) stated "A tunc much timber is destroyed by the cutting away of wooded gravel benches. The bars of the upper Nizina River were piled up with tangled masses of trees brought down by the flood of 1927." Outburst in June 1934 demolished a bridge across the Nizina River. Moderate flood hazard in Nizina River lowlands. Monitoring is recommended.	Map, McCarthy (C-4), Alaska Alaska Department of Highways (1970, p.9), Capps (1916, p.15, p.14), Hayes (1892, p.135, 154), Moffitt (1938, p.145)											
	18	Hidden Creek Lake	2.0	.8	Kennicott	Kennicott River, Chinika River, Copper River	Lake drains subglacially. Water from this lake has been observed to emerge from the "bottle" at the lower end of Kennicott Glacier since early 1900s. "In winter *** a torrent of water rushes down the Kennicott and Nizina Rivers, sometimes flooding the ice all the way to the Copper River" (Moffitt and Capps, 1911). A surge of water over the ice on the Chinika River in March 1904 (J. McKechnie, oral commun., 1970) may have been due to a release of this lake. Moderate to extreme flood hazard on Kennicott River and moderate flood hazard on Chinika River flood plains.	Map, McCarthy (C-6), Alaska Bateman (1922, p.536), Moffitt (1938, p.13, pl. 5A), Moffitt and Capps (1911), Stone (1963a) U.S. Geological Survey gaging station 15-2120											

GLACIER DAMMED LAKES AND OUTBURST FLOODS IN ALASKA

By
Austin Post and Lawrence R. Mayo
1971