DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

MISCELLANEOUS GEOLOGIC INVESTIGATIONS
MAP I-573



Base from U.S. Geological Survey, 1956

SCALE 1:250 000

5 0 5 10 15 MILES
5 0 5 10 15 KILOMETERS

CONTOUR INTERVAL 200 FEET
DATUM IS MEAN SEA LEVEL
1956 MAGNETIC DECLINATION AT SOUTH EDGE OF SHEET VARIES FROM 30°30' TO 33° EAST

Bedrock geology by Earl E. Brabb (1960-67) and Michael Churkin, Jr. (1962-63); assisted by R. N. Passero and H. J. Roepke (1960), J. C. Melik and R. L. Taylor (1961), Arne Aadland (1962), and D. L. Giles and R.H. Rohrbacher (1963). Surficial geology by Arne Aadland (1962) and Earl E. Brabb (1963)



INDEX MAP OF ALASKA



Note: surficial deposits omitted from sections



LOCATION INDEX

INDEX MAP OF CHARLEY RIVER QUADRANGLE SHOWING LOCATION OF FIELD OBSERVATION POINTS

Bedrock contacts and structures have been extended between field stations by aerial reconnaissance and by interpretation of aerial photographs. The surficial deposits, which cover about one-third of the quadrangle, have been mapped mostly from aerial photographs.



GEOLOGIC MAP OF THE CHARLEY RIVER QUADRANGLE, EAST-CENTRAL ALASKA

By
Earl E. Brabb and Michael Churkin, Jr.

1969

EXPLANATION

QUATERNARY

Qfp / Qaf / Qt / Ql — Surficial deposits
Qfp, flood-plain deposits
Qaf, alluvial fan deposits
Qt, river terrace deposits
Ql, loess

CRETACEOUS AND TERTIARY?

TKs — Sandstone, mudstone, and conglomerate — *Upper Cretaceous to Pliocene(?)*
Sandstone and conglomerate, poorly sorted, friable, composed mostly of lithic fragments and about 5 percent potash feldspar. Thin, lignitic coal beds common. Nonmarine. Thickness 800 to at least 8,000 feet

UNCONFORMITY

NORTH OF TINTINA FAULT ZONE
STRATIGRAPHIC POSITION REASONABLY CERTAIN

Kka — Kathul Graywacke — *Lower Cretaceous*
Sandstone, conglomerate, and argillite. Sandstone, dark-greenish-gray, feldspathic (about 25 percent potash and plagioclase feldspar) graywacke with secondary chlorite and epidote. Conglomerate clasts of chert, volcanic rocks, argillite, and sandstone in matrix of graywacke or argillite. Argillaceous rocks mainly dark-gray argillite but include olive-gray shale and mudstone. Contains a few stratigraphically long-ranging marine pelecypods. Seems correlative with nonmarine rocks in Eagle quadrangle to the south which contain plants of probable Albian age. Thickness several thousand feet. Named and described by Brabb (1965)

LOCAL UNCONFORMITY

Kb — Biederman Argillite — *Lower Cretaceous*
Rhythmically interbedded, dark-gray argillite and medium-gray siltstone and sandstone. Siltstone and sandstone mainly quartz arenite commonly with carbonate cement. Conglade structures and cross-laminations common. Includes a few beds of chert-pebble conglomerate, siliceous shale, chert-limestone breccia and limestone. Contains very few pelecypods of Valanginian age near base. Thickness at least 5,000 feet. Named and described by Brabb (1965)

Kkw — Keenan Quartzite — *Lower Cretaceous*
Medium-gray massive quartzite and sandstone with a few interbeds of dark-gray siltstone and argillite. Weathers white and forms resistant ridges. Locally contains abundant Buchia "subtaria" of Valanginian age. Thickness varies from 100 to 1,000 feet. Named and described by Brabb (1965)

TRIASSIC TO CRETACEOUS

Kkg — Glenn Shale — *Middle Triassic to Lower Cretaceous*
Mainly grayish-black carbonaceous shale. Minor siltstone and quartzite. Phyllite and quartz-chlorite semischist in vicinity of Indian Grave Mountain are probably local metamorphic facies. Lower few hundred feet of Glenn Shale near Nation River is mostly dark-gray limestone and oil shale with Monotis, Halobia, Daonella, Discophyllites, and Nathorstites of Middle Triassic (Ladinian) to Late Triassic (Norian) age. At other localities contains Otapiria of probable Jurassic age (Imlay, 1967, p. B7), Buchia okensis of Early Cretaceous (Berriasian) age, and Buchia cf. B. crassicollis of Early Cretaceous (probably Valanginian) age. Sparsely fossiliferous north of 65°30'N. Maximum thickness about 5,000 feet. Named and described by Brabb (1965)

UNCONFORMITY(?)

PERMIAN

Pt — Tahkandit Limestone — *Upper Pennsylvanian to Permian*
Very pale orange, coarse-grained bioclastic limestone. Forms massive cliffs. Grayish-green glauconitic sandstone and chert-pebble conglomerate in basal 5 feet. Contains abundant brachiopods and a few corals, ammonoids, foraminifers and bryozoa. Maximum thickness about 350 feet

Ps — Step Conglomerate
Chert-pebble conglomerate and chert arenite. Minor siltstone, bioclastic limestone, and quartz arenite. Has Tahkandit fauna in Step Mountains area. Nonfossiliferous and of uncertain age in western and northern part of quadrangle except for single Permian fauna from sandstone near Tahoma Bluff. Thickness about 2,000 feet. Named and described by Brabb (1969)

UNCONFORMITY

MISSISSIPPIAN

Mcb — Calico Bluff Formation — *Upper Mississippian*
Rhythmically interbedded, dark-brownish-gray and brownish-black limestone and shale. Maximum thickness approximately 1,000 feet

MDf — Ford Lake Shale — *Upper Devonian to Upper Mississippian*
Grayish-black laminated siliceous shale and chert. Minor dark-gray siltstone, quartzite and limestone. Plant stems common. Thickness about 1,000 feet. Named and described by Brabb (1969)

Dnr — Nation River Formation — *Upper Devonian*
Rhythmically interbedded, mudstone, sandstone, gritstone, and conglomerate. Graded beds common. Mudstone is olive gray and nearly everywhere contains plant fragments and spores of Late Devonian age. Sandstone is olive-gray chert-quartz arenite of late Devonian age. Contains siltstone, limestone, and conglomerate composed mostly of recrystallized chert granules and pebbles. Thickness approximately 2,000 to 4,000 feet. See reports by Brabb and Churkin (1967) and Scott and Dober (1967)

DEVONIAN

Dm — McCann Hill Chert — *Lower, Middle(?), and Upper Devonian*
Thin-bedded and laminated dark- to light-gray chert, siliceous shale, and minor chert gritstone. Contains plant fragments, poorly preserved spores, and rare conodonts, gastropods, and cephalopods. Basal part of formation has beds of dark-gray bioclastic limestone with remarkably varied fauna including corals, brachiopods, trilobites, tentaculitids, fish, and ostracods of Early Devonian (Emsian) age. Thickness 800 to 850 feet except near Jones Ridge where it may be a few thousand feet thick. Named and described by Churkin and Brabb (1965a). Additional faunas listed by Churkin and Brabb (1967)

Dl — Limestone
Medium-gray limestone and calcareous shale. Contains a few brachiopods, corals, and bryozoa of Devonian age. Thickness a few hundred feet

DISCONFORMITY

DSOr — Road River Formation — *Lower Ordovician to Lower Devonian*
Dark-gray graptolitic shale with lesser amounts of grayish-black laminated chert and very minor dark-gray limestone, greenish-gray dolomitic, chert arenite, and conglomerate. Chert, chert arenite, and chert conglomerate occur mainly in basal part of formation. Graptolites indicate that most series of the Ordovician and Silurian are represented, and that the youngest rocks are Early Devonian (pre-Emsian). Thickness 200 to 900 feet. See reports by Churkin (1966) and Churkin and Brabb (1965a, 1967)

DISCONFORMITY

OCh — Hillard Limestone — *Lower Cambrian to Lower Ordovician*
Very fine to medium-grained pale-yellowish-brown limestone. Laminated to very thick bedded, shaly to massive parting. Has interbeds of edgewise limestone conglomerate, oolitic limestone, and sandy limestone. Minor dolomite, grayish-black chert, dark-gray faustic shale and siltstone, and a basal limestone boulder conglomerate. Fossils, especially trilobites and brachiopods, are common. Age, Early, Middle, and Late Cambrian and (near Tatonduk River) earliest Ordovician. Thickness 100 to 500 feet. See reports by Brabb (1967) and Palmer (1968)

Ca — Adams Argillite — *Lower Cambrian*
Light-olive-gray argillite, siltstone, and cross-laminated quartzite. Worm(?) burrows occur in quartzite; Oldhamia (see Churkin and Brabb, 1965b) in argillite. Oolitic and sandy limestone near base has archaeocyaths, thids, and trilobites of Early Cambrian age (Palmer 1968). Thickness 300 to 600 feet. Named and described by Brabb (1967)

Cf — Funnel Creek Limestone — *Lower Cambrian*
Light-gray fine- to medium-grained laminated limestone and dolomite. Extensively silicified and commonly oolitic. Forms massive cliffs. Non-fossiliferous. Thickness 50 to approximately 1,800 feet. Named and described by Brabb (1967)

OCj — Jones Ridge Limestone — *Lower Cambrian to Lower Ordovician*
Lower 8,000 feet of lower member is mainly very light gray, massive limestone and dolomite. Extensively silicified and commonly oolitic, pisolitic, and laminated. Upper 940 feet of lower member is pale-yellowish-brown, very fine grained, thin- to thick-bedded limestone. Upper member 60 feet thick is pale-yellowish-brown thick-bedded bioclastic limestone.
Lower member has archaeocyathids of Early Cambrian age and trilobites and other fossils of Late Cambrian and Early Ordovician age. Upper member has rich fauna of Middle or Late Ordovician age. Age of Jones Ridge is therefore considered to be Early Cambrian to Middle or Late Ordovician. Total thickness about 8,000 feet. Named and described by Brabb (1967). Fossils described by Palmer (1968) and Ross and Dutro (1966)

PRECAMBRIAN

br — Basalt and red beds — *Upper Precambrian to Tindir Group*
Dark-greenish-gray basalt, commonly amygdaloidal and with pillow structures. Feldspar and pyroxene mainly unaltered but at a few localities they are chloritized, and the rock is a greenstone. Minor basaltic tuff with pebbles and cobbles of basalt. Red beds mostly grayish-red shale and siliceous shale. Minor greenish-gray shale and siliceous shale, jasper, greenstone-dolomite conglomerate, hematitic dolomarenite, and vitric tuff and lava largely replaced by hematite and carbonate. Thickness 300 to at least 8,500 feet. Units C, D, and part of G of Mertie (1938)

ds / Dd — Dolomite and shale
Light- to medium-gray laminated dolomite and grayish-black shale. Minor chert, dolarenite, and dolomite-chert gritstone. The dolomite weathers very pale orange and is cut by diabase dikes. Thickness about 2,000 feet. Part of units E, F, and G of Mertie (1938)

sh — Shale
Grayish-black carbonaceous shale with minor interbeds of quartzite, dark-gray limestone, oolitic limestone, laminated dolomite, massive dolomite, and dolomite conglomerate. Cut by diabase dikes. Shale is a few thousand feet thick. Part of units E, F, and G of Mertie (1938)

pCt — Tindir Group, undivided — *Mainly dolomite and shale*

STRATIGRAPHIC POSITION UNCERTAIN

hs — Argillite
Dark-gray argillite and chert with minor interbeds of sandstone (chert and quartz arenite) and chert-pebble conglomerate. Contains Oldhamia. Grades into Paleozoic age near intersection of Wood and Black Rivers. May include some Precambrian and Mesozoic rocks. The argillite is probably several thousand feet thick
Pzl; limestone and dolomite. Contains a few poorly preserved brachiopods. Few hundred feet thick

Pc — Circle Volcanics
Dark-greenish-gray basalt, diabase, and gabbro. Minor interbedded chert, quartzite, and argillite. Nonfossiliferous but considered Early Mississippian by Mertie (1969) based on correlation with a sparsely fossiliferous volcanic sequence near Rampart, about 175 miles east of the Charley River quadrangle. Morris A. Lanphere of the U.S. Geological Survey (written commun., 1968) obtained a minimum potassium-argon date of about 220 million years for an intrusive(?) rock within the Circle Volcanics and a minimum date of about 200 million years for a similar intrusive(?) within the volcanic sequence near Rampart indicating that these volcanic rocks could be as young as Triassic. Pending further study, the Circle Volcanics are here considered provisionally to be of late Paleozoic age

SILURIAN AND DEVONIAN / ORDOVICIAN / CAMBRIAN — *Middle Devonian*

Woodchopper Volcanics — DEVONIAN
Mostly dark-greenish-gray spilitic basalt and lavalik (q.f. Amygdules and pillow structures common. Minor thin interbeds(?) of shale, chert, sandstone, siltstone, limestone, and dolomite. Probably a few thousand feet thick. Age uncertain.
Dwc; dark-gray laminated dolomite with minor interbeds of chert. Age uncertain. Probably several hundred feet thick
Dwl; dark-gray bioclastic limestone with brachiopods, corals, and other fossils of Middle Devonian age. The relation of the limestone to the volcanic rocks is uncertain. The limestone is a few hundred feet thick
Dws; dark-gray laminated shale with minor interbeds of limestone. Age uncertain. The shale is several hundred feet thick

pCd — Dolomite — PRECAMBRIAN(?)
Light-gray laminated dolomite. Chert laminae common. Minor dolomite-cobble conglomerate, sandy dolomite, laminated limestone, sandy limestone, shale, argillite, and quartzite. Probably Precambrian but could be Paleozoic. Few thousand feet thick

pCv — Volcanic rocks — PRECAMBRIAN(?)
Amygdaloidal basalt flows, pillow lavas, tuff breccia, tuff, and minor interbeds of dolomite, limestone, chert, argillite, and quartzite. Probably Precambrian but could be Paleozoic. Few thousand feet thick

pCu — Sedimentary rocks, undivided — PRECAMBRIAN(?)
Dark-gray laminated limestone and sandy limestone; olive-gray and greenish-red shale, argillite, and phyllitic argillite; chert, chert-pebble, and quartzite; and laminated dolomite. Includes a diabase dike along Edwards Creek. Probably Precambrian but may be Paleozoic. Few thousand feet thick

SOUTH OF TINTINA FAULT ZONE

Ma — Adamellite — MESOZOIC(?)
Medium- to coarse-grained adamellite. Biotite is chief accessory mineral. Hornblende less abundant. Muscovite and garnet occur locally. Adamellite has xenoliths of schist and is cut by aplite dikes. The adamellite forms generally structureless bodies discordant with surrounding schist, but in some places the adamellite seems to grade into quartz biotite gneiss

AGE RELATIONSHIPS UNCERTAIN

gn — Gneiss — PRECAMBRIAN(?) AND PALEOZOIC(?)
Quartz-plagioclase-biotite or muscovite gneiss

sp — Serpentine
Greenish-black antigorite

ms — Mica schist
Mostly quartz-plagioclase-biotite or muscovite schist. Includes a few beds of white marble as much as 25 feet thick. The schist has thin interlayers of, or is closely associated with amphibolite that has the chemical composition of basalt. Chlorite is a common constituent in all of the rocks. Garnet, andalusite, and kaolin occur locally, particularly near the adamellite. The unit is probably a sandstone or graywacke sequence with interbeds of limestone, shale, tuffaceous sedimentary rock and basalt that has been regionally metamorphosed

ph — Phyllite
Mainly fine-grained grayish-black and grayish-green phyllite. Minor massive dark-gray chert and stretched pebble-conglomerate. The unit was originally a carbonaceous and siliceous shale with a few interbeds of chert and chert-pebble conglomerate

gs / gsl — Greenstone
Fine-grained massive greenish-gray greenstone composed mostly of amphibole, plagioclase, chlorite, epidote, and minor quartz. Chemical analyses indicate that the rock was originally a basalt. At a few localities the greenstone is well foliated
gsl; coarse-grained massive greenstone similar in composition to the fine-grained greenstone. Represents basic intrusive rocks that were regionally metamorphosed

Contact, approximately located

Fault
Dashed where approximately located; dotted where concealed. U, upthrown side; D, downthrown side

Fault zone
Inferred and approximately located; concealed beneath sedimentary rocks and alluvium

FOLDS

Anticline showing trace of axial plane and plunge of axis

Syncline showing trace of axial plane and plunge of axis

Plunge of minor anticline

Plunge of minor syncline

BEDDING

Field measurement — Photogeologic interpretation or distant measurement from helicopter

Strike and dip of beds

Strike of vertical beds

Strike and dip of overturned beds

Horizontal beds

FOLIATION

Field measurement — Photogeologic interpretation or distant measurement from helicopter

Strike and dip of foliation

Strike of vertical foliation

Horizontal foliation

Identification of fossils
(Identification and age of fossils by geologists in the U.S. Geological Survey, unless otherwise noted)
J. M. Berdan (Ordovician, Silurian, and Devonian ostracods)
H. R. Bergquist (Mesozoic foraminifers)
W. R. N. Berry (Ordovician and Silurian graptolites)
Michael Churkin, Jr. (Ordovician, Silurian and Devonian graptolites, tentaculitids and phyllocarid crustaceans; Cambrian trace fossils)
P. E. Cloud, Jr. (Cambrian trace fossils)
D. H. Dunkle, U.S. National Museum (Devonian fish)
H. M. Duncan (Devonian hyrnanid)
J. T. Dutro, Jr. (Devonian brachiopods)
Mackenzie Gordon, Jr. (Paleozoic cephalopods)
J. W. Huddle (Ordovician and Devonian conodonts)
R. E. Grant (Permian brachiopods)
R. W. Imlay (Jurassic pelecypods)
D. L. Jones (Cretaceous mollusks)
S. H. Mamay (Paleozoic plants)
W. A. Oliver, Jr. (Ordovician and Devonian corals)
A. R. Ormiston, Harvard College (Devonian trilobites)
A. R. Palmer (Cambrian trilobites, brachiopods and archaeocyathids)
John Pojeta, Jr. (Paleozoic pelecypods)
R. J. Ross, Jr. (Ordovician and Silurian trilobites, brachiopods and graptolites)
R. A. Scott (Phanerozoic spores and pollen)
N. J. Silberling (Triassic mollusks)
I. G. Sohn (Carboniferous conodonts)
R. H. Tschudy (Cretaceous spores and pollen)
J. A. Wolfe (Cretaceous and Tertiary plants)
E. L. Yochelson (Paleozoic gastropods)

REFERENCES

Brabb, E. E., 1967, Stratigraphy of the Cambrian and Ordovician rocks of east-central Alaska: U.S. Geol. Survey Prof. Paper 559–A, 30 p., 16 figs.

———1969, Six new Paleozoic and Mesozoic formations in east-central Alaska: U.S. Geol. Survey Bull., 1274–I.

Brabb, E. E., and Churkin, Michael, Jr., 1967, Stratigraphic evidence for the Late Devonian age of the Nation River Formation, east-central Alaska, in Geological Survey research 1967: U.S. Geol. Survey Prof. Paper 575–D, p. D4–D15, 5 figs.

Churkin, Michael, Jr., 1966, Morphology and stratigraphic range of the phyllocarid crustacean *Caryocaris* from Alaska and the Great Basin: Paleontology, v. 9, pt. 3, p. 371–380, 2 pls.

Churkin, Michael, Jr., and Brabb, E. E., 1965a, Ordovician, Silurian, and Devonian biostratigraphy of east-central Alaska: Am. Assoc. Petroleum Geologists Bull., v. 49, no. 2, p. 172–185, 4 figs., 3 tables.

———1965b, Occurrence and stratigraphic significance of *Oldhamia*, a Cambrian trace fossil, in east-central Alaska in Geological Survey research 1965: U.S. Geol. Survey Prof. Paper 525–D, p. D130–D124, 4 figs.

———1967, Devonian rocks of the Yukon-Porcupine Rivers area and their tectonic relation to other Devonian successes in Alaska: Alberta Soc. Petroleum Geologists, Internat. Symposium on the Devonian System, Calgary 1967, v. 2, p. 227–256.

Mertie, J. B., Jr., 1930, Geology of the Eagle-Circle district, Alaska: U.S. Geol. Survey Bull. 816, 168 p., 6 figs., 12 pls.

———1938, The Tatonduk-Nation district: U.S. Geological Survey Bull. 896–E, p. 547–648, 3 figs., 1 pl.

Palmer, A. R., 1968, Cambrian trilobites of east-central Alaska: U.S. Geol. Survey Prof. Paper 559–B, B115 p., 16 pls.

Ross, R. J., Jr., and Dutro, J. T., Jr., 1966, Silicified Ordovician brachiopods from east-central Alaska: Smithsonian Misc. Colln., v. 149, no. 7, 22 p., 3 pls.

Scott, R. A., and Dober, L. I., 1967, Palynological evidence for Devonian age of the Nation River Formation, east-central Alaska, in Geological Survey research 1967: U.S. Geol. Survey Prof. Paper 575–B, p. B45–B49, 3 figs.

Imlay, R. W., 1967, The Mesozoic pelecypods *Otapiria* Marwick and *Lupherella* Imlay, new genus, in the United States: U.S. Geol. Survey Prof. Paper 573–B, B11 p., 2 figs., 3 pls., 2 tables.

For sale by U.S. Geological Survey, Washington, D.C. 20242