

189.	<i>S. laeta</i> Rich. (including <i>S. ciliatosepala</i> Trautv.)	Shining starwort	x	x	x	x	x	x	x	x	x	0-3500	C	7	Sandy floodplains, coastal sandy beaches, alpine meadows. Habitat unknown, Umiat.	
190.	<i>S. laxmannii</i> Fisch.															
191.	<i>S. longipes</i> Goldie															
192.	<i>S. monantha</i> Hult.											2000-4000	S	7	Habitat unknown, Umiat. Dry alpine rubble slopes.	
BUTTERCUP FAMILY																
193.	<i>Aconitum delphinifolium</i> DC.	Monkshood	x	x	x	x	x	x	x	x	x	0-5000	C	7-8	Moist alpine slopes, usually growing with grasses.	
194.	<i>Anemone drummondii</i> Wats.	Anemone				H	x	x	x	x	x	350-3500	C	6	Moist sandbars, terrace meadows, rubble slopes; eastern portion of mountain front.	
195.	<i>A. multiceps</i> (Greene) Standl.	Alaska blue anemone	x			x						1000-2500	L	6	Dry rubble slopes; western portion of mountain front.	
196.	<i>A. narcissiflora</i> L.	Narcissus-flowered anemone										1500-3000	L	7	Dry alpine meadows; central portion of mountain front.	
197.	<i>A. parviflora</i> Michx.	Northern anemone	x	x	x	x	x	x	x	x	x	0-3500	C	6-7	Moist alpine meadows and valley slopes; sandbars.	
198.	<i>A. patens</i> L. ssp. <i>multifida</i> (Pritzell) Zamels	Pasque flower				H						500-1500	L	6-7	Dry hilltops; central portion of foothills.	
199.	<i>A. richardsonii</i> Hook.	Yellow anemone	x	x	x	x	x	x	x	x	x	0-4000	C	7	Stream margins in sun or shade, moist alpine meadows.	
200.	<i>Caltha palustris</i> L. var. <i>arctica</i> (R. Br.) Huth	Yellow marsh marigold														
201.	<i>Delphinium brachycentrum</i> Ledeb.	Dwarf larkspur	H	x	x	x	x	x	x	x	x	0-3500	C	7	Emergent aquatic in shallow, slowly moving water; with floating leaves at Pt. Barrow.	
202.	<i>D. glaucum</i> Wats.	Glaucous larkspur										0-4000	C	7-8	Alpine meadows, small damp stream valleys, rocky slopes.	
203.	<i>Ranunculus aquatilis</i> L. var. <i>eradicatus</i> Laest.	White water-buttercup										350	L	7	Aquatic in small oxbow lakes around Umiat (<i>R. reptans</i> in Hulten 1941-1950)	
204.	<i>R. flammula</i> L. var. <i>filiformis</i> Michx.	Creeping buttercup										350	L	7	Semiaquatic in mud; west end of Umiat airstrip.	
205.	<i>R. gelidus</i> Kar. & Kir.											2000	L	7	Dry hilltops; Nuka River.	
206.	<i>R. glacialis</i> L. var. <i>chamissonis</i> Schlecht.											4000	L	7	Rubble slope; Thunder Mountain.	
207.	<i>R. gmelini</i> DC. var. <i>yukonensis</i> (Brit.) Benson											0-3500	C	8	Emergent or submerged aquatic of lake margins.	
208.	<i>R. hyperboreus</i> Rottb.		x	x	x	x	x	x	x	x	x	1000-2500	S	8	Semiaquatic of sandy lake margins.	
209.	<i>R. lapponicus</i> L.	Lapland buttercup										100-2500	S	7	Moist mossy valleys and gullies.	
210.	<i>R. nivalis</i> L.	Snow buttercup	H	x	x	x	x	x	x	x	x	0-3000	C	6	Late snow areas, damp margins of small streams. (<i>R. nivalis</i> and <i>R. sulphureus</i> are sometimes very similar, and intermediate forms occur.)	
211.	<i>R. pallasi</i> Schlecht.	Pallas buttercup	x	x	H	x	x	x	x	x	x	0-2500	C	7-8	Emergent aquatic in lake margins.	
212.	<i>R. pedatifidus</i> J. E. Smith var. <i>affinis</i> (R. Br.) Benson	Northern buttercup										0-3500	C	6-7	Dry slopes.	
213.	<i>R. pygmaeus</i> Wahl.	Pygmy buttercup	x	x	x	x	x	x	x	x	x	0-3500	C	6-7	Late snow areas, coastal disturbed areas and sands.	
214.	<i>R. sabinei</i> R. Br.															
215.	<i>R. sulphureus</i> Soland. * <i>R. turneri</i> Greene	Sulphur buttercup	x	H	x	x	x	x	x	x	x	0-3500	C	6-7	Late snow areas, wet valleys of small alpine streams. (See <i>R. nivalis</i> note.) Habitat unknown; Arctic coast of Yukon Territory.	
216.	<i>Thalictrum alpinum</i> L.	Arctic meadow-rue										1000-4500	S	7	Moist alpine crevices in sandstone and limestone, wet terrace meadows.	
POPPY FAMILY																
217.	<i>Papaver alboroseum</i> Hult.		x	x								0-3500	L	7-8	Sand mounds at Alaktak; glacial moraine and outwash at Lake Schrader.	
218.	<i>P. macounii</i> Greene	Macoun poppy	H	x	x	x	x	x	x	x	x	0-5000	C	7	Sandbars, dry alpine meadows, crevices and rubble slopes of sandstone and shale.	
219.	<i>P. radicum</i> Rottb.	Arctic poppy	x	x	x	x	x	x	x	x	x	0-5000	C	7-8	Dry meadows, hilltops, sandbars, and outcrops.	
FUMEWORT FAMILY																
220.	<i>Corydalis pauciflora</i> (Steph.) Pers.	Few-flowered corydalis										500-3500	S	6-7	Damp mossy valleys of small alpine streams.	
MUSTARD FAMILY																
221.	<i>Arabis lyrata</i> L. ssp. <i>kamchatica</i> (Fisch.) Hult.	Rock-cress										350-500	S	6-7	Sandbars.	
222.	<i>Braya purpurea</i> (R. Br.) Bunge		H	x	x	x	x	x	x	x	x	50-2000	L	7	Sandy floodplains at Alaktak; limestone talus, Katakaturuk River.	
223.	<i>Cardamine bellidifolia</i> L.	Alpine cress	x									0-4500	S	7	Moist alpine meadows, sandbars.	
224.	<i>C. microphylla</i> Adams					H	x	x	x	x	x	1500-3500	S	7	Moist alpine meadows east of Canning River.	
225.	<i>C. pratensis</i> L.	Cuckoo flower	x	x	x	x	x	x	x	x	x	0-3500	C	7-8	Wet meadows and lake margins.	
226.	<i>C. purpurea</i> C. & S.	Purple bitter-cress										1000-2500	S	7	Open wet soil, north-facing alpine slopes; western mountain front.	
227.	<i>C. richardsonii</i> Hult.		x	x	x	x	x	x	x	x	x	0-4000	C	6-7	River terraces, grassy slopes, sedge meadows.	
228.	<i>Cochlearia officinalis</i> L.	Scurvy grass	H	x	x							0-100	L	6-7	Ocean beaches and sandy floodplains near the coast.	
229.	<i>Descurainia sophioides</i> (Fisch.) Schulz	Northern tansy mustard										0-2500	L	7-8	Shale outcrops and freshly eroded cutbanks.	
230.	<i>Draba alpina</i> L.	Alpine whitflow-grass	x	x	x	x	x	x	x	x	x	0-3500	S	7	Dry alpine meadows and dry slopes. Material partly with white flowers and partly with yellow flowers.	
231.	<i>D. chamissonis</i> G. Don											2000-3500	S	7	Dry alpine slopes and hilltops. (<i>D. caesia</i> in Hulten 1941-1950)	
232.	<i>D. cinerea</i> Adams		H	x	x							1000-3000	S	7	Dry slopes and hilltops.	
233.	<i>D. crassifolia</i> Grah.											350-3000	S	7	Dry slopes.	
234.	<i>D. exalata</i> E. Ekm.					H										
235.	<i>D. fladnizensis</i> Wulfen											350-3500	S	7	Habitat unknown; Icy Reef.	
236.	<i>D. glabella</i> Pursh		H	x	H	x	x	x	x	x	x	0-3500	C	7	Dry alpine slopes.	
237.	<i>D. lactea</i> , Adams		x	x	H	x	x	x	x	x	x	0-3000	S	7	Moist sandy cutbanks (also called <i>D. hirta</i> L.).	
238.	<i>D. lanceolata</i> Royle											50-2000	S	7	Coastal sandy beaches and dry alpine slopes.	
239.	<i>D. longipes</i> Raup											500-3500	S	7	Silty mounds along coast, cutbanks.	
240.	<i>D. macrocarpa</i> Adams		H	H												
241.	<i>D. nivalis</i> Lilj.											0-4000	S	7	Cutbanks and dry alpine slopes.	
242.	Note: <i>Ermannia borealis</i> (Greene) Hult.											2500-4500	L	7	Habitat unknown; Pt. Barrow and Camden Bay.	
243.	<i>Erysimum inconspicuum</i> (Wats) MacM.	Narrow-leaved wallflower										350-2000	S	7	Coastal sandy beaches, dry slopes, crevices in limestone. (Several other doubtful species of <i>Draba</i> occur in Arctic Alaska.)	
244.	<i>E. pallasi</i> (Pursh) Fern.	Pallas wallflower	H	x	x	x	x	x	x	x	x	350-4500	C	6-7	Barren eroding slopes of shale, and coarse limestone talus.	
245.	<i>Eutrema edwardsii</i> R. Br.											0-4000	C	6-7	Shale and gravel cutbanks; Umiat, Sadlerochit River, Shaviovik River.	
246.	<i>Lesquerella arctica</i> (Wormsk.) Wats.	Arctic bladder-pod	H	x	x	x	x	x	x	x	x	1000-3000	L	6-7	Dry sandbars, cutbanks, and rubble slopes.	
247.	<i>Parrya nudicaulis</i> (L.) Regel		x	x	x	x	x	x	x	x	x	0-4500	C	6-7	Arctic and alpine meadows.	
248.	<i>Rorippa palustris</i> (L.) Bess.	Yellow water-cress										350-1500	L	7	Well drained limestone pebble alluvial fans east of Canning River.	
249.	<i>Smelowskia calycina</i> (Steph.) C. A. Mey. var. <i>integrifolia</i> Drurey var. <i>media</i> Drurey var. <i>porsildii</i> Drurey	Siberian smelowskia										2000-5000	C	6-7	Moist meadow slopes, sandbars, rock crevices.	
250.	<i>Thlaspi arcticum</i> Pers.	Arctic penny-cress										1500	L	8	Sandy stream and lake margins.	
251.	<i>Torularia humilis</i> (C. A. Mey) Schulz	Northern rock-cress										350-800	S	7	Coarse, dry, alpine rubble slopes, and crevices in sandstone, limestone, and conglomerate. Var. <i>integrifolia</i> all along mountain front; var. <i>media</i> along eastern mountain front; var. <i>porsildii</i> in central mountain front near Anaktuvuk Pass. Only a few plants found; Sadlerochit River terrace.	
ORPINE FAMILY																
252.	<i>Sedum roseum</i> (L.) Scop. ssp. <i>integrifolium</i> (Raf.) Hult.	Rose root										0-2000	L	8	Sandbars, Umiat & Sagavanirktok River.	
SAXIFRAGE FAMILY																
253.	<i>Chryso-splenium towensis</i> Rydb.	Water-carpet	x	x								0-2000	L	7-8	Coastal sandy beaches, dry slopes, crevices in limestone. (Several other doubtful species of <i>Draba</i> occur in Arctic Alaska.)	
254.	<i>C. tetrandrum</i> Th. Fries	Northern water-carpet	x	x	x	x	x	x	x	x	x	0-4000	C	6-7	Barren eroding slopes of shale, and coarse limestone talus.	
255.	<i>C. wrightii</i> Franch. & Sav.	Bering Sea water-carpet										350-5000	S	5-7	Shale and gravel cutbanks; Umiat, Sadlerochit River, Shaviovik River.	
256.	<i>Parnassia kotzebuei</i> C. & S.	Kotzebue grass-of-parnassus	x	x	x	x	x	x	x	x	x	0-4000	C	7	Dry sandbars, cutbanks, and rubble slopes.	
257.	<i>P. palustris</i> L.	Northern grass-of-parnassus										350-3000	C	7	Arctic and alpine meadows.	
258.	<i>Saxifraga bronchialis</i> L. ssp. <i>funstonii</i> (Small) Hult.	Spotted saxifrage										350-6000	C	7	Well drained limestone pebble alluvial fans east of Canning River.	
259.	<i>S. caespitosa</i> L. ssp. <i>sileneiflora</i> (Sternb.) Hult.	Nodding saxifrage	H	x	H	x	x	x	x	x	x	0-5000	C	7	Moist meadow slopes, sandbars, rock crevices.	
260.	<i>S. cernua</i> L.		x	x	x	x	x	x	x	x	x	0-3500	C	7	Sandy stream and lake margins.	
261.	<i>S. davurica</i> Willd. ssp. <i>grandipetala</i> (Engl. & Irm.) Hult.	Purple saxifrage										1000-5000	C	7	Coarse, dry, alpine rubble slopes, and crevices in sandstone, limestone, and conglomerate.	
262.	<i>S. eschscholtzii</i> Sternb.	Ciliate saxifrage										2000-5500	C	7	Var. <i>integrifolia</i> all along mountain front; var. <i>media</i> along eastern mountain front; var. <i>porsildii</i> in central mountain front near Anaktuvuk Pass.	
263.	<i>S. flagellaris</i> Willd.	Flagellate saxifrage	x	H	x	x	x	x	x	x	x	0-5000	C	7	Only a few plants found; Sadlerochit River terrace.	
264.	<i>S. foliolosa</i> R. Br.		x	x	x	x	x	x	x	x	x	0-3000	S	7	Sandbars, Umiat & Sagavanirktok River.	
265.	<i>S. hieracifolia</i> Waldst. & Kit.	Hawkweed-leaved saxifrage	x	x	x	x	x	x	x	x	x	0-4000	C	7	Coarse, dry, alpine rubble slopes, and crevices in sandstone, limestone, and conglomerate.	
266.	<i>S. hirculus</i> L.	Yellow marsh saxifrage	x	x	x	x	x	x	x	x	x	0-4000	C	7	Dry alpine meadows, mountain slopes of limestone, sandstone, shale, conglomerate, and basalt.	
267.	<i>S. nivalis</i> L.	Alpine saxifrage										0-4000	S	7	Wet meadows, late snow areas.	
268.	<i>S. oppositifolia</i> L.	Purple mountain saxifrage	H	x	x	x	x	x	x	x	x	0-6000	C	6-7	Moist valleys and moist rolling meadows.	
269.	<i>S. punctata</i> L. ssp. <i>nelsoniana</i> (D. Don) Hult.	Brook saxifrage	x	x	x	x	x	x	x	x	x	0-4000	C	6-7	Wet sedge meadows.	
270.	<i>S. radiata</i> Small															
271.	<i>S. reflexa</i> Hook.	Yukon saxifrage										350-5000	C	6-7	Dry meadows and rubble slopes.	
272.	<i>S. rivularis</i> L.	Alpine brook saxifrage	x	x	x	x	x	x	x	x	x	0-4500	S	6-7	Moist rocky alpine slopes, crevices in sandstone, limestone, shale, schist, and basalt; late snow areas, silty frost boils in niggerhead meadows.	
273.	<i>S. serpyllifolia</i> Pursh * <i>S. spicata</i> D. Don	Thyme-leaved saxifrage										2000-6000	C	7	Valley and upland meadows, river terraces and cutbanks.	
274.	<i>S. tricuspidata</i> Retz.	Spiked saxifrage										2000	L	8	Habitat unknown; coastal.	
275.	<i>S. unalaschensis</i> Sternb.	Three-toothed saxifrage	x	x	x	x	x	x	x	x	x	350-4500	C	7	Dry rubble slopes, cutbanks, and crevices in sandstone, conglomerate, schist, and gabbro.	
276.	<i>Therophon richardsonii</i> (Hook.) Kze.	Richardson saxifrage										300-4000	C	7	Coastal sandy beaches, late snow areas in small alpine valleys.	
GOOSEBERRY FAMILY																
277.	<i>Ribes triste</i> Pallas	Red currant										350-2000	L	6-7	Moist alpine rubble slopes of sandstone and shale.	

U. S. Geological Survey
OPEN FILE REPORT
This report is preliminary and has not been edited or reviewed for conformity with Geological Survey standards or nomenclature.