DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

U.S. GEOLOGICAL SURVEY
OPEN FILE MAP
OF 77-166D

Base map from the National Atlas of the United States of America, U. S. Geological Survey, 1970

Albers Equal Area Projection

SCALE 1:1,000,000

10 0 20 40 60 80 100 MILES

10 0 50 100 150 KILOMETERS



MAP SHOWING GEOCHEMICAL ANOMALIES IN THE BROOKS RANGE, ALASKA

Compiled by Donald Grybeck

1977

REFERENCES

1. Brosgé, W. P., Reiser, H. N., and Tailleur, I. L., 1967, Copper analyses of selected samples, southwestern Brooks Range, Alaska: U.S. Geol. Survey open-file rept. 274, 1 sheet.

2. Pessel, G. H., 1976, Geochemistry of stream-sediment samples in southeastern Baird Mountains quadrangle, Alaska: Alaska Div. Geol. and Geophys. Surveys open-file rept. AOF-88, 8 sheets, scale 1:200,000.

3. Quadrangle systematically sampled and data now being processed under Alaska Mineral Resource Assessment Program (AMRAP). Geochemical results are scheduled for release in the summer of 1977 and will be included on a revision of this map in late 1977.

4. Garland, R. E., Eakins, G. R., and Trible, T. C., 1975a, Geochemical analysis of stream-sediment samples from part of the Survey Pass A-2 quadrangle, Alaska: Alaska Div. of Geol. and Geophysical Surveys open-file rept. 61, 2 p.

_____1975b, Geochemical analysis of stream-sediment samples from the Survey Pass B-3 quadrangle, Alaska: Alaska Div. Geol. and Geophysical Surveys open-file rept. 62, 2 p.

_____1975c, Geochemical analysis of rock and stream-sediment samples from the Survey Pass C-4 quadrangle, Alaska: Alaska Div. of Geol. and Geophysical Surveys open-file rept. 63, 2 p.

_____1975d, Geochemical analysis of stream-sediment samples from the Survey Pass C-5 quadrangle, Alaska: Alaska Div. of Geol. and Geophysical Surveys open-file rept. 64, 2 p.

_____1975e, Geochemical analysis of stream-sediment samples from the Survey Pass C-6 quadrangle, Alaska: Alaska Div. of Geol. and Geophysical Surveys open-file rept. 65, 2 p.

Garland, R. E., Eakins, G. R., Trible, T. C., and McClintock, W. W., 1975a, Geochemical analysis of rock and stream-sediment samples from the Survey Pass A-3 quadrangle, Alaska: Alaska Div. of Geol. and Geophysical Surveys open-file rept. 66, 2 p.

_____1975b, Geochemical analysis of rock and stream-sediment samples, Survey Pass A-4, A-5, A-6, B-4, B-5 and B-6 quadrangles, Alaska: Alaska Div. of Geol. and Geophysical Surveys open-file rept. 67, 2 p.

5. Elliott, R. L., and Miller, T. P., 1969, Results of stream-sediment sampling in the western Candle and southern Selawik quadrangles, Alaska: U.S. Geol. Survey open-file rept. 353, 61 p.

6. Miller, T. P., 1969, Results of stream sediment sampling in the northern Melozitna, the Hughes, and the southern Shungnak quadrangles, west-central Alaska: U.S. Geol. Survey open-file rept, 53 p.

7. Brosgé, W. P., and Reiser, H. N., 1970, Chemical analyses of stream sediment samples from the Chandalar and eastern Wiseman quadrangles, Alaska: U.S. Geol. Survey open-file rept., 56 p.

_____1972, Geochemical reconnaissance in the Wiseman and Chandalar districts and adjacent region, southern Brooks Range, Alaska: U.S. Geol. Survey Prof. Paper 709, 21 p.

8. Patton, W. W., Jr., and Miller, T. P., 1973, Analyses of stream-sediment samples from the Bettles and the southern part of the Wiseman quadrangles, Alaska: U.S. Geol. Survey open-file rept. 562, 52 p.

9-10. Quadrangle systematically sampled and data now being processed under Alaska Mineral Resource Assessment Program (AMRAP). Geochemical results are scheduled for release in the summer of 1977 and will be included on a revision of this map in late 1977.

11. Brosgé, W. P., and Reiser, H. N., 1976, Preliminary geologic and mineral resource maps (excluding petroleum), Arctic National Wildlife Range, Alaska: U.S. Geol. Survey open-file map 76-539, 4 sheets, scale 1:500,000.

12. Brosgé, W. P., Reiser, H. N., and Estlund, M. B., 1970, Chemical analyses of stream sediment samples from the Sadlerochit-Jago Rivers area, Mt. Michelson and Demarcation Point quadrangles, Alaska: U.S. Geol. Survey open-file rept. 409, 6 p.

13. Brosgé, W. P., and Reiser, H. N., 1968, Geochemical reconnaissance maps of granitic rocks, Coleen and Table Mountain quadrangles, Alaska: U.S. Geol. Survey open-file map, scale 1:250,000.

_____1977, Chemical analyses of stream-sediment samples from the Table Mountain and Arctic quadrangles, northern Alaska: U.S. Geol. Survey open-file rept. 77-29, 4 p.

EXPLANATION

This map attempts to summarize the scattered geochemical sampling in the Brooks Range as a tool in the mineral resource assessment of the area. Only stream sediment samples are considered.

In this report, anomalous is considered to be as (infrequently) defined by the authors of the reports cited or is established at the following values:

Silver (Ag):	≥	1 ppm
Gold (Au):	≥	0.5 ppm
Copper (Cu):	>	100 ppm
Molybdenum (Mo):	≥	20 ppm
Tin (Sn):	≥	50 ppm
Tungsten (W):	≥	100 ppm
Zinc (Zn):	≥	300 ppm

These values have been somewhat arbitrarily set high enough to exclude most uncertainty due to the accuracy and precision of the analytical techniques and to the variability in geochemical background. However, substantial mineralization frequently occurs without producing anomalies to these levels under a variety of conditions. Conversely, some of the anomalies may only represent high "normal" geochemical background or analytical error. Thus as in any interpretation of stream sediment data, the anomalies are subjective and a compromise that is here complicated by suggesting uniformity over such diverse geology and large area. The original sources should be consulted if a more refined interpretation or one with different levels of anomality seem more reasonable.

EXPLANATION OF SYMBOLS

Ag
Cu, Pb, Zn — Location of one or two closely-spaced, anomalous stream sediment samples. If more than one element is anomalous, the location of the sample is at the first elemental symbol.

Areas with a number of samples generally taken less than 5 kms apart which are anomalous in one or more metals.

(3) Outline of area with systematic network of stream sediment samples generally taken less than 5 kms apart. The number refers to the reference(s) pertaining to that area. (Note that the geochemical data is yet to be released in areas 3, 9, and 10).

(7) Outline of area with an erratic network of stream sediment samples or a few isolated samples. No inference should be made within these areas about the probable absence of deposits or area of influence of the anomalous sample(s) noted without consulting the original references. The number refers to the reference(s) pertaining to that area.

Boundary of Brooks Range Regional Mineral Resource Assessment Program (RAMRAP) as used in this study.

Outline of area selected under sec. 17(d)(2) of the Alaska Native Claims Settlement Act. Numbers refer to the Four National Systems proposal of the Department of the Interior, Dec. 18, 1973. Compiled from maps prepared by the U.S. Bureau of Land Management, December 1973 and March 1974.

1. Gates of the Arctic National Park
2. Kobuk Valley National Monument
3. Cape Krusenstern National Monument
13. Arctic National Wildlife Refuge Additions
15. Selawik National Wildlife Refuge
19. Noatak National Arctic Range

Approximate route of Alyeska pipeline

This map is preliminary and has not been reviewed for conformity with U. S. Geological Survey standards and nomenclature.