DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

FOLIO OF THE
GOODNEWS AND HAGEMEISTER ISLAND QUADRANGLES REGION

OPEN FILE REPORT 78-9-N

HESSIN AND OTHERS -- GEOCHEMICAL DISTRIBUTION OF LEAD

PROPERTY OF USGS LIBRARY
RESTON, VIRGINIA

HISTOGRAMS

PERCENTILE
95 97

THESE MANUALLY DERIVED HISTOGRAMS GRAPHICALLY ILLUSTRATE THE FREQUENCY AND ABUNDANCE OF LEAD CONCENTRATIONS REPORTED IN TABLES OF GEOCHEMICAL ANALYSES (HESSIN AND OTHERS, 1978; EAKINS, 1968, 1969). COMPUTER-GENERATED HISTOGRAMS AND RELATED STATISTICS FROM DATA OF HESSIN AND OTHERS (1978) ARE INCLUDED WITH THE GEOCHEMICAL ANALYSES IN THEIR REPORT.

PAN CONCENTRATE
NON-MAGNETIC FRACTION

MINUS-80-MESH STREAM SEDIMENT

(ATOMIC ABSORPTION)

MEAN: 44 PPM
MEAN: 18 PPM
MEAN: 23 PPM

(OXALIC ACID LEACHATE)

MEAN: 51 PPM

WILLOW LEAVES

(ATOMIC ABSORPTION)

MEAN: 15 PPM

FREQUENCY
PARTS PER MILLION

Mean 20-30
Wood River Lakes Area
(Eakins, 1968,Fig.7)
Anomalous over 50 ppm
NUMBER OF SAMPLES
CONCENTRATION(PPM)

(AREA OF REPORT: Barnes and others,1978)

(AREA OF REPORT: CLARK, GRYBECK. GREENWOOD and others. 1978
CLARK, GRYBECK, HESSIN and others, 1978)

(AREA OF REPORTS: Eakins, 1968 and 1969)

EXPLANATION

GEOLOGY GENERALIZED FROM HOARE AND COONRAD (1978)

CORRELATION OF MAP UNITS

SURFICIAL DEPOSITS

Qa	QUATERNARY

VOLCANIC, SEDIMENTARY, AND METAMORPHIC ROCKS

Qtb	Pleistocene → QUATERNARY
QTa	Plio-Pleistocene → QUATERNARY or TERTIARY
Tv	Early Tertiary → TERTIARY
Kk	Late Cretaceous Meadinchuan ?
Ks	Upper and Lower Cretaceous
Kn Kg Kis Kcg	Lower Cretaceous → CRETACEOUS
KJvs	Lower Cretaceous to Middle Jurassic → CRETACEOUS AND JURASSIC
Jk Jvs	Middle to Lower Upper Jurassic → JURASSIC
Jtv	Lower Jurassic → JURASSIC
MzPz	Lower Cretaceous to Lower Ordovician ? → MESOZOIC AND PALEOZOIC
Pzss	PALEOZOIC
pCk	PRE-CAMBRIAN

INTRUSIVE ROCKS

Td Tm Tx	TERTIARY
TKg	TERTIARY AND CRETACEOUS
Kg Jum Jt	JURASSIC AND JURASSIC ?
Pzmg	PALEOZOIC ?

DESCRIPTION OF MAP UNITS

SURFICIAL DEPOSITS

Qu UNCONSOLIDATED SEDIMENTARY DEPOSITS

SEDIMENTARY, VOLCANIC, AND METAMORPHIC ROCKS

Qtb TOKSHA BASALT

QTa UNCONSOLIDATED MARINE BEACH SEDIMENTS

Tv VOLCANIC ROCKS AND VOLCANOGENIC SEDIMENTS - Chiefly andesitic flows and tuffs

Kk SUMMIT ISLAND FORMATION - Nonmarine conglomerate, sandstone, shale, and carbonaceous mudstone

Ka HISKOKHOK GROUP - Conglomerate overlain by interbedded greywacke, siltstone, and shale; commonly calcareous; mostly marine

Ks BUCKIN RIDGE GRAYWACKE - Chiefly interbedded calcareous greywacke, siltstone, and conglomerate with local crinoidal beds of Aucella shells

Klg LIMY GRIT AND LIMESTONE - Chiefly angular grit cemented by bioclastic limestone

Kis TUFFS AND SEDIMENTARY ROCKS - Varied assemblage of andesitic tuffs, greywacke, siltstone, impure limestone, and tuffaceous chert; tuff and tuffaceous sediments commonly laumontitized

Kcg GRAYWACKE AND CONGLOMERATE - Marine greywacke, siltstone, and conglomerate; commonly calcareous

KJvs VOLCANIC AND SEDIMENTARY ROCKS - Interbedded intermediate to mafic flows, tuffs, tuffaceous sedimentary rocks, and argillite; intermediate composition tuffaceous rocks commonly laumontitized

Kn KULUKAK GRAYWACKE - Chiefly very hard lithic greywacke and siltstone with local conglomerate

Jvs VOLCANIC AND SEDIMENTARY ROCKS - Andesitic, trachytic, and basaltic flows and breccias interbedded with volcanogenic sedimentary rocks

Jtvs VOLCANIC AND SEDIMENTARY ROCKS - Mafic flows and breccias interbedded with volcanogenic sedimentary rocks; fractures commonly coated with lepidolite

MzPz MESOZOIC AND PALEOZOIC ROCKS UNDIVIDED - Widespread marine unit including mafic to intermediate volcanic rocks, tuffaceous sedimentary rocks, chert, argillite, siltstone, greywacke, conglomerate, and limestone

Mvs VOLCANIC AND SEDIMENTARY ROCKS (Upper Triassic) - Locally differentiated marine unit of chert, tuffaceous cherty rocks, argillite, siltstone, wacke, conglomerate, limestone, and mafic flows and breccia

Pv VOLCANIC ROCKS (Permian) - Locally differentiated marine unit of pillow basalts, massive mafic flows, breccia, and tuff

Pl LIMESTONE (Permian) - Locally differentiated thin unit of marine limestone; generally tuffaceous, commonly has strong fetid odor

DOS LIMESTONE - Thin-bedded to massive limestone with minor interbedded tuff and mafic flows; locally recrystallized to marble with interbedded quartzite and quartz-chlorite schist

Pzcs CALCAREOUS SCHIST (Permian or older?)

pCm KANAYUT METAMORPHIC COMPLEX - Gneiss, schist, amphibolite, and marble; upper greenschist to lower amphibolite facies

pCl MARBLEIZED LIMESTONE - Locally differentiated

INTRUSIVE ROCKS

Tri FELSIC INTRUSIVE ROCKS - Chiefly rhyolitic to dacitic dikes and sills; locally mapped

Tim MAFIC INTRUSIVE ROCKS - Diabase, basalt, dioritic, and gabbroic dikes and sills locally mapped

Tx WINDGAMM COMPLEX - Quartz-rich porphyritic felsite intrusive-extrusive complex of dikes, sills, tuff, and breccias

TKg GRANITIC ROCKS - Chiefly quartz monzonite, granodiorite, and quartz diorite stocks

Jv GABBROIC ROCKS - Commonly shows compositional layering and generally associated with ultramafic rocks

Jum ULTRAMAFIC ROCKS - Serpentinite, dunite, and websterite

Jt TRONDHJEMITE - Associated with serpentinite and gabbro

Pzmg METAGABBRO AND GREENSTONE - Probable dismembered ophiolite complex of mafic flows, dikes, volcaniclastic rocks, and gabbro altered by greenschist facies metamorphism and calcium metasomatism

GEOLOGIC SYMBOLS

Contact. Known, approximately located, gradational, and inferred. Most contacts between bedded rock units are probably faults

Fault or fault zone. Dashed where approximately located, inferred, or concealed

Thrust fault. Dashed where approximately located, inferred, or concealed. Sawteeth on upper plate

Hornfels

GEOCHEMICAL SYMBOLS

RED MOUNTAIN ULTRAMAFIC BODY

RED MOUNTAIN CONTACT ZONE

GEOCHEMICAL SAMPLE SITES

- ROCK (CLARK, GRYBECK, GREENWOOD, AND OTHERS, 1978; COONRAD, AND OTHERS, 1978)
- CONCENTRATE (OVERSTREET, AND OTHERS, 1973)
- OFFSHORE AND ONSHORE SEDIMENTS (BARNES, AND OTHERS, 1978)
- BEACH AND STREAM SEDIMENTS (BERRYHILL, 1963)
- STREAM DRAINAGE SEDIMENT (HESSIN, AND OTHERS, 1978)
- STREAM DRAINAGE SEDIMENT (CLARK, GRYBECK, HESSIN, AND OTHERS, 1978)
- STREAM DRAINAGE SEDIMENT (EAKINS, 1968, 1969)

ABUNDANCE

- ▼50 NUMBER WITH SOLID ROCK SAMPLE-SITE SYMBOL REPRESENTS GEOCHEMICAL CONCENTRATION IN PARTS PER MILLION (PPM); ✱ INDICATES ATOMIC-ABSORPTION)
- ● SOLID STREAM-DRAINAGE SAMPLE-SITE SYMBOL INDICATES GEOCHEMICAL ABUNDANCE OF 90TH PERCENTILE VALUE OR GREATER IN TWO OR MORE GEOCHEMICAL DETERMINATIONS AS SHOWN IN HISTOGRAMS

○ □ ○ ○ △ ◇ ABUNDANCE SYMBOLS REPRESENTING 95TH PERCENTILE OR GREATER CONCENTRATIONS DETERMINED IN STREAM-DRAINAGE SITE SAMPLES ARE SHOWN WITH HISTOGRAMS

DISCUSSION OF GEOCHEMISTRY

THIS REPORT SHOWS THE DISTRIBUTION AND ABUNDANCE OF LEAD AS GEOCHEMICALLY DETERMINED IN VARIOUS SAMPLE MEDIA COLLECTED FROM LOCATIONS THROUGHOUT THE GOODNEWS AND HAGEMEISTER ISLAND QUADRANGLES REGION. DATA PRESENTED HAVE BEEN COMPILED FROM ANALYSES AND LOCATIONS REPORTED BY BARNES AND OTHERS (1978); CLARK, GRYBECK, GREENWOOD, AND OTHERS (1978); CLARK, GRYBECK, HESSIN, AND OTHERS (1978); COONRAD AND OTHERS (1978); EAKINS (1968, 1969); HESSIN AND OTHERS (1978); AND OVERSTREET AND OTHERS (1973).

THE INCLUDED HISTOGRAMS HAVE BEEN USED TO IDENTIFY CONCENTRATIONS OF LEAD THAT MIGHT BE ANOMALOUS. HOWEVER, THE GENERALIZED GEOLOGIC MAP DATA INCLUDED IN THIS REPORT AND THE MORE DETAILED GEOLOGIC MAP OF THE REGION (HOARE AND COONRAD, 1978) SHOULD BE UTILIZED IN CONSIDERING POSSIBLE SOURCE ROCKS AND SIGNIFICANCE OF THE LEAD THAT HAS BEEN DETECTED IN VARIOUS GEOCHEMICAL SAMPLES.

REFERENCES CITED

Berryhill, R. V., 1963, Reconnaissance of beach sands, Bristol Bay, Alaska: U.S. Bur. Mines Rept. Inv. 6214, 48 p.

Barnes, P. W., Tagg, A. R., and Coonrad, W. L., 1978, Nearshore survey for platinum off Goodnews Bay, southwestern Alaska: U.S. Geol. Survey Open-file Rept. 78-9-E, 5 sheets, scale 1:125,000.

Clark, A. L., Grybeck, Donald, Greenwood, W. R., Taufen, P. M., Yount, M. E., and Coonrad, W. L., 1978, Geochemical analyses of rock samples in the vicinity of Platinum, Goodnews and Hagemeister Island quadrangles region, southwestern Alaska: U.S. Geol. Survey Open-file rept. 78-9-F, scale 1:125,000.

Clark, A. L., Grybeck, Donald, Hessin, T. D., Yount, M. E., and Coonrad, W. L., 1978, Geochemical analyses of stream-sediment samples in the vicinity of Platinum, Goodnews and Hagemeister Island quadrangles region, southwestern Alaska: U.S. Geol. Survey Open-file Rept. 78-9-G, scale 1:125,000.

Coonrad, W. L., Hoare, J. M., Taufen, P. M., and Hessin, T. D., 1978, Geochemical analyses of rock samples in the Goodnews and Hagemeister Island quadrangles region, southwestern Alaska: U.S. Geol. Survey Open-file Rept. 78-9-H, 2 sheets, scale 1:250,000.

Eakins, G. R., 1968, A geochemical investigation of the Wood River-Tikchik Lakes area, southwestern Alaska: Alaska Div. Mines and Minerals Geochem. Rept. 17, 31 p.

_____1969, A geochemical investigation of the Wood River-Tikchik Lakes area: Alaska Div. Mines and Geology Geochem. Rept. 17, Suppl., 6 p.

Hessin, T. D., O'Leary, R. M., Cooley, E. F., McDanal, S. K., McDougal, C. M., and Yount, M. E., 1978, Geochemical analyses of stream-sediment and vegetation samples in the Goodnews and Hagemeister Island quadrangles region, southwestern Alaska: U.S. Geol. Survey Open-file Rept. 78-9-I, scale 1:250,000.

Hoare, J. M., and Coonrad, W. L., 1978, Geologic map of the Goodnews and Hagemeister Island quadrangles region, southwestern Alaska: U.S. Geol. Survey Open-file Rept. 78-9-D, 1 sheet, scale 1:250,000.

Hoare, J. M., Coonrad, W. L., Hessin, T. D., Barnes, D. F., Clark, A. L., Grissom, Andrew, Singer, D. A., and Steele, W. C., 1978, The Alaskan Mineral Resources Assessment Program; background information to accompany folio of geologic, geochemical, geophysical, and mineral resource maps of the Goodnews and Hagemeister Island quadrangles region, southwestern Alaska: U.S. Geol. Survey Open-file Rept. 78-9.

Overstreet, W. C., Hamilton, J. C., Boerngen, J. G., Rosenblum, Sam Marsh, W. R., and Salnesburg, C. L., 1973, Minor elements in non-magnetic concentrates from Alaska: U.S. Geol. Survey open-file rept., 440 p. (also available NTIS (1975) PB-238 989/AS, 440 p.)

BASE MAP INDEX

Base compiled from U.S. Geological Survey 1:63,360 (A) and 1:250,000 (B) scale topographic series 1952-1957 and advance field completion material (C) 1976 by Menlo Park Base Map Section

SCALE 1:250 000
CONTOUR INTERVALS 50, 100 AND 200 FEET
DATUM IS MEAN SEA LEVEL

GEOCHEMICAL AND GENERALIZED GEOLOGICAL MAP SHOWING DISTRIBUTION AND ABUNDANCE OF LEAD IN THE GOODNEWS AND HAGEMEISTER ISLAND QUADRANGLES REGION, SOUTHWESTERN ALASKA

BY

T. D. HESSIN, P. M. TAUFEN, J. C. SEWARD, S. J. QUINTANA, A. L. CLARK, DONALD GRYBECK, J. M. HOARE, AND W. L. COONRAD

1978

BACKGROUND INFORMATION RELATING TO THIS REPORT IS PUBLISHED AS U.S. GEOLOGICAL SURVEY OPEN-FILE REPORT 78-9 (AVAILABLE FROM THE SAME SOURCE AS THE MAP)

This report is preliminary and has not been edited or reviewed for conformity with Geological Survey standards and nomenclature.