

**CORRELATION OF MAP UNITS**

Qa1	QUATERNARY
Tt	TERTIARY (?)
Kd	CRETACEOUS (?)
Kc	CRETACEOUS
Kf	CRETACEOUS
Kj	CRETACEOUS
Kk	CRETACEOUS
Kl	CRETACEOUS
Km	CRETACEOUS AND JURASSIC
Kn	CRETACEOUS
Ko	CRETACEOUS
Kp	CRETACEOUS
Kq	CRETACEOUS
Kr	CRETACEOUS
Ks	CRETACEOUS
Kt	CRETACEOUS
Ku	CRETACEOUS
Kv	CRETACEOUS
Kw	CRETACEOUS
Kx	CRETACEOUS
Ky	CRETACEOUS
Kz	CRETACEOUS
Ma	MESOZOIC AND PALEOZOIC

**LIST OF MAP UNITS**

Qa1	Alluvial deposits, undivided
Tt	Tertiary plutonic rocks
Kd	Mafic plutonic rocks
Kc	Diabase, extensively altered
Kf	Sites Group
Kj	Site Group
Kk	Site Group
Kl	Site Group
Km	Site Group
Kn	Site Group
Ko	Site Group
Kp	Site Group
Kq	Site Group
Kr	Site Group
Ks	Site Group
Kt	Site Group
Ku	Site Group
Kv	Site Group
Kw	Site Group
Kx	Site Group
Ky	Site Group
Kz	Site Group
Ma	Individual metamorphic, metasedimentary, and magmatic rocks

**SYMBOLS**

- Contact, approximately located, dotted where concealed
- Boundary of study area
- Geochemical sample site
- Copper (AA) = 200 ppm
- Copper (AA) = 300-500 ppm
- Copper (AA) > 500 ppm
- Copper (Spec) = 300 ppm
- Copper (Spec) = 500-700 ppm
- Copper (Spec) > 700 ppm

**STUDIES RELATED TO WILDERNESS**

The Wilderness Act (Public Law 88-577, September 3, 1964) and related acts require the U.S. Geological Survey and the U.S. Bureau of Mines to survey certain areas on Federal lands to determine their mineral resource potential. Results must be made available to the public and be submitted to the President and the Congress. This report presents the results of a geological survey of the Western Chichagof-Yakobi Islands Wilderness Study Area in the Tongass National Forest, Alaska. About 55 percent of the study area was established as a wilderness on December 7, 1980, under the Alaska National Interest Lands Conservation Act (P.L. 96-487).

In the course of the U.S. Geological Survey investigation of the Western Chichagof-Yakobi Islands Wilderness Study Area, 2,135 bedrock geochemical samples were collected. Samples were analyzed for 31 elements by a fast, sequential, semi-quantitative spectrographic method (Grimes and Morrison, 1965) and for 4 elements by atomic absorption spectrophotometry (Ward and others, 1969). Complete analytical data, element coordinates, and a station location map are available in two reports: Johnson, 1982, and Johnson and Elliott, 1984. A map and discussion of the mineral resource potential of the study area is also available (Johnson, Kibball, and Still, 1982).

Background levels for each element vary for different lithologies in the study area. Because of this and variability introduced from other sources such as sampling technique, analytical variation, and chemical weathering, it is impossible to select a specific analytical level above which values indicate mineralization. Higher values may indicate a greater likelihood of bedrock mineralization, but confidence levels are low for single element high values and results which are not supported by neighboring values. This map shows the distribution of high analytical values for the element copper by two analytical techniques as well as the location of all 2,135 samples. Multiple symbols for a single analytical technique at one sample site represent multiple samples at that site.

**REFERENCES CITED**

Grimes, D. J., and Morrison, A. F., 1965, Direct-current arc and alternating-current spark emission spectrographic field methods for the semi-quantitative analysis of geologic materials: U.S. Geological Survey Circular 591, 8 p.





Johnson, B. R., 1982, Magnetic tape containing trace element data for bedrock geochemical samples from the West Chichagof-Yakobi Islands Wilderness Study Area, southeastern Alaska: National Technical Information Service Report No. D85-02-82-005, computer tape, 1 reel.

Johnson, B. R., and Elliott, G. S., 1984, Map showing bedrock geochemical station locations, Western Chichagof-Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Open-File Report 81-0027-A, scale 1:125,000.

Johnson, B. R., and Still, J. W., 1982, Mesozoic geologic map of the Western Chichagof and Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1476-A, scale 1:125,000.

Johnson, B. R., Kibball, A. L., and Still, J. W., 1982, Mineral resource potential map of the Western Chichagof and Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1476-B, scale 1:125,000.

Ward, F. N., Nakagawa, M. N., Harris, T. F., and Van Stolle, G. R., 1969, Atomic absorption method of analysis useful in geochemical exploration: U.S. Geological Survey Bulletin 1289, 45 p.



MAP SHOWING THE DISTRIBUTION AND ABUNDANCE OF COPPER IN BEDROCK SAMPLES, WESTERN CHICHAGOF AND YAKOBI ISLANDS WILDERNESS STUDY AREA, SOUTHEASTERN ALASKA  
By  
Bruce R. Johnson and Geoffrey S. Elliott  
1984

This map is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards, but the stratigraphic nomenclature has been approved previously.