

CORRELATION OF MAP UNITS

Qal	QUATERNARY
T1, Tn	TERTIARY (?)
K2	CRETACEOUS (?)
K1	CRETACEOUS
K2, K1, J2, J1	CRETACEOUS AND JURASSIC
M2	MESOZOIC (?)
M1, P2, P1	MESOZOIC AND PALEOZOIC

LIST OF MAP UNITS

Qal	Alluvial deposits, undivided
T1	Tertiary plutonic rocks
Tn	Neofly plutonic rocks
K2	Metre silt, extensively altered
K1	Slate, gneiss
K2, K1	Granite Group
K1, J1	Plutonic plutonic rocks
K1, J1	Metre plutonic rocks
M2	Whitecap Marble
M1	Green Hill Gneiss
P2, P1	Undivided metamorphic, metasedimentary, and magmatic rocks

SYMBOLS

-----	Contact, approximately located, dotted where concealed
-----	Boundary of study area
*	Geochemical sample site
△	Strontium = 1500 ppm
△	Strontium > 1500 ppm
□	Lanthanum = 50 ppm
□	Lanthanum > 50 ppm

STUDIES RELATED TO WILDERNESS

The Wilderness Act (Public Law 89-377, September 3, 1964) and related acts require the U.S. Geological Survey and the U.S. Bureau of Mines to determine certain areas on Federal lands to determine their mineral resource potential. Results must be made available to the public and be submitted to the President and the Congress. This report presents the results of a geochemical survey of the Western Chichagof-Yakobi Islands Wilderness Study Area in the Tongass National Forest, Alaska. About 70 percent of the study area was established as a wilderness on December 2, 1980, under the Alaska National Interest Lands Conservation Act (P.L. 96-487).

Investigations of the Western Chichagof-Yakobi Islands Wilderness Study Area, 2,230 bedrock geochemical samples were collected. Samples were analyzed for 31 elements by a 6-step, semi-quantitative spectrographic method (Grimes and Morrison, 1968) and for 4 elements by atomic absorption spectrophotometry (Ward and others, 1969). Complete analytical data, station coordinates, and a station location map are available in two reports: Johnson, 1982, and Johnson and Elliott, 1984. A map and discussion of the mineral resource potential of the study area is also available (Johnson, Kimball, and Sillit, 1982).

Background levels for each element vary for different lithologies in the study area. Because of this and variability introduced from other sources such as sampling technique, analytical variance, and chemical weathering, it is impossible to select a specific analytical level above which values indicate mineralization. Higher values may indicate a greater likelihood of bedrock mineralization, but confidence levels are low for single element high values and results which are not supported by neighboring values. This map shows the distribution of high analytical values for the elements strontium and lanthanum as well as the location of all 2,230 samples. Multiple symbols for a single element at one sample site represent multiple samples at that site. Although not of contrast of elements themselves, these patterns, along with other, may prove to be useful in other elements. In a statistical analysis of similar geochemical data from Glacier Bay National Monument, high values for strontium and lanthanum were found to correlate with high values for zinc (Johnson, 1978).

REFERENCES CITED

Grimes, R. J., and Morrison, A. F., 1968, Direct-current arc and alternating-current spark emission spectrographic field methods for the semi-quantitative analysis of geological materials: U.S. Geological Survey Circular 591, 8 p.





Johnson, R. S., 1978, Statistical analysis of geochemical data from Glacier Bay National Monument, Alaska: U.S. Geological Survey Open-File Report 78-185, 18 p.

Johnson, R. S., 1982, Magnetic tape containing trace element data for bedrock geochemical samples from the West Chichagof-Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Open-File Report 81-0027-A, scale 1:125,000.

Johnson, R. S., and Sillit, G. S., 1982, Geochemical geologic map of the Western Chichagof and Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1474-A, scale 1:125,000.

Johnson, R. S., Kimball, A. L., and Sillit, G. S., 1982, Mineral resource potential map of the Western Chichagof and Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1476-A, scale 1:125,000.

Ward, J. W., Nakagawa, H. M., Harms, T. F., and Van Stickle, G. W., 1969, Atomic absorption methods of analysis useful in geochemical exploration: U.S. Geological Survey Bulletin 1289, 45 p.



MAP SHOWING THE DISTRIBUTION AND ABUNDANCE OF STRONTIUM AND LANTHANUM IN BEDROCK SAMPLES, WESTERN CHICHAGOF AND YAKOBI ISLANDS WILDERNESS STUDY AREA, SOUTHEASTERN ALASKA
By
Bruce R. Johnson and Geoffrey S. Elliott
1984

This map is preliminary and has not been reviewed for compliance with U.S. Geological Survey editorial standards, but the stratigraphic nomenclature has been approved previously.