

DISCUSSION

This map shows the distribution and abundance of copper, lead, zinc, and barium in 287 nonmagnetic, heavy-mineral concentrate samples collected during the 1978-1979 field seasons in the West Chichagof-Yakobi Wilderness Study Area, Sitka quadrangle, southeastern Alaska.

The area of this project is about 700 square miles (1800 km²). The heavy-mineral concentrate samples were taken from the active channels of streams draining areas of about 2 square miles (5 km²).

The preparation of the heavy-mineral concentrates prior to analysis involved panning stream sediment in the field to remove most of the low-density minerals. The panned sample was sieved through a 20-mesh (0.8 mm) sieve and the minus 20-mesh (0.8 mm) fraction was separated into light- and heavy-mineral fractions using bromoform (specific gravity of 2.96). The high density or heavy-mineral fraction was further prepared by removing magnetite and other strongly magnetic minerals by using a hand magnet.

The remaining heavy-mineral fraction was passed through a Frantz Isodynamic Separator at a setting of 0.6 amperes, thereby separating the magnetic fractions from the nonmagnetic. A split of the nonmagnetic fraction was pulverized with mortar and pestle and analyzed by semi-quantitative emission spectroscopy (Grimes and Marranzino, 1968).

The analytical data were compiled and used in preparing the histograms and geochemical map. The histograms were used to identify the range of anomalous values for each element.

A listing of results from the analysis of the various sample media collected in this Wilderness Study Area appears in Hessin and others, (1980).

REFERENCES CITED

- Grimes, D. J., and Marranzino, A. P., 1968, Direct-current and alternating-current spark emission spectrographic field methods for the semi-quantitative analysis of geologic materials: U.S. Geological Survey Circular 591, 6 p.
- Hessin, T. D., Smetan, V. S., Crenshaw, G. L., Hoffman, J. D., and Cooley, E. F., 1980, Analytical results of various types of samples taken in the West Chichagof-Yakobi Wilderness Study Area, Sitka quadrangle, southeastern Alaska, Open-File Report 80-905.

Use of brand names in this report is for descriptive purposes only and does not constitute endorsement by the U.S. Geological Survey.

Geochemical Symbols (size of symbol explained on histogram)

- △ Anomalous copper value (figure 1)
- Anomalous lead value (figure 2)
- Anomalous zinc value (figure 3)
- Anomalous barium value (figure 4)




Figure 1.--Histogram showing copper in 287, nonmagnetic, heavy-mineral concentrates from the West Chichagof-Yakobi Wilderness Study Area. Analysis by optical emission spectroscopy (Grimes and Marranzino, 1968). Triangles indicate anomalous concentrations and class percentages computed on total sample population.




Figure 2.--Histogram showing lead in 287, nonmagnetic, heavy-mineral concentrates from the Chichagof-Yakobi Wilderness Study Area. Analysis by optical emission spectroscopy (Grimes and Marranzino, 1968). Squares indicate anomalous concentrations and class percentages computed on total sample population.




Figure 3.--Histogram showing zinc in 287, nonmagnetic, heavy-mineral concentrates from the Chichagof-Yakobi Wilderness Study Area. Analysis by optical emission spectroscopy (Grimes and Marranzino, 1968). Hexagon indicates anomalous concentrations and class percentages computed on total sample population.




Figure 4.--Histogram showing barium in 287, nonmagnetic, heavy-mineral concentrates from the Chichagof-Yakobi Wilderness Study Area. Analysis by optical emission spectroscopy (Grimes and Marranzino, 1968). Circles indicate anomalous concentrations and class percentages computed on total sample population.

CORRELATION OF MAP UNITS

Qal	QUATERNARY
Tf	TERTIARY(?)
Kd	CRETACEOUS(?)
Ks	CRETACEOUS
Kkb	CRETACEOUS
Kjf	CRETACEOUS AND JURASSIC
Kjm	CRETACEOUS AND JURASSIC
Trw	TRIASSIC(?)
Trg	TRIASSIC(?)
MzPzu	MESOZOIC AND PALEOZOIC(?)

LIST OF MAP UNITS

Qal	ALLUVIAL DEPOSITS--Undivided
Tf	PELVIC PLUTONIC ROCKS--Dominantly tonalitic
Tm	MAFIC PLUTONIC ROCKS--Dominantly gabbroic
Kd	DIORITE SILL--Extensively altered
Ks	SITKA GRAYWAGE
Kkb	KELP BAY GROUP--Metasediments and metavolcanics
Kjf	PELVIC PLUTONIC ROCKS--Dominantly granodiorite
Kjm	MAFIC PLUTONIC ROCKS--Dominantly quartz diorite, diorite, and gabbro
Trw	WHITESTRIPE MARBLE
Trg	GOOSE DIP GREENSTONE
MzPzu	UNDIVIDED METASEDIMENTARY--Metavolcanic and metaplutonic rocks

GEOCHEMICAL MAP SHOWING DISTRIBUTION AND ABUNDANCE OF COPPER, LEAD, ZINC, AND BARIUM IN THE NONMAGNETIC, HEAVY-MINERAL CONCENTRATE SAMPLES IN THE WEST CHICHAGOF-YAKOBI WILDERNESS STUDY AREA, SITKA QUADRANGLE, SOUTHEASTERN ALASKA

By
T.D. Hessin, M.G. Maslowski and E.F. Cooley
1981

Studies Related to Wilderness
The Wilderness Act (Public Law 88-577, Sept. 3, 1964) and related Acts require the U.S. Geological Survey to survey certain areas on Federal lands to determine their mineral resource potential. Results must be made available to the public and be submitted to the President and the Congress. This report presents the results of a geochemical survey of the West Chichagof-Yakobi Wilderness Study Area, Sitka quadrangle, southeastern Alaska.

Base from U. S. Geological Survey 1:250,000, 1951
Sitka; Mt. Fairweather

