



OPHIOLITIC COMPLEXES OF THE GULF OF ALASKA by Steven W. Nelson

	A	B	C	D	E
	LITHOLOGY	THICK-NESS	AGE	MINERAL DEPOSITS ASSOCIATED WITH OPHIOLITIC TERRANES	GEOCHEMISTRY OF MAFIC, ULTRAMAFIC, AND ASSOCIATED ROCKS
Mchugh Complex	Disrupted and metamorphosed broken formation (melange) consisting of blocks (up to several kilometers in size) of chert, pillow basalt, marble, and schist in an argillaceous matrix, and a more coherent clastic section of graywacke, siltstone, and conglomerate. Prehnite-pumpellyite to greenschist-grade metamorphism. Structurally overlain by Border Ranges ultramafic and mafic complex locally. Clark, 1973; Connelly, 1976; Cowan and Boss, 1976; Burns, 1985.	Structural thickness up to 20km	Mesozoic Argillaceous matrix and chert blocks-Late Triassic (Carnian) to Late Cretaceous (Cenomanian) radiolarian ages. Marble blocks-Late Mississippian conodonts and Late Pennsylvanian fusulinids. K-Ar age of granitic clast from conglomerate of 150 Ma, and 161 Ma to 192 Ma ages from metamorphic rocks. Accretionary age early Late Cretaceous. Karl and others, 1979; Nelson and others, 1986, 1987; Winkler and others, 1981; Plafker and others, 1989.	No deposits known	Figure 1. AMF diagram of mafic volcanic blocks from the Mchugh Complex and correlative units in Alaska. Hill, 1979; Nelson and Blome, 1991; Plafker and others, 1989. Boundary line after Irvine and Baragar (1971). Figure 2. FeO*/MgO vs TiO <sub>2</sub> variation diagram for mafic volcanic rocks from the Mchugh Complex. Ol-ocean island; MORB-mid-ocean ridge basalt; IAT-island arc tholeiite, after Glassey, 1974. Nelson and Blome, 1991; Plafker and others, 1989.
Valdez Group	Predominantly interbedded sandstone, siltstone, and minor mudstone and conglomerate. This sequence of flysch consists of fan, slope, and basin-plain deposits and has a dominantly magmatic-arc provenance. Mafic volcanic rocks, consist of massive greenstone, pillow basalt, gabbro, tuffaceous units, minor sheeted dikes. Rare serpenitized dunite and peridotite. The entire sequence is extensively folded and faulted. Metamorphosed to grades ranging from zeolite to lower greenschist facies commonly and to amphibolite facies locally. Schrader, 1900; Budnik, 1974; Plafker and others, 1977; Zuffa and others, 1980; Nilsen and Zuffa, 1982; Dumoulin, 1987.	2 to 5km	Late Cretaceous Campanian (?) and Maestrichtian pelceopyods. Jones and Clark, 1973; Tysdal and Plafker, 1978.	Au-Gold-bearing quartz veins in flysch with associated placers. Cu-Massive sulfide deposit associated with mafic volcanic rocks. One major mine. Jansons and others, 1984; Nelson and others, 1984.	Figure 3. Range of chondrite-normalized REE distribution for blocks of mafic volcanic rock from the Mchugh Complex. Hill, 1979; Nelson and Blome, 1991; Plafker and others, 1989. Figure 4. Ternary diagram of (Hf/3)-Th-Ta for blocks of mafic volcanic rock from the Mchugh Complex and correlative units in Alaska. Boundaries after Wood, 1980. Hill, 1979; Nelson and Blome, 1991; Plafker and others, 1989.
Orca Group	Complexly deformed flysch and mafic igneous rocks. Sedimentary sequence composed of sandstone, siltstone, mudstone, and conglomerate with structures indicative of deposition by turbidity currents. Facies study indicates deposition in inner and middle parts of a complex system of submarine fans. Source area of sediments was plutonic-metamorphic roots of eroded arc. Mafic igneous rocks divided into Ophiolite sequence and an Outboard (non-ophiolite) sequence. Metamorphic grade ranges from laumontite to upper greenschist facies and local areas of hornblende hornfels. Grant and Higgins, 1910; Winkler, 1976; Winkler and Plafker, 1981; Nelson and others, 1985; Goldfarb and others, 1986; Dumoulin, 1987; Nelson, unpub.	2-3km 1-5km 1-5km 6-10km up to 30m 0-2km	Paleocene to middle Eocene Age based on microfossil and megafossil assemblages and intrusion by 53-50 Ma granitic plutons. Plafker and Langphere, 1974; Plafker and others, 1985; Nelson and others, 1985.  K-Ar whole rock dates Glacier Is. -42.5 ± 6.9 Ma Elamar -44.7 ± 5.0 Ma Miller, 1984.  U-Pb zircon date from plagiogranite Resurrection Peninsula -57 ± 1.0 Ma Nelson and others, 1989.	Cu, Au, and Ag-Produced in Prince William Sound area from volcanogenic massive sulfide deposits hosted in turbidites associated with ophiolite.  Knight Island -43 mines and prospects developed for copper. Resurrection Peninsula-8 prospects; known occurrences of Cu, Pb, Zn. Jansons and others, 1984; Nelson and others, 1984.	Figure 5. AMF diagram showing variation of rocks from the three sequences from Prince William Sound region. Nelson, unpub.; Plafker and others, 1989. Boundary line after Irvine and Baragar (1971). Figure 6. MgO-CaO-Al <sub>2</sub> O <sub>3</sub> variation diagram showing rocks from the three sequences from Prince William Sound region. Boundaries after Coleman, 1977. Nelson, unpub.; Plafker and others, 1989. Figure 7. Hf/3-Th-Ta discrimination diagram for rocks from the three sequences from Prince William Sound region. Data from Nelson, unpub.; and Plafker and others, 1989. CA-calc-alkaline, PAT-primitive arc tholeiite, WP-within plate, N-MORB-normal mid-ocean ridge basalt, E-MORB-enriched MORB (after Wood, 1980). Figure 8. Chondrite-normalized REE abundances of rocks from Prince William Sound region. Orca Group samples from Nelson, unpub.; Valdez Group samples from Nelson, unpub.; Plafker and others, 1989.
	Pillow Basalt-Includes hyaloclastite, massive basalt flows, breccia, and minor interbedded turbidite. Tholeiitic to basaltic andesite, porphyritic, vesicular, and amygdaloidal. Metamorphosed to prehnite-pumpellyite facies.  Sheeted Dike-Pyroxene gabbro to rare hornblende diorite; aphanitic to porphyritic and diabasic textures. Width of dikes varies from 10cm-2m, most vertical, although low angle cross cutting dikes locally present. Includes peridotite xenoliths. Dike orientation on Resurrection Peninsula-NS, W, and NW; on Knight Island NE, W, and NW.  Plagiogranite-On Knight Island, occurs as "lenses" within the sheeted dike unit; at Resurrection Peninsula intrudes gabbro and sheeted dikes, occurs as "lenses" in sheeted dikes, and is intruded by basalt dikes. Medium grained equigranular and porphyritic.  Gabbro-On Resurrection Peninsula, layered gabbro with alternating plagioclase-rich and clinopyroxene-rich layers and fault bounded block of coarse-grained gabbro with ophiolite and pegmatitic phases; includes xenoliths of partially serpenitized ultramafic rock. On Knight Island, gabbro occurs as small (<1km <sup>2</sup> ) bodies in sheeted dikes.  Ultramafic Rocks-On Knight Island, occur as xenoliths in sheeted dikes and larger (up to 200m x 400m) tectonic inclusions in Port Audrey shear zone (other inclusions include amphibolite, greenschist, siliceous mylonites, pillow basalt and metasediments). Olivine altered to serpentine, chlorite, and talc; clinopyroxene altered to actinolite, hornblende, and chlorite. Dunite, wehrlite, and olivine clinopyroxene found in float. On Resurrection Peninsula occur as xenoliths in gabbro (dunite and pyroxenite), float (harzburgite), and serpenitized peridotite bodies in turbidites. Richter, 1965; Winkler, 1973; Tysdal and others, 1977; Nelson and others, 1985, 1989.				

EXPLANATION

- |                                           |                                     |
|-------------------------------------------|-------------------------------------|
| Mudstone, shale                           | Pillow basalt                       |
| Argillite matrix in melange               | Massive basalt/greenstone           |
| Siltstone                                 | Pillow breccia                      |
| Sandstone(graywacke)                      | Sheeted dikes                       |
| Conglomerate                              | Plagiogranite                       |
| Limestone                                 | Clinopyroxene gabbro stocks         |
| Bedded chert                              | Layered gabbro                      |
| Greenschist, slate                        | Massive gabbro                      |
| Faults                                    | Serpenitized ultramafic             |
| Regional scale thrust fault (see fig. 12) | Ultramafic inclusions and xenoliths |
| Local normal fault                        |                                     |



**REGIONAL SETTING AND GEOLOGY**

Figure 12. Simplified map and section in southern Alaska showing geologic setting of ophiolite terranes of Gulf of Alaska. After Plafker and others, 1989.

Figure 13. Geologic map of the Resurrection Peninsula ophiolite.

Figure 14. Geologic map of the Knight Island ophiolite.

Figure 15. Cross sections through the Resurrection Peninsula and Knight Island ophiolites.

**Diagram symbols**

**Orca Group**

- Ophiolite sequence
  - Group 1-pillow basalt, sheeted dikes, gabbro
  - Group 2-plagiogranite
  - Group 3-diorite, pillowed basaltic andesite
  - Group 4-ultramafic rocks
- Outboard sequence
  - Massive basalt

**Valdez Group**

- Valdez Group ophiolite sequence
  - Pillow basalt, massive basalt, gabbro
  - Ultramafic rocks

**Geologic Explanation**

- Intrusive Rocks (Eocene and Oligocene)
  - Granite and Granodiorite
- Orca Group (Paleocene to middle Eocene)
  - Sedimentary Rocks
  - Pillow Basalt
  - Sedimentary and Volcanic Rocks
  - Sheeted Dikes
  - Gabbro
  - Ultramafic Rocks
  - Metasedimentary Rocks
- Other features:
  - Contact - Dashed where approximate
  - Fault - Dotted where concealed, dashed where inferred
  - Thrust Fault
  - Lineament - Inferred from well developed bathymetric contours
  - Port Audrey Shear Zone
  - Dip and Strike of Bedding
  - Dip and Strike of Cleavage
  - Dip and Strike of Dikes
  - Strike of Vertical Dikes

This report is preliminary and has not been edited or reviewed for conformity with U.S. Geological Survey editorial standards. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.