



U.S. DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY
in cooperation with
ALASKA DIVISION OF GEOLOGICAL AND GEOPHYSICAL SURVEYS

A	B	C	D	E
LITHOLOGY	THICKNESS (EST.)	AGE	MINERAL DEPOSITS	COMPOSITIONAL DATA
	Volcanogenic and sedimentary unit: Andesitic, dacitic, and minor basaltic tuffs, tuff breccias, flows, flow breccias, and domes of the Talkeetna Formation. Interbedded siltstone, and shale, and volcanoclastic sandstone and conglomerate are also present, particularly in the upper sections. Base not clearly exposed. The Talkeetna Formation is intruded by quartz diorite and tonalite with minor granodiorite and trondhjemite. Plutons are Jurassic in age.	> 6 km (?)	Late Triassic-Early Jurassic age determinations for megafossils within sedimentary units (Mittay and Detterman, 1973; Detterman and Reed, 1980). K-Ar ages on intermediate plutonic rocks are 163 to 194 Ma (Hill, 1979; Winkler and others, 1981; Pavlis, 1982b; Burns and others, 1991).	Ag, Cu, Pb, Zn - is interpreted to be the host rock for volcanogenic massive sulphide prospect (Johnson River) on the west side of Cook Inlet. Volcanogenic massive sulphide prospects in the Lake Clark quadrangle may be in the Talkeetna Formation. However, age control is poor, and the prospects may be in younger volcanic rocks (Newberry, pers. comm., 1990). Ag, minor Au - Geochemical indicators and subaerial volcanic rocks suggest that epithermal silver deposits may be present in the southern Talkeetna Mountains (Newberry, pers. comm., 1990).
	Sheeted dike complex: Composed of multiple intrusions (1 cm to 2 m thick) of diabase, gabbro, hornblende pyroxene gabbro, hornblende gabbro, and quartz diorite.	1 km (?)	No deposits reported	
	Massive gabbroic complex: Chiefly gabbro, leucogabbro, magnetite gabbro, and hornblende pyroxene gabbro. Intrudes amphibolite, schist, and minor marble; protolith of the metamorphic rocks is interpreted to be dominantly mafic volcanic rocks, metachert, and minor impure limestones. These metamorphic rocks are also intruded by quartz diorites and trondhjemites.	4 km	K-Ar dates on hornblende from gabbroic rocks range from 154 to 188 Ma (Winkler and others, 1981) and are interpreted to represent cooling ages of young phases. Quartz diorite ages as given above. K-Ar ages on biotite, hornblende, and muscovite from the tonalite-trondhjemite suite are 135 to 110 Ma (Pavlis, 1983; Winkler, 1989). A zircon age is 103 Ma (Winkler, unpub. data). Permian age fusulinids were found in a marble near the Eklutna complex (Clark, 1972a); the marble may be correlative with the metavolcanic and metasedimentary rocks intruded by the gabbroic rocks.	Cu - A few small magmatic copper prospects are known in the gabbroic rocks.
	Lined layered gabbroic complex: Chiefly gabbro, leucogabbro, thin layers of anorthosite, garnet-bearing layers locally. Includes rare pods of dunite, clinopyroxenite in Nelchina River Gabbro. Intruded by narrow clinopyroxenite and later websterite dikes in the Tonsina area.	2 km	No age data	
	Cumulate ultramafic complex: Eklutna: Thick sequence of partly serpentinized chromite-bearing dunite at base, with progressively more wehrlite and clinopyroxenite up section. Minor websterite. Gabbroites interlayered with clinopyroxenites near top. Red Mountain: Klippe composed of thick section of dunite, chromite, and subordinate clinopyroxenite. Local garnet clinopyroxenite. Klippe rests on McHugh Complex. Wolverine: Thin layers (fault slices?) of dunite, clinopyroxenite, wehrlite, altered gabbroite; extremely faulted, tectonized area (Burns and others, 1991).	0.5 km (?)	No age data	Cr - Wolverine complex - Chromite bands and stringers up to 8 cm thick occur in dunite and minor wehrlite. The west zone the complex grades 10 to 20% Cr. The east zone has a grade of 10 to 15% Cr (Foley and Barker, 1985; Newberry, 1986). Cr - Eklutna complex - An estimated 1000 t of Cr ₂ O ₃ in segregations of disseminated and banded chromite within serpentinized dunite (Foley and Barker, 1985). Cr - Red Mountain - Podiform chromite deposits in dunite and wehrlite. Reserve estimates include 1.6 million st of contained Cr ₂ O ₃ (Foley and Barker, 1985). Cr - Bernard Mountain - Three exposed deposits at Bernard Mountain contain identified resources equal to 343,000 tons of Cr ₂ O ₃ . Each deposit consists of numerous, thin lenses or bands of chromite up to several inches thick in dunite gangue. Chromite also occurs as disseminated grains ranging from 0.1 to 2 mm (Foley and Barker, 1985).



- EXPLANATION**
- Volcanoclastic sedimentary rocks, including mainly graywacke, sandstone, mudstone, and subordinate conglomerate
 - Hyaloclastite
 - Tuff, tuff breccia, lignimritic tuff
 - Andesite
 - Basalt
 - Trondhjemite-tonalite dikes and pluton
 - Quartz diorite, tonalite, minor granodiorite
 - Hornblende diabase dikes, boudinaged and partially assimilated by quartz diorite and gabbroic rocks
 - Plutonic dike complex composed of hornblende gabbro, pyroxene hornblende gabbro, magnetite gabbro, and quartz diorite
 - Metasomatized gabbroic rocks which contain abundant hornblende, chlorite, epidote, and magnetite
 - Inclusions of amphibolite, schist, and minor marble
 - Gabbroite and magnetite gabbroite - massive
 - Gabbroite, magnetite gabbroite, leucogabbroite, and anorthosite - foliated
 - Dunite, wehrlite, clinopyroxenite, and minor websterite
 - Veins of clinopyroxenite, orthopyroxenite and websterite
 - Harzburgitic dunite and harzburgite

- EXPLANATION**
- TKsu - Sedimentary rocks (Tertiary and Cretaceous)
 - Mzm - McHugh Complex - Extremely deformed greenstones and cherts (Mesozoic)
 - Jtv - Talkeetna Formation volcanic rocks (Jurassic)
 - Jts - Talkeetna Formation sedimentary rocks (Jurassic)
 - Jqd - Quartz diorite, tonalite (Jurassic)
 - Jdc - Plutonic dike complex composed of quartz diorite, hornblende gabbro, pyroxene hornblende gabbro, gabbroite, and diabase (Jurassic)
 - Jg - Gabbroite and magnetite gabbroite with subordinate pyroxene hornblende gabbroite and anorthosite (Jurassic)
 - Jum - Dunite, wehrlite, clinopyroxene, and minor websterite (Jurassic)
 - Jh - Harzburgitic dunite and harzburgite (Jurassic)

OPHIOLITIC COMPLEXES NEAR THE BORDER RANGES FAULT ZONE, SOUTHCENTRAL ALASKA

BY
LAUREL E. BURNS
1992

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.