

UNITED STATES DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY

U, Th, AND K ANALYSES OF SELECTED PLUTONIC ROCKS FROM  
WEST-CENTRAL ALASKA

By

Thomas P. Miller and Carl M. Bunker

Open-file report  
75-216  
1975

This report is preliminary  
and has not been edited or  
reviewed for conformity with  
Geological Survey standards

## Introduction

Numerous samples of plutonic rocks collected during reconnaissance mapping in western Alaska have been analyzed for K, U, and Th. The U and Th content of the plutonic rocks from the southeastern Seward Peninsula have been discussed in a separate report (Miller and Bunker, 1975); because of the current interest in U and Th, the analyses of the remaining samples are given in this report.

The analyzed samples (table 1) were selected from a group of samples originally collected for general petrologic studies. The samples are grab samples and are considered to be generally representative of the particular unit from which they were collected. References to pertinent reports that describe the geology and petrology of the sampled units are given in Table 1.

Analyses were by gamma-ray spectrometry, and the basic operational procedures and calibration techniques have been described by Bunker and Bush (1966, 1967). Uranium concentrations are determined indirectly by measuring the radium daughter products to obtain radium equivalent uranium (RaeU) values. Radium equivalent uranium is the amount of uranium, assuming radioactive equilibrium, required to support the amount of daughter products that emit the radioactivity measured in a sample. Although thorium is also measured from daughter products, disequilibrium is improbable because of short half-lives; therefore, the concentrations are considered to be a direct measurement of parent thorium. Potassium is determined from its  $K^{40}$  constituent, which is proportional to the total potassium. The coefficients of variation for the accuracy of the data included in this report are about 3 percent for

uranium and thorium and 1 percent for potassium when compared to standards analyzed by isotope dilution and flame photometer methods.

## References Cited

- Bunker, C. M., and Bush, C. A., 1966, Uranium, thorium, and radium analyses by gamma-ray spectrometry (0.184-0.352 million electron volts), in U. S. Geol. Survey Research 1966: U. S. Geol. Survey Prof. Paper 550-B, p. B176-B181.
- \_\_\_\_\_ 1967, A comparison on potassium analyses by gamma-ray spectrometry and other techniques: U. S. Geol. Survey Prof. Paper 575-B, p. B164-B169.
- Miller, T. P., 1971, Petrology of the plutonic rocks of west-central Alaska: U. S. Geol. Survey open-file report, 132 p.
- \_\_\_\_\_ 1972, Potassium-rich alkaline intrusive rocks of western Alaska: Geol. Soc. America Bull., v. 83, p. 2111-2128.
- Miller, T. P., and Bunker, C. M., 1975, A reconnaissance study of the U and Th content of plutonic rocks in the southeastern Seward Peninsula, Alaska: U. S. Geol. Survey open-file report #75-217, 33 p.
- Miller, T. P., and Ferrians, O. J., Jr., 1968, Suggested areas for prospecting in the central Koyukuk River region, Alaska: U. S. Geol. Survey Circ. 570, 12 p.
- Sainsbury, C. L., 1974, Geologic map of the Bendeleben quadrangle, Seward Peninsula, Alaska: The Mapmakers, Anchorage, Alaska, 31 p.

Table 1. Radioactivity parameters of selected plutonic rocks from west-central Alaska.

HAP NO.	FIELD NO.	LOCATION	U (ppm)	Th (ppm)	K (pct)	HEAT CONTENT (u cal/gm yr)	Th/U	U/Kx10 <sup>-4</sup>	Th/Kx10 <sup>-4</sup>	SAMPLE DESCRIPTION
1	68ANw324	65°41.9'N 162°35.7'W	7.28	36.21	2.93	13.35	4.97	2.48	12.36	Granodiorite; medium-grained, leucocratic, hornblende-biotite. Kagruk pluton, Bendeleben quadrangle (Sainsbury, 1974).
2	69ANw59	65°24.3'N 161°16.7'W	16.36	58.11	6.16	25.23	3.55	2.66	9.43	Pseudoleucite porphyry. Nepheline syenite zone, southwestern part of Granite Mountain pluton, Candle quadrangle, (Miller, 1971; 1972).
3	69ANw28	65°24.1'N 161°16.5'W	9.49	39.16	5.37	16.21	4.13	1.77	7.29	Foyaita. Nepheline syenite zone, southwestern part of Granite Mountain pluton, Candle quadrangle (Miller, 1971; 1972).
4	68ANw391A	65°27.2'N 161°15.4'W	5.29	10.84	5.46	7.50	2.05	0.97	1.99	Garnet syenite. Garnet syenite zone, western Granite Mountain pluton, Candle quadrangle (Miller, 1971; 1972).
5	69ANw24	65°25.8'N 161°15.4'W	5.50	24.30	4.69	10.14	4.42	1.17	5.18	Monzonite. Monzonite zone, central Granite Mountain pluton, Candle quadrangle (Miller, 1971; 1972).
6	69ANw30A	65°25.4'N 161°08.9'W	4.24	21.71	2.99	8.24	5.12	1.42	7.26	Quartz monzonite. Quartz monzonite zone, eastern Granite Mountain pluton (Miller, 1972).
7	67ANw113	66°06.8'N 160°47.3'W	9.68	32.97	4.46	14.86	3.41	2.17	7.39	Porphyritic quartz monzonite. Western Selawik Hills pluton, Selawik quadrangle (Miller, 1971).
8	66ANw146	66°12.1'N 160°39.3'W	6.77	42.98	4.38	14.72	6.35	1.55	9.81	Foyaita. Probable dike cutting monzonite-syenite zone of northwestern Selawik Hills pluton, Selawik quadrangle (Miller, 1971).
9	66ANw162	66°06.7'N 160°21.3'W	7.74	45.42	6.54	16.50	5.87	1.18	6.94	Monzonite; gneissic, hornblende-pyroxene, medium-grained. Monzonite-syenite zone, north-central Selawik Hills pluton, Selawik quadrangle (Miller, 1971).
10	66ANw163B	66°06.7'N 160°16.3'W	19.63	66.16	5.08	28.93	3.37	3.86	13.02	Monzonite; coarse-grained, trachytoid, hornblende-rich. Monzonite-syenite zone, north-central Selawik Hills quadrangle (Miller, 1971).
11	66APw117	66°09.7'N 159°49'W	3.05	21.56	4.82	7.84	7.07	0.63	4.47	Quartz monzonite, fine-grained, massive. Eastern Selawik Hills pluton, Selawik quadrangle (Miller, 1971).
12	67ANw410	66°02.2'N 159°50.3'W	2.49	9.65	3.20	4.61	3.88	0.78	3.02	Malignite. Hunt alkaline complex. Selawik quadrangle (Miller, 1971; 1972).
13	66ANw110E2	66°11.2'N 159°49.2'W	5.82	22.07	6.16	10.33	3.79	0.94	3.58	Malignite. Hunt alkaline complex. Selawik quadrangle (Miller, 1971; 1972).
14	66ANw120A2	66°19.9'N 160°23.7'W	10.87	20.67	13.83	15.80	1.90	0.79	1.49	Juvite. Selawik Lake alkaline complex. Selawik quadrangle (Miller, 1971; 1972).
15	67ANw360	66°18.5'N 159°43.5'W	31.20	177.83	5.04	59.70	5.70	6.19	35.28	Pulaskite; fluorite-bearing, possibly a dike. Inland Lake alkaline complex, Selawik quadrangle (Miller, 1971; 1972).
16	67ANw136	66°10.9'N 156°02.5'W	6.94	25.56	3.01	10.99	3.68	2.31	8.49	Granodiorite; medium-grained, biotite. Biotite granodiorite zone, south-central Zana Hills pluton, Shungnak quadrangle (Miller, 1971).
17	67ANw151	66°11.2'N 155°51.9'W	12.42	123.92	5.07	35.22	9.98	2.45	24.44	Monzonite; coarse-grained, porphyritic and trachytoid. Marginal monzonite-syenite unit, east-central Zana Hills pluton, Hughes quadrangle (Miller and Ferrans, 1968; Miller, 1971). Unit locally contains minor amounts of uraninite, thorite, and beciferite (K.H. Staatz, personal communication, 1975).

Analyses were done by C. M. Bunker and C. A. Bush.




Figure 1. Map showing location of analyzed samples. Shaded areas denote plutons.