UNITED STATES DEPARTMENT OF THE INTERIOR GEOLOGICAL SURVEY

HISTOGRAMS AND CORRELATION TABLE FOR
SELECTED ELEMENTS IN SAMPLES OF THE
ASH OF GROUND BIRCH LEAVES FROM THE
TANACROSS QUADRANGLE, ALASKA

by

G. C. Curtin, G. W. Day, and R. B. Tripp

Open-File Report 80-656

CONTENTS

		Page
Discussio	η	1
Reference	S	3
	FIGURES	
Figure 1.	Histogram showing iron, in percent, in samples of the ash	
	of ground birch leaves from the Tanacross quadrangle,	
	Alaska	4
2.	Histogram showing magnesium, in percent, in samples of the	
	ash of ground birch leaves from the Tanacross quadrangle,	
	Alaska	5
3.	Histogram showing titanium, in percent in samples of the ash	
	of ground birch leaves from the Tanacross quadrangle,	
	Alaska	6
4.	Histogram showing manganese, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	7
5.	Histogram showing silver, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	8
6.	Histogram showing gold, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	8
7.	Histogram showing boron, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	9

CONTENTS--Continued

FIGURES--Continued

		Page
Figure	8. Histogram showing barium, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	10
	9. Histogram showing cadmium, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	11
1	O. Histogram showing cobalt, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	12
1	1. Histogram showing chromium, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	12
1	2. Histogram showing copper, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	13
1	3. Histogram showing molybdenum, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	14
1	4. Histogram showing nickel, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	15
۱	5. Histogram showing lead, in parts per million (ppm), in	
	samples of the ash of ground birch leaves from the	
	Tanacross quadrangle, Alaska	36

CONTENTS--Continued

FIGURES--Continued

		•	Page
Figure	16.	Histogram showing strontium, in parts per million (ppm), in	
		samples of the ash of ground birch leaves from the	
		Tanacross quadrangle, Alaska	17
	17.	Histogram showing tin, in parts per million (ppm), in	
		samples of the ash of ground birch leaves from the	
		Tanacross quadrangle, Alaska	18
	18.	Histogram showing tungsten, in parts per million (ppm), in	
		samples of the ash of ground birch leaves from the	
		Tanacross quadrangle, Alaska	18
	19.	Histogram showing vanadium, in parts per million (ppm), in	
		samples of the ash of ground birch leaves from the	
		Tanacross quadrangle, Alaska	19
	20.	Histogram showing zinc, in parts per million (ppm), in	
		samples of the ash of ground birch leaves from the	
		Tanacross quadrangle, Alaska	20
	21.	Histogram showing zirconium, in parts per million (ppm), in	
		samples of the ash of ground birch leaves from the	
		Tanacross quadrangle, Alaska	23

CONTENTS--Continued

TABLE

	Page	•
Table 1.	Simple linear correlation coefficients between logarithmic	
	values of the element concentrations in the ash of 478	
	samples of ground birch leaves collected in the	
	Tanacross quadrangle, Alaska 22	,

DISCUSSION

These histograms and correlation table show the distribution and abundance of, and correlations between, the populations of 21 elements in the ash of ground birch (<u>Betula rotundifolia Sarg.</u>) leaves collected throughout most of the Tanacross 1:250,000-scale quadrangle. The samples were collected during reconnaissance geochemical studies as part of the U.S. Geological Survey's Alaskan Mineral Resource Assessment Program. Related data for ground birch are presented in other reports (Day, Curtin, and Tripp, 1979 a,b,c), and background information on the mineral resource assessment of the Tanacross quadrangle is presented in a U.S. Geological Survey Circular (Foster and others, 1976).

COLLECTION, PREPARATION, AND ANALYSIS OF SAMPLES

Samples of ground birch leaves were collected along streams near sites where stream sediments, heavy-mineral concentrates, and other samples were obtained.

Several branches from each of 2 or 3 bushes were sampled at each site to produce a composite sample. The present years' growth of leaves and twigs were pruned from the branches, placed in cloth bags and allowed to dry. The leaves were separated from the twigs and finely chopped in a blender. This composition was then ashed in a furnace at a peak temperature of approximately 500°C for 24 hours.

The ashed material was analyzed by an optical emission spectrographic method for plant materials (Mosier, 1972) for 29 elements including the 21 elements whose distributions are shown in the histograms. Elements not present in measurable amounts are: arsenic, beryllium, bismuth, lanthanum, niobium, antimony, scandium, and yttrium.

The results were entered into the computerized Rock Analysis Storage System (RASS) of the U.S. Geological Survey (VanTrump and Miesch, 1977) and data sets were analyzed by statistical programs in the U.S. Geological Survey's STATPAC system to produce the tabular statistics for the histograms and the correlation table. Arithmetic means, standard deviations, geometric means, and geometric deviations are shown for those element distributions in which there are less than 5 percent censored values. The correlation coefficients in the correlation table (table 1) that are in bold type are statistically significant at the 5 percent error level.

REFERENCES

- Day, G. W., Curtin, G. C., and Tripp, R. B., 1979a, Geochemical maps showing the distribution and abundance of lead, zinc, and cadmium in the ash of black spruce needles and ground birch leaves from the Tanacross quadrangle, Alaska: U.S. Geological Survey Open-file Report 79-1319.

- Foster, H. L., Albert, N. R. D., Barnes, D. R., Curtin, G. C., Griscom, Andrew, Singer, D. A., and Smith, J. G., 1976, The Alaskan Mineral Resource Assessment Program: Background information to accompany folio of geologic and mineral resource maps of the Tanacross quadrangle, Alaska: U.S. Geological Survey Circular 734, 23p.
- Mosier, E. L., 1972, A method for semiquantitative spectrographic analysis of plant ash for use in biogeochemical and environmental studies: Applied Spectroscopy, v. 26, no.6, p. 636-640.
- VanTrump, George, Jr., and Miesch, A. T., 1977, the U.S. Geological Survey

 RASS-STATPAC system for management and statistical reduction of geochemical

 data: Computers and Geosciences. v. 3, 0. 475-488.




Figure 1.--Histogram showing iron, in percent, in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 2.--Histogram showing magnesium, in percent, in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 3.--Histogram showing titanium, in percent in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 4.--Histogram showing manganese, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 5.--Histogram showing silver, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 6.--Histogram showing gold, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 7.--Ristogram showing boron, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 8.--Histogram showing barium, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 9.--Histogram showing cadmium, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 10.--Histogram showing cobalt, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.

د: د: ۲

Frequency Reported Reported frequency value 50 60 70 80 90 30 (mqq) 460 Not detected 4 < 5 5 12 1 7 0 10 1 15

Figure 11.--Histogram showing chromium, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 12.--Histogram showing copper, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 13.--Histogram showing molybdenum, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 14.--Histogram showing nickel, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 15.--Histogram showing lead, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 16.--Histogram showing strontium, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 17.--Histogram showing tin, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 18.--Histogram showing tungsten, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 19.--Histogram showing vanadium, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 20.--Histogram showing zinc, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.




Figure 21.--Histogram showing zirconium, in parts per million (ppm), in samples of the ash of ground birch leaves from the Tanacross quadrangle, Alaska.

Table 1.--Simple linear correlation coefficients between logarithmic values of the element concentrations in the ash of 478 samples of ground birch leaves collected in the Tanacross quadrangle, Alaska.

[Upper half of the table contains correlation coefficients, multiplied by 100; lower half is the number of pairs of values used to compute coefficients. ****indicates carrelation coefficient was not computed. Method of analysis indicated in the row and column headings: S * Emission spectrography. Boidface figures indicate significant positive correlation.]

	S-FEX	S-HCX	S-fix	- 1 2	Ω.	S-A:J	ζη.	a e	S-6	02-5	S-C4	s-C:J	1 1 10	==	5e-5	5-5	-		S-2N	æ	¥
		I			S-£3		S-5	S-94				~		111-5 6			Y-8 6	2-5	$\overline{}$	S-2R	<u>*</u>
S-2R	- 99	-99	-99	•	:	:	100	-99	100	•	•	-99	***	-99	-99	-99	-99	•	-99	. X	S-ZR
NZ - S	30	49	13	20	-12	-10	34	13	18	1	-58	50	4	1.1	38	32	1 14	81	×	2	NZ-S
 S	-27	24	-75	:	*	:	40	<u>.</u>	-54	*	++++	-30	0	-58	4	9	47	×	S	0	A-S
N-8	31	24	15	53	9-	-2	30	21	49	39	-70	11	0	7	21	-	×	5	478	2	S-v
S-SR	42	46	19	3	-	63	17	47	-10	-2	7.7	32	7	2.9	16	×	473	2	471	2	S-5R
S-PB	46	40	36	-	92-	21	18	2	13	19	-24	45	-:-	19	*	471	478	5	478	2	8-5
S-NI S	45	34	17	22	4-	63	7	40	ಬ	22	<u></u>	44	4	**	478	47:	478	S	478	2	S-N(
S-MD	===	4	7	-16	-1-	7	۲۷)	4	2	- 2	v	12	×	350	350	347	350	2	350		S-MO S
S-CU 3-	58	65	17	0	-23	-36	32	46	-		20	×	350	478	478	471	478	រា	478 1	2	S-CU S
S-CR S-	-2	4.1	30	- 96	001	:	82	57	-57	-45.	-	<u>*</u>	4	14	14	7	14	0	14	0	
	15		20	4	0	:		10	32	-	¥ ∞	217	173	217	217	215	217	4	217	_	0 S-CR
S-C0	8	0	_	23	 9	25 ••	P)	ស		¥ 9	4		-		_			4		2	S-C0
S-co							_	_	*	206	-	431	332	438	431	426	431		431	_	03-5
S-84	27	53	7	35	-13	49	13	<u>*</u>	431	217	14	478	350	478	478	471	478	-2	478	2	S-3A
S-8	19	82	^	28	-13	8	Ž	478	431	217	4	478	350	478	478	471	478	ري 	478	-2	5-B
S-AU	47	-17	74	73	65	Š	ß	5	Ŋ	2	٥	വ	ഹ	S.	IJ	ស	ß	0	v	٥	S-AU
S-AG	-27	-20	-12	011	X	5	59	60	58	33	2	59	46	59	59	59	59	0	59	-	S-AG
×-5	2	20	8-	<u>×</u>	21	2	186	186	150	56	8	186	94	186	186	183	186	2	186	-	S-MN
S-TIX	49	22	×	107	43	4	326	326	297	156	4	326	261	326	326	323	326	4	326	2	STIX
S-HGX	65	×	315	185	27	5	466	466	420	209	13	997	338	466	466	459	466	4	766	2	S-MCX S
S-FEX	×	464	326	135	59	ß	476	9/7	430	217	4	476	350	476	476	697	476	ß	476	7	S-FEX 18
*	3-EX	S-40x	\$-11 x	S-K.N	31-S	S-AU	S O	S-34	D0~8	%-C0	S-CR	S-C1	S-MO	. N. D	8d-8	S-5R	۸-5	X-S	N2-5	S-ZR	