UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

SUMMARIES OF DATA ON AND LISTS OF REFERENCES TO
METALLIC AND SELECTED NONMETALLIC MINERAL DEPOSITS
IN THE BIG DELTA AND TANACROSS QUADRANGLES, ALASKA

Compiled by
Edward H. Cobb
and
G. Donald Eberlein

**PROPERTY OF
LIBRARY
STATE OF ALASKA
DIVISION OF
GEOLOGICAL SURVEY**
Open-File Report 80-1086
1980

This report is preliminary
and has not been reviewed
for conformity with U.S.
Geological Survey editorial
standards.

Introduction

These summaries of data on metallic and selected nonmetallic mineral occurrences and lists of selected references to them in Geological Survey, U.S. Bureau of Mines, and State of Alaska Division of Geological and Geophysical Surveys (and predecessor State and Territorial agencies) reports and maps released before Jan. 1, 1980, are designed to aid in library research on the mineral resources of the Big Delta and Tanacross quadrangles, Alaska. The references are selected in the sense that mainly statistical reports such as the annual Minerals Yearbook of the U.S. Bureau of Mines and many annual and biennial reports of the Alaska Division of Geological and Geophysical Surveys and its predecessor agencies are not included. Also not included are data on many claims about which little more than their locations is known (for example, localities 16 to 22 in Menzie and Foster, 1979 (OF 78-529 D), p. 31). These omissions should not be interpreted as a judgement that the claims are not on valid mineral occurrences, but only that there are insufficient data to describe any mineral deposit that might be present. Geochemical anomalies determined by analysis of rock and stream-sediment samples in which no metallic mineral was identified are also omitted.

This report is divided into three parts: a section made up of summaries of data and reference lists arranged alphabetically by occurrence name for each quadrangle; a second section that lists, by quadrangle, synonyms for names in the first section, the names of owners and operators of mines and prospects, and claim names; and a final section that for each quadrangle lists alphabetically by author all references mentioned in the first section and in these introductory paragraphs.

The first section consists of data on each occurrence, for each of which there is a page that gives the name of the occurrence, the mineral commodities present (listed alphabetically); the mining district (Ransome and Kerns, 1954 (IC 7679)) in which the occurrence is located; the name of the 1:250,000-scale topographic quadrangle; co-

ordinates (as described by Cobb and Kachadoorian, 1961 (B 1139), p. 3-4); the number of a map (if there is one) on which the occurrence is shown and the number of the occurrence on that map; and the latitude and longitude of the occurrence. These data, presented at the top of the page, are followed by a short, general summary of the published data on the occurrence, with the source or sources of the information given in brackets. This is followed by a list, arranged chronologically, of references to the occurrence.

Proper names of mines, prospects, and other mineral occurrences are given if such names appear in the reports cited. If a deposit does not have such a name, but is near a named geographic feature, the name of that feature is shown in parentheses in lieu of a proper name if no ambiguity would result. All placer mines and prospects are considered under the name of the stream on which they are located. If a deposit has no proper name and is not near a named geographic feature, it is titled "Unnamed occurrence" and appears at the end of the list for the quadrangle. If a part of a proper name is not always used in references, that part of the name is shown in parentheses.

Citations are given in standard bibliographic format with the exception that references to reports and maps in numbered publication series also show in parentheses an abbreviation for the report or map series and the report or map number. Abbreviations used are:

B	U.S. Geological Survey Bulletin
BMB	U.S. Bureau of Mines Bulletin
C	U.S. Geological Survey Circular
GC	Alaska Division of Mines and Minerals Geochemical Report
GR	Alaska Division of Geological and Geophysical Surveys (and predecessor State agencies) Geologic Report
I	U.S. Geological Survey Miscellaneous Geologic Investigations Map
IC	U.S. Bureau of Mines Information Circular
OF	U.S. Geological Survey Open-file Report
MF	U.S. Geological Survey Miscellaneous Field Studies Map
P	U.S. Geological Survey Professional Paper
RI	U.S. Bureau of Mines Report of Investigations
TDM	Alaska Territorial Department of Mines Pamphlet
USBM OF	U.S. Bureau of Mines Open-file Report

Index map

(Banner Cr.)

Gold, Monazite, Tungsten

Fairbanks district
MF-388, loc. 13

Bog Delta (S.1, S.3)
64°18'N, 145°20'W

Summary: Bedrock is Cretaceous? rhyolite porphyry. Gold discovered in 1905 or shortly thereafter. Gold from this and neighboring creeks ranged from 639½ to 785 in fineness. Open-pit mining. Concentrate from churn-drill hole in bedrock near placer-mining area contained monazite and scheelite. Claim staking, prospecting, and possibly minor mining in area in 1970's. [Menzie and Foster, 1979 (OF 78-529 D); Saunders, 1965 (GC 3); Bundtzen and Reger, 1977 (GR 55)]

References

- Brooks, 1908 (B 345), p. 44
Brooks, 1909 (B 379), p. 55
Ellsworth, 1910 (B 442), p. 245
Ellsworth and Parker, 1911 (B 480), p. 167
Prindle and Katz, 1913 (B 525), p. 113, 141
Chapin, 1914 (B 592), p. 361
Smith, 1917 (BMB 142), p. 25
Brooks, 1918 (B 662), p. 60
Wedow and others, 1954 (C 331), p. 11
Saunders, 1965 (GC 3), p. 2, 4, 10
Cobb, 1972 (MF-388), loc. 13
Mulligan, 1974 (IC 8626), p. 14
Bundtzen and Reger, 1977 (GR 55), p. 29-30, 33
Cobb, 1977 (OF 77-168 B), p. 6, loc. 9
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 1
MacKevett and Holloway, 1977 (OF 77-169A), p. 12, loc. 4
Foster and others, 1979 (B 783), p. 13
Menzie and Foster, 1979 (OF 78-529 D), p. 27, loc. 1

(Beaver Cr.)

Gold, Tin

Fairbanks district
MF-388, loc. 8

Big Delta (4.6, 13.4)
64°47'N, 146°24'W

Summary: Ground averaging about \$1 per square foot of bedrock [gold at \$20.67] said to have been found in 1911-12 and several windlass dumps taken out. Scarce cassiterite in placer. [Ellsworth and Davenport, 1913 (B 542); Joesting, 1942 (TDM 1)]

References

- Ellsworth and Davenport, 1913 (B 542), p. 208
Joesting, 1942 (TDM 1), p. 34
Cobb, 1972 (MF-388), loc. 8
Cobb, 1977 (OF 77-168 B), p. 6, loc. 1
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 2
Menzie and Foster, 1979 (OF 78-529 D), p. 27, loc. 2

Blue Lead

Antimony, Copper, Gold, Silver

Goodpaster district
MF-388, loc. 5

Big Delta (21.2, 6.8)
64°22'N, 144°12'W

Summary: Gold-bearing quartz vein about 2½ ft thick in Cretaceous? granitic rocks near contact with Paleozoic and/or Precambrian? metamorphic rocks; faults and shearing common. Free gold in quartz containing sulfides; gold content decreases as sulfide content increases. Sulfides include stibnite. Sample of table concentrate from mill at Blue Lead contained pyrite, stibnite, covellite, and digenite. Jamesonite in wall rocks. Mine operated in 1938-39. Production from this and other mines (Gray Lead, Grizzly Bear) in area estimated at 132 oz gold and 25 oz silver from 150 tons of ore. More than 775 ft of underground workings at Blue Lead. [Thomas, 1970 (USBM OF 14-70); Menzie and Foster, 1979 (OF 78-529 D)]

References

- Smith, 1938 (B 897-A), p. 22-23
Smith, 1939 (B 917-A), p. 30
Berg and Cobb, 1967 (B 1246), p. 222
Thomas, 1970 (USBM OF 14-70), p. 7, 10-11
Cobb, 1972 (MF-388), loc. 5
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 3
Foster and others, 1979 (C 783), p. 4
Menzie and Foster, 1979 (OF 78-529 D), p. 28, loc. 4

(Boulder Cr.)

Molybdenum

Goodpaster district
MF-388, loc. 7

Big Delta (21.75, 6.8)
64°22'N, 144°08'W

Summary: Molybdenite occurs sparingly in small quartz veins in
Cretaceous? granitic rock. [Menzie and Foster, 1979 (OF
78-529 D)]

References

- Joesting, 1942 (TDM 1), p. 29
Berg and Cobb, 1967 (B 1246), p. 223
Cobb, 1972 (MF-388), loc. 7
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 4
Menzie and Foster, 1979 (OF 78-529 D), p. 29, loc. 5

(Buckeye Cr.)

Gold, Lead, Tin, Tungsten

Fairbanks district
MF-388, loc. 11

Big Delta (5.25, 5.5)
64°19'N, 145°19'W

Summary: Bedrock Cretaceous? rhyolite porphyry. Placer gold first mined in early 1900's. Concentrates contained gold, galena, scheelite, tourmaline, and cassiterite. [Menzie and Foster, 1979 (OF 78-529 D)] See also (Banner Cr.).

References

- Prindle and Katz, 1913 (B 525), p. 113, 142
Smith, 1917 (BMB 142), p. 25
Brooks, 1918 (B 662), p. 60
Smith, 1930 (B 813), p. 29
Smith, 1932 (B 824), p. 34
Wedow and others, 1954 (C 331), p. 11, 13
Saunders, 1965 (GC 3), p. 4
Koschmann and Bergendahl, 1968 (P 610), p. 31
Cobb, 1972 (MF-388), loc. 11
Mulligan, 1974 (IC 8626), p. 14
Bundtzen and Reger, 1977 (GR 55), p. 29
Cobb, 1977 (OF 77-168 B), p. 6, loc. 9
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 1
MacKevett and Holloway, 1977 (OF 77-169A), p. 12, loc. 3
Menzie and Foster, 1979 (OF 78-529 D), p. 27, loc. 1

(Butte Cr.)

Gold

Fairbanks district
MF-388, loc. 15

Big Delta (9.55, 12.9)
64°44'N, 145°43'W

Summary: Bedrock mostly garnetiferous quartz-mica schist and quartz-ite (Paleozoic?). Staked and some development work, 1905. About 20 ft of gravel overlain by 6 ft of muck. Considerable prospecting, but little production. [Menzie and Foster, 1979 (OF 78-529 D)]

References

- Prindle, 1906 (B 284), p. 123-125
Prindle, 1906 (B 295), p. 26
Brooks, 1908 (B 345), p. 43
Ellsworth, 1910 (B 442), p. 245
Ellsworth and Parker, 1911 (B 480), p. 167
Prindle, 1913 (B 538), p. 80
Cobb, 1972 (MF-388), loc. 15
Cobb, 1977 (OF 77-168 B), p. 6, loc. 3
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 5
Foster and others, 1979 (C 783), p. 4
Menzie and Foster, 1979 (OF 78-529 D), p. 29, loc. 6

(Caribou Cr.)

Bismuth, Gold, Tin, Tungsten

Fairbanks district
MF-388, loc. 16

Big Delta (9.6-10.2, 12.1-12.6)
64°41'-64°43'N, 145°38'-145°43'W

Summary: Bedrock mostly amphibolite-facies garnetiferous quartz-mica schist (Paleozoic?). First mining was in 1905. Dredge was installed in 1940's; worked most of the length of the creek. Scarce cassiterite and scheelite and small amounts of native bismuth in concentrates. [Menzie and Foster, 1979 (OF 78-529 D)]

References

- Prindle, 1906 (B 284), p. 123-125
Prindle, 1906 (B 295), p. 26
Brooks, 1908 (B 345), p. 43
Ellsworth, 1910 (B 442), p. 245
Ellsworth and Parker, 1911 (B 480), p. 167
Prindle, 1913 (B 538), p. 80
Brooks, 1916 (B 642), p. 67
Brooks, 1923 (B 739), p. 33
Smith, 1926 (B 783), p. 15
Smith, 1939 (B 910-A), p. 48
Smith, 1939 (B 917-A), p. 44, 46-47
Smith, 1941 (B 926-A), p. 40, 43
Joesting, 1942 (TDM 1), p. 34, 39
Cobb, 1972 (MF-388), loc. 16
Cobb, 1973 (B 1374), p. 129
Hasler and others, 1973 (P 820), p. 98
Cobb, 1977 (OF 77-168 B), p. 6, loc. 4
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 6
Foster and others, 1979 (C 783), p. 4, 13
Menzie and Foster, 1979 (OF 78-529 D), p. 29, loc. 8

(Central Cr.)

Gold

Goodpaster district
OF 78-529 D, loc. 9

Big Delta (19.4, 6.4) approx.
63°20'N, 144°27'W approx.

Summary: Placer ground being developed in 1927. Placer claims staked, 1939. Lode and placer claims staked, 1974-78. [Smith, 1930 (B 810); Menzie and Foster, 1979 (OF 78-529 D)]

References

Smith, 1930 (B 810), p. 26

Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 7

Menzie and Foster, 1979 (OF 78-529 D), p. 29, loc. 9

Democrat

Gold

Fairbanks district

Big Delta (4.9, 5.9)

MF-388, loc. 1

64°20'N, 146°21'W

Summary: Adit of unknown length driven prior to 1921; gold in rhyolite; no known production, though a mill was built on the property. Prospecting and staking in area in the 1960's and 1975-78. [Menzie and Foster, 1979 (OF 78-529 D)] Includes references to lode on Democrat Cr. See also (Democrat Cr.).

References

Martin, 1919 (B 692), p. 39

Brooks, 1923 (B 739), p. 33

Saunders, 1965 (GC 3), p. 2

Berg and Cobb, 1967 (B 1246), p. 218

Cobb, 1972 (MF-388), loc. 1

Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 8

MacKevett and Holloway, 1977 (OF 77-169A), p. 12, loc. 1

Menzie and Foster, 1979 (OF 78-529 D), p. 30, loc. 11

(Democrat Cr.) (Gulch) (Pup) Gold, Silver

Fairbanks district Big Delta (4.9, 5.9)
MF-388, loc. 10 64°20'N, 145°21'W

Summary: Deposit at least in part a residual placer on Cretaceous? rhyolite porphyry, a random sample of which contained 0.24 oz silver per ton and a trace of gold. Has been placer mining. [Saunders, 1965 (GC 3); Menzie and Foster, 1979 (OF 78-529 D)] Report of scheelite in pan concentrate [Bundtzen and Reger, 1977 (GR 55), p. 30] evidently is in error; the cited source of data [Saunders, 1965 (GC 3)] does not mention scheelite in description of sample from Democrat Cr. See also (Banner Cr.), Democrat.

References

- Brooks, 1909 (B 379), p. 55
Ellsworth, 1910 (B 442), p. 245
Ellsworth and Parker, 1911 (B 480), p. 167
Ellsworth and Davenport, 1913 (B 542), p. 222
Prindle and Katz, 1913 (B 525), p. 113, 142
Chapin, 1914 (B 592), p. 361
Eakin, 1915 (B 622), p. 235
Smith, 1917 (BMB 142), p. 25
Brooks, 1923 (B 739), p. 33
Smith, 1926 (B 783), p. 15
Smith, 1929 (B 797), p. 20
Smith, 1930 (B 810), p. 26
Smith, 1930 (B 813), p. 29
Smith, 1932 (B 824), p. 34
Smith, 1933 (B 836), p. 34
Saunders, 1965 (GC 3), p. 2, 4
Koschmann and Bergendahl, 1968 (P 610), p. 31
Cobb, 1972 (MF-388), loc. 10
Mulligan, 1974 (IC 8626), p. 14
Bundtzen and Reger, 1977 (GR 55), p. 30, 33
Cobb, 1977 (OF 77-168 B), p. 6, loc. 8
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 1
MacKevett and Holloway, 1977 (OF 77-169A), p. 12, loc. 2
Menzie and Foster, 1979 (OF 78-529 D), p. 17, loc. 1

(Gold Run Cr.)

Gold

Fairbanks district
OF 77-168 B, loc. 7

Big Delta (5.0, 7.25)
64°25'N, 146°21'W

Summary: Gold prospects were found, but there was no mining.
[Mulligan, 1974 (IC 8626)]

References

Mulligan, 1974 (IC 8626), p. 14
Cobb, 1977 (OF 77-168 B), p. 6, loc. 7

(Granite Cr.)

Gold(?)

Goodpaster district

Big Delta (15.5, 4.25) approx.

64°13'N, 145°00'W approx.

Summary: Tributary of South Fork of Goodpaster R. [not shown on modern maps]. In 1916 very coarse gold reported to have been found in gravels said to be 12-25 ft thick. [Brooks, 1918 (B 662)] Description same as that for Michigan Cr.

References

Brooks, 1918 (B 662), p. 60

Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 11

Menzie and Foster, 1979 (OF 78-529 D), p. 31, loc. 23

Gray (Quartz) Lead

Gold

Goodpaster district

Big Delta (20.8, 6.5)

MF-388, loc. 3

64°21'N, 144°15'W

Summary: Steeply dipping vein about 2 ft thick along contact between Cretaceous? granitic rocks and Paleozoic and/or Precambrian? gneissic schist; intense faulting and shearing along contact. Deposit explored and worked, 1939-41; about 1,400 ft of drifts, crosscuts, and raises. Gold fine and generally visible only after ore is crushed and panned. Minor exploration and attempt at reworking tailings in 1970's. [Menzie and Foster, 1979 (OF 78-529 D); Thomas, 1970 (USBM OF 14-70)] See also Blue Lead.

References

Thomas, 1970 (USBM OF 14-70), p. 7-9, 11

Cobb, 1972 (MF-388), loc. 3

Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 9

Foster and others, 1979 (C 783), p. 4

Menzie and Foster, 1979 (OF 78-529 D), p. 30, loc. 14

Grizzly Bear

Gold

Goodpaster district
MF-388, loc. 4

Big Delta (21.15, 6.6)
64°21'N, 144°12'W

Summary: Gold mined from steeply dipping quartz vein $1\frac{1}{2}$ to $2\frac{1}{2}$ in wide. Gold near hanging wall and coarser than in other veins is area. More than 300 ft of underground workings. Amount of ore mined small. [Eberlein and others, 1977 (OF 77-168 D)] See also Blue Lead.

References

- Smith, 1939 (B 917-A), p. 30
Berg and Cobb, 1967 (B 1246), p. 222
Thomas, 1970 (USBM OF 14-70), p. 11
Cobb, 1972 (MF-388). loc. 4
Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 10
Foster and others, 1979 (C 783), p. 4
Menzie and Foster, 1979 (OF 78-529 D), p. 30, loc. 15

(Hinkley Gulch)

Gold, Tin

Fairbanks district
MF-388, loc. 12

Big Delta (5-25, 5.4)
64°18'N, 145°19'W

Summary: Bedrock Cretaceous? rhyolite porphyry; deposit at least in part a residual placer. Cassiterite as well as gold in concentrates. Mining as recently as 1950. [Menzie and Foster, 1979 (OF 78-529 D); Saunders, 1965 (GC 3)] See also: (Banner Cr.), (Buckeye Cr.).

References

- Smith, 1933 (B 836), p. 34
Saunders, 1965 (GC 3), p. 2, 4
Cobb, 1972 (MF-388), loc. 12
Bundtzen and Reger, 1977 (GR 55), p. 29-30, 33
Cobb, 1977 (OF 77-168 B), p. 6, loc. 9
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 1
MacKevett and Holloway, 1977 (OF 77-169A), p. 12, loc. 3
Menzie and Foster, 1979 (OF 78-529 D), p. 27, loc. 1

(Kenyon Cr.)

Gold

Goodpaster district

Big Delta(?)

SW $\frac{1}{4}$ SE $\frac{1}{4}$ SE $\frac{1}{4}$ quad.(?)

Summary: Tributary of Healy Lake; may be in Mt. Hayes quad. Gold and garnet in gravels about 125 ft. deep. Discovered in 1914; a little mining reported in 1915. [Cobb, 1979, (OF 79-238)]

References

Brooks, 1915 (B 622), p. 50-51

Brooks, 1916 (B 642), p. 67

Brooks, 1918 (B 662), p. 60

Cobb, 1973 (B 1374), p. 137

Cobb, 1979 (OF 79-238), p. 41

(Michigan Cr.)

Gold

Goodpaster district
MF-388, loc. 18

Big Delta (18.4, 3.05) approx.
64°09'N, 144°35'W approx.

Summary: Gold discovered in 1915, setting off a short-lived stampede. Gold coarse (one \$9 nugget [gold at \$20.67] found). Gravel reported to be 12-25 ft thick. Deposits low grade. [Thomas, 1970 (USBM OF 14-70); Brooks, 1918 (B 662)]

References

- Brooks, 1918 (B 662), p. 60
Smith, 1930 (B 810), p. 26
Thomas, 1970 (USBM OF 14-70), p. 7
Cobb, 1972 (MF-388), loc. 18
Cobb, 1974 (B 1374), p. 137
Cobb, 1977 (OF 77-168 B), p. 6, loc. 11
Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 11
Menzie and Foster, 1979 (OF 78-529 D), p. 31, loc. 23

(Mineral Cr.)

Gold(?)

Fairbanks district

Big Delta
N $\frac{1}{2}$ SW $\frac{1}{4}$ quad.

Summary: Indications [of placer gold] found in 1922; sufficiently favorable to warrant further investigation. [Brooks and Capps, 1924 (B 755)] Only location is Richardson district; may be tributary of Banner Cr. or Buckeye Cr.

Reference

Brooks and Capps, 1924 (B 755), p. 38

(Minton Cr.)

Tungsten

Fairbanks district

Big Delta (5.75, 6.7)

64°23'N, 146°15'W

Summary: Pan concentrate from geochemical sample contained a piece of scheelite. Other heavy minerals included magnetite, ilmenite, and zircon. [Saunders, 1965 (GC 3)]

Reference

Saunders, 1965 (GC 3), p. 10, sample 109

(Moore Cr.)

Gold

Fairbanks district
MF-388, loc. 11

Big Delta (5.5, 5.5) approx.
64°20'N, 146°17'W approx.

Summary: Small tributary of Buckeye Cr. on which there has been placer gold mining. [Menzie and Foster, 1979 (OF 78-168 D)] See also: (Banner Cr.), (Buckeye Cr.).

References

- Smith, 1930 (B 810), p. 26
Cobb, 1972 (MF-388), loc. 11
Cobb, 1977 (OF 77-168 B), p. 6, loc. 9
Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 1
MacKevett and Holloway, 1977 (OF 77-169A), p. 12, loc. 3
Menzie and Foster, 1979 (OF 78-529 D), p. 27, loc. 1

(Nickel Cr.)

Chromite, Nickel

Fairbanks district

Big Delta (15.55, 13.4) approx.

MF-388, loc. 2

64°45'N, 144°55'W approx.

Summary: Silica-carbonate zone adjacent to serpentized peridotite contains traces of nickel. Peridotite in this region is reported to have small areas with segregations of high-grade chromite 1 ft thick and 3 ft or so long, but mostly the chromite content is less than 1 percent. [Eberlein and others, 1977 (OF 77-168 D); Joesting, 1942 (TDM 1)]

References

Joesting, 1942 (TDM 1). p. 16, 18

Berg and Cobb, 1967 (B 1246), p. 221

Cobb, 1972 (MF-388), loc. 2

Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 12

Menzie and Foster, 1979 (OF 78-529 D), p. 32, loc. 26

(No Grub Cr.)

Bismuth, Gold, Tungsten

Fairbanks district
MF-388, loc. 17

Big Delta (10.4, 11.9)
64°40'N, 145°37'W

Summary: Bedrock mostly quartz-mica schist of Paleozoic age. Intermittent mining from early 1900's to as recently as 1977. Considerable native bismuth found with gold and subordinate scheelite. Gold reported to be coarse and irregular and derived from gold- and bismuth-bearing quartz veins. [Menzie and Foster, 1979 (OF 78-529 D)]

References

- Ellsworth and Parker, 1911 (B 480), p. 167
Prindle, 1913 (B 538), p. 80
Brooks, 1916 (B 642), p. 67
Smith, 1942 (B 933-A), p. 39
Cobb, 1972 (MF-388), loc. 17
Hasler and others, 1973 (P 820), p. 98
Cobb, 1977 (OF 77-168 B), p. 6, loc. 5
Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 13
Foster and others, 1979 (C 783), p. 5
Menzie and Foster, 1979 (OF 78-529 D), p. 32, loc. 27

(Pasco Cr.)

Gold

Fairbanks district
OF 78-529 D, loc. 29

Big Delta (11.0, 12.0) approx.
64°41'N, 145°32'W approx.

Summary: Signs of small placer-mining activity; first record of claims, 1954; some claims active, 1976-77. [Eberlein and others, 1977 (OF 77-168 D); Menzie and Foster, 1979 (OF 78-529 D)]

References

Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 14
Menzie and Foster, 1979 (OF 78-529 D), p. 32, loc. 29

(Pine Cr.)

Gold, Tin

Fairbanks district
MF-388, loc. 9

Big Delta (5.1, 13.0)
64°44'N, 146°19'W

Summary: Scarce placer cassiterite reported; presence of gold assumed. Claims for lode gold have been staked in area.
[Menzie and Foster, 1979 (OF 78-529 D)] Includes reference to (Pyne Cr.).

References

Joesting, 1942 (TDM 1), p. 34
Cobb, 1972 (MF-388), loc. 9
Cobb, 1977 (OF 77-168 B), p. 6, loc. 2
Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 15
Menzie and Foster, 1979 (OF 78-529 D), p. 32, loc. 31

(Redemption Cr.)

Gold(?)

Fairbanks distroct(?)

Big Delta(?)

SW $\frac{1}{4}$ quad.(?)

Summary: Placer gold prospecting reported, 1926. [Smith, 1929
(B 797)] Creek may be in Mt. Hayes quad. and may be in
either Delta River or Goodpaster district.

Reference

Smith, 1929 (B 797), p. 20

(Redmond Cr.)

Gold

Fairbanks district Big Delta (3.0-3.75, 6.6-7.8) approx.
OF 78-529 D, loc. 33 (in part) 64°23'-64°27'N, 146°30'-146°36'W approx.

Summary: Prospecting in 1908-09. Depth to bedrock at mouth of Mosquito Cr. is said to be 40-50 ft. Has been mining on headwater tributaries. Gold found during prospect drilling, but not enough to stimulate mining. [Ellsworth, 1910 (B 442); Mulligan, 1974 (IC 8626); Wedow and others, 1954 (C 331)]

References

- Ellsworth, 1910 (B 442), p. 245
Wedow and others, 1954 (C 331), p. 11
Mulligan, 1974 (IC 8626), p. 14
Cobb, 1977 (OF 77-168 B), p. 6, loc. 6
Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 16
Menzie and Foster, 1979 (OF 78-529 D), p. 33, loc. 33

(Salcha R.)

Gold

Fairbanks district

Big Delta

OF 78-529 D, loc. 17 (in part)

N $\frac{1}{2}$ quad.

Summary: Prospecting above and claim staking at mouth of No Grub Cr. [Ellsworth and Parker, 1911 (B 480); Menzie and Foster, 1979 (OF 78-529 D)] Mining reported in references listed below as in the Salcha R. drainage was probably mainly on Butte, Caribou, and No Grub Creeks. See also: (Butte Cr.), (Caribou Cr.), (No Grub Cr.).

References

Brooks, 1906 (B 284), p. 7

Ellsworth and Parker, 1911 (B 480), p. 167

Smith, 1933 (B 844-A), p. 34

Smith, 1934 (B 857-A), p. 31

Smith, 1934 (B 864-A), p. 36

Smith, 1936 (B 868-A), p. 37

Smith, 1937 (B 880-A), p. 41

Menzie and Foster, 1979 (OF 78-529 D), p. 32, loc. 27

(Savage Gulch)

Gold(?)

Fairbanks district

Big Delta
N $\frac{1}{2}$ SW $\frac{1}{4}$ quad.

Summary: Indications [of placer gold] found in 1912; sufficiently favorable to warrant further investigation. [Brooks and Capps, 1924 (B 755)] Only location is Richardson district; may be tributary of Banner Cr. or Buckeye Cr.

Reference

Brooks and Capps, 1924 (B 755), p. 38

(Tenderfoot Cr.)

Gold, Lead, Silver

Fairbanks district

Big Delta (5.7-6.1, 4.7-5.1)

MF-388, loc. 14

64°16'-64°17'N, 145°12'-145°15'W

Summary: Area underlain by metasedimentary and metaigneous rocks of "Birch Creek Schist" with common small quartz veins and stringers. Gold (low grade with much alloyed silver) was discovered in 1905. Alluvium and loess near head of creek 30 ft thick; more than 155 ft thick near mouth. Creek was worked mainly by drifting, during which gold-bearing galena float was found. Stream was major producing creek in old Richardson gold district, the total production of which was about 95,000 oz of gold and 24,000 oz of silver before 1930 and an unknown amount since then. As of 1978 there has been recent prospecting, claim staking, and possibly a little mining. [Menzie and Foster, 1979 (OF 78-529 D); Saunders, 1965 (GC 3)]

References

- Prindle, 1906 (B 284), p. 123-124
Brooks, 1907 (B 314), p. 37
Brooks, 1908 (B 345), p. 44
Prindle, 1908 (B 337), p. 29
Brooks, 1909 (B 379), p. 55
Ellsworth, 1910 (B 442), p. 245
Ellsworth and Parker, 1911 (B 480), p. 167
Ellsworth and Davenport, 1913 (B 542), p. 222
Prindle and Katz, 1913 (B 525), p. 113, 141
Chapin, 1914 (B 592), p. 361
Eakin, 1915 (B 622), p. 235
Brooks, 1916 (B 642), p. 67
Smith, 1917 (BMB 142), p. 25
Brooks, 1918 (B 662), p. 60
Brooks, 1923 (B 739), p. 33
Smith, 1929 (B 797), p. 20
Smith, 1930 (B 810), p. 26
Smith, 1930 (B 813), p. 29
Smith, 1932 (B 824), p. 34
Smith, 1933 (B 836), p. 34
Wedow and others, 1954 (C 331), p. 11
Saunders, 1965 (GC 3), p. 2, 4
Koschmann and Bergendahl, 1968 (P 610), p. 31
Cobb, 1972 (MF-388), loc. 14
Mulligan, 1974 (IC 8626), p. 14
Bundtzen and Reger, 1977 (GR 55), p. 29, 33
Cobb, 1977 (OF 77-168 B), p. 6, loc. 10
Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 17

(Tenderfoot Cr.) -- cont.

MacKevett and Holloway, 1977 (OF 77-169A), p. 12, loc. 5

Eberlein and Menzie, 1978 (OF 78-1-D), p. 31

Foster and others, 1979 (C 783), p. 4, 13

Menzie and Foster, 1979 (OF 78-529 D), p. 33, loc. 36

(Tibbs Cr.)

Antimony, Gold, Molybdenum

Goospaster district
MF-388, locs. 3, 5, 6, 19

Big Delta (20.75-21.25, 6.5-7.1)
64°21'-64°23'N, 144°11'-144°15'W

Summary: Has been limited gold placer mining [MF-388, loc. 19]. Quartz veins in granitic rocks carry gold, jamesonite, stibnite, and molybdenite. For description of a typical occurrence see Blue Lead sheet. [Menzie and Foster, 1979 (OF 78-529 D); Cobb, 1973 (B 1374); Joesting, 1942 (TDM 1); Thomas, 1970 (USBM OF 14-70)] Includes references to: Hidden Treasure, Michigan Lead. See also: Blue Lead, Gray Quartz Lead, Grizzly Bear.

References

- Smith, 1941 (B 926-A), p. 28
Joesting, 1942 (TDM 1), p. 14, 29
Berg and Cobb, 1967 (B 1246), p. 222
Thomas, 1970 (USBM OF 14-70), p. 11
Cobb, 1972 (MF-388), locs. 3, 5, 6, 19
Cobb, 1973 (B 1374), p. 136-137
Cobb, 1977 (OF 77-168 B), p. 6, loc. 12
Eberlein and others, 1977 (OF 77-168 D), p. 13, loc. 18
Eberlein and Menzie, 1978 (OF 78-1-D), p. 35
Foster and others, 1979 (C 783), p. 4-5
Menzie and Foster, 1979 (OF 78-529 D), p. 34, loc. 37; p. 38, locs. 43, 46

(Twentymile Cr.)

Gold(?)

Fairbanks district

Big Delta

N $\frac{1}{2}$ quad.

Summary: Prospecting in 1910; tributary of Salcha R. [Ellsworth and Parker, 1911 (B 480)] Creek not shown on modern maps; may be the stream now called McCoy Cr.

References

Ellsworth and Parker, 1911 (B 480), p. 167

Eberlein and others, 1977 (OF 77-168 D), p. 12, loc. 5

Menzie and Foster, 1979 (OF 78-529 D), p. 29, loc. 6

(Wheeler Cr.)

Copper

Fairbanks district

Big Delta (7.4, 14.65)

64°50'N, 146°01'W

Summary: Quartz vein contains chalcocite, bornite, azurite, and malachite. [Menzie and Foster, 1979 (OF 78-529 D)]

Reference

Menzie and Foster, 1979 (OF 78-529 D), p. 46, loc. 182

Unnamed occurrence

Gold(?)

Fairbanks district

Big Delta (12.05, 12.1)
64°41'N, 145°24'W

Summary: Black iridescent rock cuts garnetiferous amphibole gneiss; appears to contain gold specks. [Menzie and Foster, 1979 (OF 78-529 D)]

Reference

Menzie and Foster, 1979 (OF 78-529 D), p. 40, loc. 90

Unnamed occurrence

Lead

Fairbanks district

Big Delta (19.2, 10.75)

Goodpaster district

64°36'N, 144°27'W

Summary: Galena in quartzite and quartz-mica schist. [Menzie and Foster, 1979 (OF 78-529 D)] Occurrence on boundary between Fairbanks and Goodpaster districts.

Reference

Menzie and Foster, 1979 (OF 78-529 D), p. 38, loc. 61

Unnamed occurrence

Lead, Zinc(?)

Fairbanks district

Big Delta (10.25, 15.6)

64°53'N, 145°37'W

Summary: Galena and sphalerite(?) in secondary silica layer in
impure limestone. [Menzie and Foster, 1979 (OF 78-529 D)]

Reference

Menzie and Foster, 1979 (OF 78-529 D), p. 44, loc. 153

Unnamed occurrence

Molybdenum

Goodpaster district

Big Delta (21.15, 5.6)

64°18'N, 144°13'W

Summary: Pyrite and molybdenite in quartz in a mineralized area in granitic rocks. Euhedral quartz crystals in vugs. [Menzie and Foster, 1979 (OF 78-529 D)]

Reference

Menzie and Foster, 1979 (OF 78-529 D), p. 36, loc. 21

ASARCO

Copper, Molybdenum

Fortymile district
MF-767E, loc. 20

Tanacross (11.8, 6.6)
63°22'N, 142°30'W

Summary: Silicified and deeply leached Tertiary quartz porphyry hypabyssal intrusive and, in part, extrusive rocks contain disseminated molybdenite and copper sulfides. Company sampling and drilling program in 1970. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 20

Eberlein and others, 1977 (OF 77-168D), p. 101, loc. 1

B.C.

Gold

Fortymile district
MF-767E, loc. 14

Tanacross (23.7, 1.85)
63°15'N, 141°01'W

Summary: Eluvial and gold-bearing bench gravels about 20 ft thick overlie decomposed Mesozoic granodiorite that contains gold quartz veins. The eluvial gold has delicate wire and crystal forms. Other creeks that are tributary to McArthur Cr. may have placer gold deposits; has been recent claim-staking activity. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 14
Eberlein and others, 1977 (OF 77-168D), p. 101, loc. 2
MacKevett and Holloway, 1977 (OF 77-169A), p. 75, loc. 6

(Big Cr.)

Lead, Zinc

Fortymile district
MF-767E, loc. 12

Tanacross (22.15, 9.45)
63°31'N, 141°10'W

Summary: Paleozoic and/or Precambrian? quartz-mica schist contains disseminated pyrite and lead and zinc sulfides. In an area of possible stratiform sulfide deposits. Claim-staking activity, 1975-76. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 12
Eberlein and others, 1977 (OF 77-168D), p. 101, loc. 3

Bluff

Copper, Molybdenum

Fortymile district
MF-767E, loc. 9

Tanacross (19.6, 11.45)
63°38'N, 141°29'W

Summary: Disseminated pyrite, chalcopyrite, molybdenite, and magnetite in porphyritic quartz monzonite, granodiorite, and quartz porphyry high-level intrusives of mid-Cretaceous—early Tertiary age; intense hydrothermal alteration. Numerous faults and dikes. Has been explored by surface sampling, diamond drilling, and geophysical methods. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 9
Eberlein and others, 1977 (OF 77-168D), p. 101, loc. 4

(Boulder Cr.)

Antimony, Gold, Silver

Tok district
MF-767E, loc. 22

Tanacross (1.6, 4.35)
63°15'N, 143°48'W

Summary: Quartz-stibnite vein between 2 faults that converge upward; at creek level is nearly 20 ft across. Central 6-ft thickness is high-grade stibnite (chiefly bladed crystals) with a little quartz; border zones have less stibnite and more quartz and a few grains of pyrite. Ore carries about \$1.00 in gold [at \$35] per ton and traces of silver and arsenic. Country rock is Paleozoic or Precambrian schist; much folded, faulted, and cut by granitic rocks. Some ore was mined, but not shipped in 1940-41. Recent production reported. [Moffit, 1954 (B 989-D); Joesting, 1942 (TDM 1); MacKevett and Holloway, 1977 (OF 77-166A)] Includes references to: A Lucky Leak, Stibnite, (Stibnite Cr.), antimony on Tok R.

References

- Brooks, 1916 (B 649), p. 63
Moffit, 1938 (B 904), p. 43-45
Joesting, 1942 (TDM 1), p. 13
Smith, 1942 (B 933-A), p. 89
Moffit, 1944 (B 943-B), p. 44-45
Ebbley and Wright, 1948 (RI 4173), p. 4, 30-35
Moffit, 1954 (B 989-D), p. 190, 201, 207-208
Berg and Cobb, 1967 (B 1246), p. 238
Foster, 1970 (I-593)
Cobb, 1972 (MF-383)
Foster and others, 1976 (C 734), p. 2
Singer and others, 1976 (MF-767E), loc. 22
MacKevett and Holloway, 1977 (OF 77-169A), p. 75, loc. 1

Fairplay

Copper, Lead, Molybdenum, Zinc

Fortymile district

Tanacross (13.75, 12.2)

MF-767E, loc. 5

63°41'N, 142°12'W

Summary: Disseminated copper, molybdenum, lead, and zinc sulfides in hydrothermally altered Tertiary hypabyssal intrusive and associated felsic volcanic rocks. Claims staked in early 1970's. Other similar prospects in neighborhood. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 5

Eberlein and others, 1977 (OF 77-168D), p. 101, loc. 6

Fishhook

Silver

Fortymile district
OF 77-168D, loc. 7

Tanacross (20.3, 14.9) approx.
63°50'N, 141°23'W approx.

Summary: Prospect located on basis of geochemical anomalies. Selected grab samples containsulfosalts and assay significant silver. Bedrock is augen gneiss (Paleozoic and/or Precambrian in age). [Eberlein and others, 1977 (OF 77-168D)]

Reference

Eberlein and others, 1977 (OF 77-168D), p. 101, loc. 7

(Fortymile R., Dennison Fork) Gold(?)
Fortymile district Tanacross
 SW $\frac{1}{4}$ SE $\frac{1}{4}$ quad.

Summary: Man who prospected somewhere in upper drainage of the
Dennison Fork in 1912 reported "very generally distributed"
fine gold with some colors from 10¢ to 30¢ in value [gold
at \$20.67]. [Ellsworth and Davenport, 1913 (B 542)]

Reference

Ellsworth and Davenport, 1913 (B 542), p. 215

Frey

Building stone

Chistochina district
MF-767E, loc. 24

Tanacross (0.4, 0.15)
63°01'N, 143°57'W

Summary: Marble consisting of thin laminae, lenses, and veins of magnesite, dolomite, and quartz in shades of green, cream, and white makes up a band 3 to 9 m thick and probably 1.6 km long. In terrane of Paleozoic and/or Precambrian metamorphic rocks. Stone has been quarried for ornamental use. [Singer and others, 1976 (MF-767E); Foster and others, 1976 (C 734)] Name used only in Richter, 1967 (GR 30).

References

Richter, 1967 (GR 30), p. 19

Foster, 1970 (I-593)

Foster and others, 1976 (C 734), p. 2

Singer and others, 1976 (MF-767E), loc. 24

Mosquito

Copper, Molybdenum

Fortymile district

Tanacross (4.1, 15.55)

MF-767E, loc. 1

63°53'N, 143°28'W

Summary: Disseminated chalcopyrite, molybdenite, and pyrite in hydrothermally altered Upper Cretaceous-lower Tertiary quartz monzonite and quartz latite porphyry. Were company drilling and sampling programs, 1974-75. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 1

Eberlein and others, 1977 (OF 77-168D), p. 102, loc. 9

Move

Gold(?)

Fortymile district
MF-767E, loc. 15

Tanacross (23.75, 1.1)
63°02'N, 141°01'W

Summary: Gold-bearing quartz veins in Mesozoic or Tertiary granitic rocks; other prospects nearby. [MacKevett and Holloway, 1977 (OF 77-169A)]. Claim staking, 1976 and probably more recently. No information on nature and extent of mineralization. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 15
Eberlein and others, 1977 (OF 77-168D), p. 102, loc. 10
MacKevett and Holloway, 1977 (OF 77-169A), p. 75, loc. 7

Noah

Copper(?)

Tok district

Tanacross (4.65, 2.85)

MF-767E, loc. 21

63°10'N, 143°35'W

Summary: Copper prospect in schist near a Mesozoic or Tertiary pluton.
[MacKevett and Holloway, 1977 (OF 77-169A)]

References

Singer and others, 1976 (MF-767E), loc. 21

MacKevett and Holloway, 1977 (OF 77-169A), p. 75, loc. 2

Paternie

Copper, Molybdenum, Zinc

Fortymile district
MF-767E, loc. 2

Tanacross (9.65, 10.8)
63°37'N, 142°46'W

Summary: Disseminated sulfide minerals occur in altered quartz monzonite and latite porphyry (Late Cretaceous-early Tertiary in age). Bedrock samples contain values of molybdenum and zinc. Company drilling and sampling program, 1974; exploration activity, 1971. Resources column of table lists Mo, Cu, Zn. [Eberlein and others, 1977 (OF 77-168D)] Copper mineral(s) and molybdenite probably present; zinc mineral may have been recognized or Zn may have been found only in analyses. Includes reference to Paternie.

References

Singer and others, 1976 (MF-767E), loc. 2
Eberlein and others, 1977 (OF 77-168D), p. 103, loc. 11

(Pitka Canyon)

Copper, Zinc

Fortymile district
MF-767E, loc. 7

Tanacross (20.1, 14.85)
63°49'N, 141°24'W

Summary: Disseminated pyrite, chalcopyrite, and sphalerite in float of altered granodiorite and/or gneiss (Paleozoic and/or Precambrian in age). Staked 1975 or 1976. Extent of mineralization not known. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 7
Eberlein and others, 1977 (OF 77-168D), p. 102, loc. 12

Pushbush

Copper, Molybdenum

Fortymile district
MF-767E, loc 11

Tanacross (21.0, 9.56)
63°32'N, 141°19'W

Summary: Disseminated pyrite, chalcopyrite, and molybdenite in Tertiary mafic and felsic volcanic rocks and biotite gneiss and schist (Paleozoic and/or Precambrian? in age). Extent of mineralization and other relations not known. Exposed by trenches. [Eberlein and others, 1977 (OF 77-168D)]

References

Singer and others, 1976 (MF-767E), loc. 11
Eberlein and others, 1977 (OF 77-168D), p. 102, loc. 13

(Sixtymile R.)

Gold

Fortymile district(?)

Tanacross(?)

NE $\frac{1}{4}$ NE $\frac{1}{4}$ quad.(?)

Summary: Rich placer gold strike near head of Sixtymile R. reported in 1911. Ground said to be about 20 ft. deep and to average \$1 per square foot of bedrock [gold at \$20.67]. Not known if strike was in Alaska or Yukon. [Porter, 1912 (B 520)]

References

Porter, 1912 (B 520), p. 212

Singer and others, 1976 (MF-767E)

Taurus

Copper, Molybdenum

Fortymile district
MF-767E, loc. 10

Tanacross (20.9, 11.9)
63°39'N, 141°19'W

Summary: Discovered in 1970 in eastern part of altered porphyry system 8 mi long and a little less than a mile wide along trend of Dawson Range (Yukon Territory) where numerous porphyry copper-molybdenum deposits are known. Disseminated chalcopyrite, pyrite, molybdenite, chalcocite, and magnetite are present in hydrothermally altered quartz monzonite to quartz porphyry high-level intrusives (mid-Cretaceous to early Tertiary in age) into upper green-schist-facies metamorphic rocks and gneiss (Paleozoic and/or Precambrian? in age). Extrusive phases may also be present. Numerous faults. Potassic, phyllic, argillic, and propylitic alteration zones recognized. May be near top of a porphyry system. Tourmaline and fluorite are present. Explored by geochemical sampling and surface and airborne geophysical work. East Taurus has been drilled. Partial to complete leaching extends to depth of about 200 ft. Chalcocite mostly above 350 ft. Chalcopyrite-molybdenite mineralization known to 918 ft, the maximum depth drilled. Of 21 diamond drill holes and numerous rotary drill holes, six have reportedly intersected intervals of 0.1% Cu or better. [Eberlein and others, 1977 (OF 77-168D)] Includes references to: East Taurus, West Taurus.

References

- Singer and others, 1976 (MF-767E), loc. 10
Eberlein and others, 1977 (OF 77-169D), p. 101, loc. 5

(Timber Cr.)

Asbestos

Tok district

Tanacross

SW $\frac{1}{4}$ SW $\frac{1}{4}$ SW $\frac{1}{4}$ quad.

Summary: Sam Gamblin reported finding asbestos in general Timber Cr. area, which is underlain by Paleozoic or Precambrian metamorphic rocks. [Joesting, 1942 (TDM 1); Singer and others, 1976 (MF-767E)]

References

Joesting, 1942 (TDM 1), p. 44

Singer and others, 1976 (MF-767E) [deposit not shown; cited for geology]

Concentrate samples

Molybdenum(?), Tin, Tungsten

Fortymile district

Tanacross

Cassiterite and scheelite were identified in the nonmagnetic fractions of heavy-mineral concentrates of samples of stream sediments or rocks from the following localities. Powellite probably is present in some or all samples.

(14.8, 8.7)	63°29'N, 142°07'W
(17.4, 1.7)	63°05'N, 141°49'W
(19.3, 0.8)	63°02'N, 141°35'W
(19.35, 1.9)	63°05'N, 141°34'W
(19.35, 2.25)	63°06'N, 141°34'W
(19.4, 1.65)	63°04'N, 141°34'W
(19.4, 3.0)	63°09'N, 141°33'W
(21.1, 7.35)	63°24'N, 141°20'W
(21.3, 14.9)	63°49'N, 141°15'W
(21.4, 9.45)	63°31'N, 141°16'W
(21.7, 14.2)	63°47'N, 141°12'W

Reference

Tripp and others, 1976 (MF-767 O), map A

Concentrate samples

Molybdenum, Tungsten

Fortymile district

Tanacross

Scheelite was identified in the nonmagnetic fractions of heavy-mineral concentrates of samples of stream sediments or rocks from the following localities. Powellite is present in some or all samples, but data are not adequate to specify which samples certainly contain powellite.

(1.8, 16.25)	63°56'N, 143°46'W
(1.8, 16.55)	63°57'N, 143°46'W
(2.65, 17.5)	64°00'N, 143°40'W
(3.15, 16.85)	63°58'N, 143°36'W
(3.55, 16.75)	63°57'N, 143°32'W
(4.3, 15.1)	63°52'N, 143°27'W
(4.4, 15.4)	63°53'N, 143°26'W
(4.55, 16.75)	63°57'N, 143°25'W
(4.9, 15.1)	63°52'N, 143°22'W
(4.9, 15.8)	63°54'N, 143°22'W
(5.1, 17.2)	63°59'N, 143°20'W
(5.15, 13.65)	63°47'N, 143°20'W
(5.8, 17.3)	63°59'N, 143°15'W
(6.35, 14.7)	63°50'N, 143°11'W
(6.7, 14.1)	63°48'N, 143°08'W
(6.7, 14.8)	63°50'N, 143°08'W
(7.1, 12.0)	63°41'N, 143°06'W
(7.15, 14.75)	63°50'N, 143°04'W
(7.6, 15.1)	63°51'N, 143°01'W
(8.15, 13.5)	63°46'N, 142°57'W
(8.2, 12.45)	63°42'N, 142°57'W
(8.25, 13.1)	63°45'N, 142°56'W
(8.55, 14.2)	63°48'N, 142°54'W
(8.8, 13.3)	63°45'N, 142°52'W
(9.25, 11.0)	63°37'N, 142°49'W
(9.35, 12.3)	63°42'N, 142°48'W
(9.6, 10.6)	63°36'N, 142°46'W
(9.7, 12.6)	63°43'N, 142°45'W
(9.8, 11.85)	63°40'N, 142°44'W
(9.85, 12.9)	63°44'N, 142°44'W
(10.35, 12.7)	63°43'N, 142°40'W
(10.5, 11.65)	63°39'N, 142°39'W
(10.55, 12.55)	63°43'N, 142°38'W
(10.7, 13.65)	63°46'N, 142°37'W
(11.15, 10.15)	63°34'N, 142°34'W
(11.2, 14.3)	63°49'N, 142°33'W
(11.3, 8.75)	63°30'N, 142°33'W
(11.45, 12.6)	63°43'N, 142°32'W
(11.5, 10.9)	63°37'N, 142°32'W

Concentrate samples

Molybdenum, Tungsten (cont.)

Fortymile district

Tanacross

(11.65, 6.75)	63°23'N, 142°31'W
(11.75, 14.8)	63°50'N, 142°29'W
(11.85, 9.4)	63°32'N, 142°29'W
(11.9, 12.25)	63°42'N, 142°28'W
(12.6, 6.45)	63°21'N, 142°24'W
(12.65, 11.5)	63°39'N, 142°22'W
(12.75, 12.0)	63°41'N, 142°22'W
(12.8, 10.95)	63°37'N, 142°21'W
(13.0, 13.7)	63°47'N, 142°19'W
(13.2, 6.2)	63°21'N, 142°19'W
(13.25, 15.1)	63°51'N, 142°17'W
(13.45, 10.6)	63°36'N, 142°17'W
(14.0, 15.75)	63°53'N, 142°11'W
(14.05, 5.0)	63°17'N, 142°13'W
(14.1, 6.45)	63°21'N, 142°13'W
(14.2, 11.15)	63°37'N, 142°11'W
(14.3, 6.6)	63°22'N, 142°11'W
(14.45, 12.4)	63°41'N, 142°09'W
(14.5, 15.3)	63°52'N, 142°07'W
(15.0, 10.9)	63°36'N, 142°05'W
(15.1, 15.75)	63°53'N, 142°03'W
(15.3, 5.6)	63°18'N, 142°03'W
(15.35, 14.35)	63°48'N, 142°01'W
(15.45, 4.7)	63°15'N, 142°03'W
(15.6, 16.15)	63°54'N, 141°59'W
(15.65, 14.65)	63°49'N, 141°59'W
(15.75, 3.35)	63°11'N, 142°01'W
(15.8, 13.75)	63°46'N, 141°58'W
(15.9, 12.55)	63°42'N, 151°57'W
(16.1, 7.65)	63°25'N, 141°57'W
(16.5, 16.2)	63°55'N, 141°52'W
(16.7, 11.9)	63°40'N, 141°51'W
(16.75, 11.6)	63°39'N, 141°51'W
(16.9, 12.4)	63°41'N, 141°50'W
(16.9, 13.6)	63°45'N, 141°49'W
(17.0, 4.0)	63°13'N, 141°51'W
(17.0, 5.35)	63°17'N, 141°51'W
(17.05, 5.05)	63°16'N, 141°50'W
(17.3, 3.25)	63°10'N, 141°49'W
(17.45, 4.5)	63°14'N, 141°48'W
(17.5, 3.65)	63°11'N, 141°47'W
(17.7, 8.9)	63°29'N, 141°45'W
(17.7, 15.75)	63°53'N, 141°42'W
(17.75, 11.25)	63°37'N, 141°43'W
(17.8, 7.85)	63°26'N, 141°44'W
(17.8, 8.7)	63°29'N, 141°44'W

Concentrate samples

Molybdenum, Tungsten

(cont.)

Fortymile district

Tanacross

(18.15, 7.6)	63°25'N, 141°41'W
(18.2, 7.3)	63°24'N, 141°41'W
(18.3, 11.9)	63°40'N, 141°39'W
(18.4, 15.75)	63°53'N, 141°37'W
(19.05, 12.2)	63°40'N, 141°33'W
(19.2, 7.95)	63°26'N, 141°34'W
(19.2, 11.0)	63°36'N, 141°32'W
(19.25, 12.9)	63°43'N, 141°31'W
(19.3, 17.6)	63°59'N, 141°30'W
(19.5, 15.4)	63°51'N, 141°29'W
(19.7, 11.65)	63°38'N, 141°29'W
(19.7, 12.2)	63°40'N, 141°28'W
(19.8, 3.85)	63°12'N, 141°30'W
(19.85, 3.55)	63°11'N, 141°30'W
(19.9, 8.15)	63°27'N, 141°28'W
(19.9, 8.85)	63°29'N, 141°28'W
(10.0, 8.55)	63°28'N, 141°27'W
(20.2, 13.6)	63°45'N, 141°24'W
(20.3, 12.6)	63°42'N, 141°23'W
(20.35, 13.15)	63°44'N, 141°23'W
(20.45, 15.55)	63°52'N, 141°21'W
(20.5, 13.85)	63°46'N, 141°21'W
(20.55, 13.65)	63°45'N, 141°21'W
(20.6, 13.3)	63°44'N, 141°21'W
(20.75, 6.0)	63°19'N, 141°22'W
(20.85, 9.9)	63°32'N, 141°20'W
(20.85, 10.5)	63°35'N, 141°20'W
(21.3, 10.35)	63°34'N, 141°16'W
(21.5, 13.05)	63°43'N, 141°14'W
(21.65, 12.1)	63°40'N, 141°13'W
(21.85, 15.55)	63°52'N, 141°10'W
(21.9, 17.6)	63°59'N, 141°09'W
(22.0, 12.8)	63°41'N, 141°10'W
(22.0, 13.15)	63°43'N, 141°10'W
(22.0, 18.0)	64°00'N, 141°09'W
(22.1, 15.7)	63°52'N, 141°09'W
(22.25, 6.2)	63°20'N, 141°11'W
(22.3, 9.8)	63°32'N, 141°09'W
(22.5, 5.0)	63°15'N, 141°09'W
(22.6, 1.7)	63°04'N, 141°09'W
(22.7, 14.35)	63°47'N, 141°04'W
(22.75, 1.2)	63°03'N, 141°08'W
(22.75, 2.6)	63°07'N, 141°08'W
(22.75, 4.0)	63°12'N, 141°07'W
(22.85, 3.55)	63°11'N, 141°07'W

Concentrate samples Molybdenum, Tungsten (cont.)

Fortymile district Tanacross

(22.85, 3.55)	63°11'N, 141°07'W
(22.85, 10.65)	63°35'N, 141°04'W
(22.9, 11.9)	63°39'N, 141°04'W
(22.9, 14.6)	63°48'N, 141°03'W
(22.95, 17.9)	63°59'N, 141°01'W
(23.0, 0.3)	63°02'N, 141°06'W
(23.0, 3.15)	63°09'N, 141°06'W
(23.05, 1.8)	63°05'N, 141°06'W
(23.05, 8.2)	63°26'N, 141°04'W
(23.2, 5.45)	63°17'N, 141°04'W
(23.2, 10.0)	63°32'N, 141°02'W
(23.2, 11.05)	63°36'N, 141°01'W
(23.2, 12.25)	63°40'N, 141°01'W
(23.25, 1.1)	63°02'N, 141°05'W
(23.4, 6.65)	63°21'N, 141°02'W

Reference

Tripp and others, 1976 (MF-767 O), map A

Concentrate samples

Molybdenum, Tungsten

Goodpaster district

Tanacross

Scheelite was identified in the nonmagnetic fractions of heavy-mineral concentrates of samples of stream sediments or rocks from the following localities. Powellite is present in some or all samples, but data are not adequate to specify which samples certainly contain powellite.

(0.25, 15.3)	63°52'N, 143°58'W
(0.7, 4.45)	63°49'N, 143°55'W
(0.8, 16.85)	63°58'N, 143°54'W
(0.85, 16.0)	63°55'N, 143°53'W
(1.0, 14.8)	63°51'N, 143°52'W
(1.05, 17.3)	63°59'N, 143°51'W
(1.1, 12.75)	63°44'N, 143°51'W
(1.3, 14.15)	63°45'N, 143°50'W
(1.7, 12.5)	63°43'N, 143°47'W
(1.9, 12.05)	63°41'N, 143°45'W
(1.9, 14.65)	63°50'N, 143°45'W
(2.1, 11.35)	63°39'N, 143°44'W
(2.15, 13.6)	63°47'N, 143°43'W
(2.2, 14.8)	63°51'N, 143°43'W
(2.3, 15.25)	63°52'N, 143°42'W
(2.4, 10.55)	63°36'N, 143°42'W
(2.4, 15.0)	63°51'N, 143°41'W
(2.6, 13.2)	63°45'N, 143°40'W
(2.65, 9.65)	63°33'N, 143°40'W
(2.7, 12.85)	63°44'N, 143°39'W
(2.75, 8.2)	63°28'N, 143°39'W
(2.75, 15.5)	63°53'N, 143°38'W
(2.8, 14.25)	63°49'N, 143°38'W
(2.85, 14.6)	63°50'N, 143°38'W
(3.05, 8.35)	63°28'N, 143°37'W
(3.1, 15.75)	63°54'N, 143°36'W
(3.2, 15.9)	63°54'N, 143°34'W
(3.4, 10.6)	63°36'N, 143°34'W
(3.4, 11.2)	63°38'N, 143°33'W
(3.4, 12.75)	63°43'N, 143°34'W
(3.5, 9.0)	63°31'N, 143°33'W
(3.5, 12.45)	63°43'N, 143°33'W
(3.6, 8.15)	63°28'N, 143°32'W
(3.6, 8.7)	63°30'N, 143°32'W
(3.8, 7.9)	63°27'N, 143°31'W
(3.85, 13.2)	63°45'N, 143°30'W
(4.05, 12.9)	63°44'N, 143°29'W
(4.25, 8.7)	63°30'N, 143°27'W
(4.8, 11.7)	63°40'N, 143°23'W

Concentrate samples Molybdenum, Tungsten (cont.)

Goodpaster district Tanacross

(4.9, 9.05)	63°31'N, 143°22'W
(5.05, 8.85)	63°30'N, 143°21'W
(5.15, 9.85)	63°39'N, 143°20'W
(5.3, 10.35)	63°35'N, 143°19'W
(5.5, 11.9)	63°41'N, 143°18'W
(6.0, 8.85)	63°30'N, 143°14'W
(6.3, 12.0)	63°41'N, 143°11'W
(7.4, 8.7)	63°30'N, 143°03'W

Reference

Tripp and others, 1976 (MF-767 O), map A

Concentrate samples

Molybdenum, Tungsten

Tok district

Tanacross

Scheelite was identified in the nonmagnetic fractions of heavy-mineral concentrates of samples of stream sediments or rocks from the following localities. Powellite is present in some or all samples, but data are not adequate to specify which samples certainly contain powellite.

(0.1, 3.4)	63°12'N, 143°59'W
(0.4, 5.9)	63°20'N, 143°57'W
(0.4, 8.0)	63°27'N, 143°57'W
(0.7, 4.4)	63°15'N, 143°55'W
(0.85, 4.15)	63°14'N, 143°54'W
(0.85, 5.4)	63°18'N, 143°54'W
(0.95, 6.85)	63°23'N, 143°53'W
(1.55, 4.4)	63°15'N, 143°48'W
(1.6, 4.6)	63°16'N, 143°48'W
(1.65, 5.1)	63°17'N, 143°47'W
(1.8, 4.0)	63°14'N, 143°46'W
(1.8, 6.2)	63°21'N, 143°46'W
(2.25, 6.45)	63°22'N, 143°43'W
(2.45, 4.5)	63°45'N, 143°41'W
(2.75, 5.9)	63°20'N, 143°39'W
(2.95, 6.1)	63°21'N, 143°37'W
(3.0, 5.8)	63°20'N, 143°37'W
(3.4, 5.3)	63°18'N, 143°34'W
(3.6, 6.45)	63°22'N, 143°33'W
(4.35, 5.65)	63°19'N, 143°27'W
(5.25, 4.9)	63°17'N, 143°20'W

Reference

Tripp and others, 1976 (MF-767 0), map A

Concentrate samples

Tin

Fortymile district

Tanacross

Cassiterite was identified in the nonmagnetic fractions of heavy-mineral concentrates of samples of stream sediments or rocks from the following localities.

(14.9, 8.2)	63°27'N, 142°06'W
(19.15, 3.9)	63°12'N, 141°35'W
(21.9, 8.3)	63°27'N, 141°12'W
(22.15, 7.4)	63°24'N, 141°11'W
(22.2, 7.85)	63°25'N, 141°10'W

Reference

Tripp and others, 1976 (MF-767 O), map A

Synonyms, Owners, Operators, and Claim Names

American Smelting & Refining Co. -- see Blue Lead
Caribou Gold Dredging Co. -- see (Caribou Cr.)
Caribou Mining Co. -- see (Caribou Cr.)
Goodpaster Exploration -- see Gray Quartz Lead
Hidden Treasure -- see (Tibbs Cr.)
Hirschberger & Johnson -- see Blue Lead
Martha -- see (Banner Cr.)
McCord -- see Blue Lead
Michigan Lead -- see (Tibbs Cr.)
Noribu -- see (Nickel Cr.)
Puntilla -- see (Tenderfoot Cr.)
(Pyne Cr.) -- see (Pine Cr.)
Ricks -- see (Nickel Cr.)
Salcha Mining Co. -- see (Caribou Cr.)
Shuff -- see Democrat
Stevens & Johnson -- see (Caribou Cr.)
Tibbett -- see Blue Lead

Synonyms, Owners, Operators, and Claim Names

A Lucky Leak -- see (Boulder Cr.)
Caulk -- see (Boulder Cr.)
East Taurus -- see Taurus
Gamblin (& Murhoffer) -- see (Boulder Cr.)
(Lost Cr.) -- see Frey
Paternie -- see Peternie
Rambler -- see (Boulder Cr.)
Stibnite -- see (Boulder Cr.)
(Stibnite Cr.) -- see (Boulder Cr.)
(Tok R.) -- see (Boulder Cr.)
West Taurus -- see Taurus

References Cited

Big Delta quadrangle

Berg, H. C., and Cobb, E. H., 1967, Metalliferous lode deposits of Alaska: U.S. Geological Survey Bulletin 1246, 254 p.

Brooks, A. H., 1906, The mining industry in 1905: U.S. Geological Survey Bulletin 284, p. 4-9.

Brooks, A. H., 1907, The mining industry in 1906: U.S. Geological Survey Bulletin 314, p. 19-39.

Brooks, A. H., 1908, The mining industry in 1907: U.S. Geological Survey Bulletin 345, p. 30-53.

Brooks, A. H., 1909, The mining industry in 1908: U.S. Geological Survey Bulletin 379, p. 21-62.

Brooks, A. H., 1915, The Alaskan mining industry in 1914: U.S. Geological Survey Bulletin 622, p. 15-68.

Brooks, A. H., 1916, The Alaskan mining industry in 1915: U.S. Geological Survey Bulletin 642, p. 16-71.

Brooks, A. H., 1918, The Alaskan mining industry in 1916: U.S. Geological Survey Bulletin 662, p. 11-62.

Brooks, A. H., 1923, The Alaska mining industry in 1921: U.S. Geological Survey Bulletin 739, p. 1-44.

Brooks, A. H., and Capps, S. R., 1924, The Alaska mining industry in 1922: U.S. Geological Survey Bulletin 755, p. 3-49.

Bundtzen, T. K., and Reger, R. D., 1977, The Richardson lineament - a structural control for gold deposits in the Richardson mining district, Alaska: Alaska Division of Geological and Geophysical Surveys Geologic Report 55, p. 29-34.

Chapin, Theodore, 1914, Placer mining in the Yukon-Tanana region: U.S. Geological Survey Bulletin 592, p. 357-362.

Cobb, E. H., 1972, Metallic mineral resources map of the Big Delta quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-388, 1 sheet, scale 1:250,000.

Big Delta quadrangle

Cobb, E. H., 1973, Placer deposits of Alaska: U.S. Geological Survey Bulletin 1374, 213 p.

Cobb, E. H., 1977, Placer deposits map of central Alaska: U.S. Geological Survey Open-file Report 77-168B, 64 p. - map, scale 1:1,000,000.

Cobb, E. H., 1979, Summary of references to mineral occurrences (other than mineral fuels and construction materials) in the Mount Hayes quadrangle, Alaska: U.S. Geological Survey Open-file Report 79-238, 140 p.

Cobb, E. H., and Kachadoorian, Reuben, 1961, Index of metallic and nonmetallic mineral deposits of Alaska compiled from published reports of Federal and State agencies through 1959: U.S. Geological Survey Bulletin 1139, 363 p.

Eakin, H. M., 1915, Mining in the Fairbanks district: U.S. Geological Survey Bulletin 622, p. 229-238.

Eberlein, G. D., Chapman, R. M., Foster, H. L., and Gassaway, J. S., 1977, Map and table describing known metalliferous and selected nonmetalliferous mineral deposits in central Alaska: U.S. Geological Survey Open-file Report 77-168D, 132 p. + map, scale 1:1,000,000.

Eberlein, G. D., and Menzie, W. D., 1978, Maps and tables describing areas of metalliferous mineral resource potential of central Alaska: U.S. Geological Survey Open-file Report 78-1-D, 43 p. + 2 pl., scale 1:1,000,000.

Ellsworth, C. E., 1910, Placer mining in the Yukon-Tanana region: U.S. Geological Survey Bulletin 442, p. 230-245.

Ellsworth, C. E., and Davenport, R. W., 1913, Placer mining in the Yukon-Tanana region: U.S. Geological Survey Bulletin 542, p. 203-222.

Ellsworth, C. E., and Parker, G. L., 1911, Placer mining in the Yukon-Tanana region: U.S. Geological Survey Bulletin 480, p. 153-172.

- Foster, H. L., Albert, N. R. D., Griscom, Andrew, Hessin, T. D., Menzie, W. D., Turner, D. L., and Wilson, F. H., 1979, The Alaskan Mineral Resource Assessment Program: Background information to accompany folio of geologic and mineral resource maps of the Big Delta quadrangle, Alaska: U.S. Geological Survey Circular 783, 19 p.
- Hasler, J. W., Miller, M. H., and Chapman, R. M., 1973, Bismuth, in Brobst, D. A., and Pratt, W. P., eds., United States mineral resources: U.S. Geological Survey Professional Paper 820, p. 95-98.
- Joesting, H. R., 1942, Strategic mineral occurrences in interior Alaska: Alaska Department of Mines Pamphlet 1, 46 p.
- Koschmann, A. H., and Bergendahl, M. H., 1968, Principal gold-producing districts of the United States: U.S. Geological Survey Professional Paper 610, 283 p.
- MacKevett, E. M., Jr., and Holloway, C. D., 1977, Map showing metal-liferous and selected nonmetalliferous mineral deposits in the eastern part of southern Alaska: U.S. Geological Survey Open-file Report 77-169A, 1 sheet + 99 p. tabular material, scale 1:1,000,000.
- Martin, G. C., 1919, The Alaskan mining industry in 1917: U.S. Geological Survey Bulletin 692, p. 11-42.
- Menzie, W. D., and Foster, H. L., 1979, Metalliferous and selected nonmetalliferous mineral resource potential in the Big Delta quadrangle, Alaska: U.S. Geological Survey Open-file Report 78-529 D, 61 p. + 1 sheet, scale 1:250,000.
- Mulligan, J. J., 1974, Mineral resources of the trans-Alaska pipeline corridor: U.S. Bureau of Mines Information Circular 8626, 24 p.
- Prindle, L. M., 1906, Yukon placer fields: U.S. Geological Survey Bulletin 284, p. 109-127.
- Prindle, L. M., 1906, The Yukon-Tanana region, Alaska: Description of Circle quadrangle: U.S. Geological Survey Bulletin 295, 27 p.

Big Delta quadrangle

- Prindle, L. M., 1908, The Fairbanks and Rampart quadrangles, Yukon-Tanana region, Alaska, with a section on the Rampart placers, by F. L. Hess, and a paper on the water supply of the Fairbanks region, by C. C. Covert: U.S. Geological Survey Bulletin 337, 102 p.
- Prindle, L. M., 1913, A geologic reconnaissance of the Circle quadrangle, Alaska: U.S. Geological Survey Bulletin 538, 82 p.
- Prindle, L. M., and Katz, F. J., 1913, Geology of the Fairbanks district, in Prindle, L. M., A geologic reconnaissance of the Fairbanks quadrangle, Alaska: U.S. Geological Survey Bulletin 525, p. 59-152.
- Ransome, A. L., and Kerns, W. H., 1954, Names and definitions of regions, districts, and subdistricts in Alaska (used by the Bureau of Mines in statistical and economic studies covering the mineral industry of the Territory): U.S. Bureau of Mines Information Circular 7679, 91 p.
- Saunders, R. H., 1965, A geochemical investigation in the Richardson area, Big Delta quadrangle, Alaska: Alaska Division of Mines and Minerals Geochemical Report 3, 11 p.
- Smith, P. S., 1926, Mineral industry of Alaska in 1924: U.S. Geological Survey Bulletin 783, p. 1-30.
- Smith, P. S., 1929, Mineral industry of Alaska in 1926: U.S. Geological Survey Bulletin 797, p. 1-50.
- Smith, P. S., 1930, Mineral industry of Alaska in 1927: U.S. Geological Survey Bulletin 810, p. 1-64.
- Smith, P. S., 1930, Mineral industry of Alaska in 1928: U.S. Geological Survey Bulletin 813, p. 1-72.
- Smith, P. S., 1932, Mineral industry of Alaska in 1929: U.S. Geological Survey Bulletin 824, p. 1-61.
- Smith, P. S., 1933, Mineral industry of Alaska in 1930: U.S. Geological Survey Bulletin 836, p. 1-83.
- Smith, P. S., 1933, Mineral industry of Alaska in 1931: U.S. Geological Survey Bulletin 844-A, p. 1-82.

- Smith, P. S., 1934, Mineral industry of Alaska in 1932: U.S. Geological Survey Bulletin 857-A, p. 1-91.
- Smith, P. S., 1934, Mineral industry of Alaska in 1933: U.S. Geological Survey Bulletin 861-A, p. 1-94.
- Smith, P. S., 1936, Mineral industry of Alaska in 1934: U.S. Geological Survey Bulletin 868-A, p. 1-91.
- Smith, P. S., 1937, Mineral industry of Alaska in 1935: U.S. Geological Survey Bulletin 880-A, p. 1-95.
- Smith, P. S., 1938, Mineral industry of Alaska in 1936: U.S. Geological Survey Bulletin 897-A, p. 1-107.
- Smith, P. S., 1939, Mineral industry of Alaska in 1937: U.S. Geological Survey Bulletin 910-A, p. 1-113.
- Smith, P. S., 1939, Mineral industry of Alaska in 1938: U.S. Geological Survey Bulletin 917-A, p. 1-113.
- Smith, P. S., 1941, Mineral industry of Alaska in 1939: U.S. Geological Survey Bulletin 926-A, p. 1-106.
- Smith, P. S., 1942, Mineral industry of Alaska in 1940: U.S. Geological Survey Bulletin 933-A, p. 1-102.
- Smith, S. S., 1917, The mining industry in the Territory of Alaska during the calendar year 1915: U.S. Bureau of Mines Bulletin 142, 66 p.
- Thomas, B. I., 1970, Reconnaissance of the gold-bearing quartz veins in the Tibbs Creek area, Goodpaster River, Big Delta quadrangle, central Alaska: U.S. Bureau of Mines Open-file Report 14-70, 12 p.
- Wedow, Helmuth, Jr., Killeen, P. L., and others, 1954, Reconnaissance for radioactive deposits in eastern interior Alaska, 1946: U.S. Geological Survey Circular 331, 36 p.

- MacKevett, E. M., Jr., and Holloway, C. D., 1977, Map showing metal-liferous and selected nonmetalliferous mineral deposits in the eastern part of southern Alaska: U.S. Geological Survey Open-file Report 77-169A, 1 sheet + 99 p. tabular material, scale 1:1,000,000.
- Moffit, F. H., 1938, Geology of the Slana-Tok district, Alaska: U.S. Geological Survey Bulletin 904, 54 p.
- Moffit, F. H., 1944, Mining in the northern Copper River region, Alaska: U.S. Geological Survey Bulletin 943-B, p. 25-47.
- Moffit, F. H., 1954, Geology of the eastern part of the Alaska Range and adjacent area: U.S. Geological Survey Bulletin 989-D, p. 63-218.
- Porter, E. A., 1912, Placer mining in the Fortymile, Eagle, and Seventymile River districts: U.S. Geological Survey Bulletin 520, p. 211-218.
- Ransome, A. L., and Kerns, W. H., 1954, Names and definitions of regions, districts, and subdistricts in Alaska (used by the Bureau of Mines in statistical and economic studies covering the mineral industry of the Territory): U.S. Bureau of Mines Information Circular 7679, 91 p.
- Richter, D. H., 1967, Geology of the upper Slana-Mentasta Pass area, southcentral Alaska: Alaska Division of Mines and Minerals Geologic Report 30, 25 p.
- Singer, D. A., Curtin, G. C., and Foster, H. L., 1976, Mineral resources map of the Tanacross quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-767E, 1 sheet, scale 1:25,000.
- Smith, P. S., 1942, Mineral industry of Alaska in 1940: U.S. Geological Survey Bulletin 933-A, p. 1-102.
- Tripp, R. B., Curtin, G. C., Day, G. W., Karlson, R. C., and Marsh, S. P., 1976, Maps showing mineralogical and geochemical data for heavy-mineral concentrates in the Tanacross quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-767 O, 2 sheets, scale 1:500,000.

References Cited

Tanacross quadrangle

Berg, H. C., and Cobb, E. H., 1967, Metalliferous lode deposits of Alaska: U.S. Geological Survey Bulletin 1246, 254 p.

Brooks, A. H., 1916, Antimony deposits of Alaska: U.S. Geological Survey Bulletin 649, 67 p.

Cobb, E. H., 1972, Metallic mineral resources map of the Tanacross quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-383, 1 sheet, scale 1:250,000.

Cobb, E. H., and Kachadoorian, Reuben, 1961, Index of metallic and nonmetallic mineral deposits of Alaska compiled from published reports of Federal and State agencies through 1959: U.S. Geological Survey Bulletin 1139, 363 p.

Eberlein, G. D., Chapman, R. M., Foster, H. L., and Gassaway, J. S., 1977, Map and table describing known metalliferous and selected nonmetalliferous mineral deposits in central Alaska: U.S. Geological Survey Open-file Report 77-168D, 132 p. + map, scale 1:1,000,000.

Ebbley, Norman, Jr., and Wright, W. S., 1948, Antimony deposits in Alaska: U.S. Bureau of Mines Report of Investigations 4173, 41 p.

Ellsworth, C. E., and Davenport, R. W., 1913, Placer mining in the Yukon-Tanana region: U.S. Geological Survey Bulletin 542, p. 203-222.

Foster, H. L., 1970, Reconnaissance geologic map of the Tanacross quadrangle, Alaska: U.S. Geological Survey Miscellaneous Geological Investigations Map I-593, 1 sheet, scale 1:250,000.

Foster, H. L., Albert, N. R. D., Barnes, D. F., Curtin, G. C., Griscom, Andrew, Singer, D. A., and Smith, J. G., 1976, The Alaskan Mineral Resource Assessment Program: Background information to accompany folio of geologic and mineral resource maps of the Tanacross quadrangle, Alaska: U.S. Geological Survey Circular 734, 23 p.

Joesting, H. R., 1942, Strategic mineral occurrences in interior Alaska: Alaska Department of Mines Pamphlet 1, 46 p.