

UNITED STATES DEPARTMENT OF THE INTERIOR

GEOLOGICAL SUREVY

PETROLEUM GEOLOGY OF ROCKS EXPOSED ON KODIAK  
ISLAND, ALASKA:  
A REPORT FOR THE FEDERAL LANDS ASSESSMENT  
PROGRAM (FLAP)

by

Michael A. Fisher <sup>1</sup>

Open File Report 88-385

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or stratigraphic nomenclature. Any use of trade names is for purposes of identification only and does not imply endorsement by the U.S. Geological Survey.

<sup>1</sup>U. S. Geological Survey  
345 Middlefield Rd  
MS 999  
Menlo Park, CA 94025






Figure 1. Index map of Kodiak islands and the Gulf of Alaska

## INTRODUCTION

### Geography and History of Petroleum Exploration

Kodiak Island forms part of the western rim of the Gulf of Alaska (Figure 1). This island, together with two other main and several lesser islands (Figure 2), cluster within an offshore area that was once explored intensively for petroleum. This exploration entailed primarily collecting seismic reflection data, but also six Continental Offshore Stratigraphic Test (COST) wells were drilled in Federal waters through the continental shelf that borders these islands on the southeast. Despite initially high optimism for finding hydrocarbons within this offshore area, the petroleum industry's attention shifted elsewhere. After polling petroleum exploration companies in 1986, the Minerals Management Service of the Department of the Interior decided that the low level of

interest did not warrant a sale of offshore leases.

Some of the six offshore COST wells bottomed in Eocene rocks that dip northwest and apparently correlate with rocks of the same age exposed extensively onshore (Turner et al., 1987). Although offshore seismic reflection data do not reveal the structure of the Eocene rocks, these data show that highly reflective Miocene and younger (McClellan et al., 1980; Keller et al., Turner et al., 1987) rocks unconformably overlie the Eocene units and fill three deep (about 5 km) basins under the continental shelf (Fisher and von Huene, 1980). This thick Neogene section is known from seismic reflection data to thin sharply toward the islands, so that rocks of this age outcrop onshore at only scattered, local sites, where they form a thin veneer that unconformably overlies the much more extensively exposed, highly deformed Cretaceous and Paleogene rocks. Hence, offshore well data contribute little to the petroleum assessment of the onshore rocks.




Figure 2. Generalized geologic map of Kodiak Island and adjacent islands. Patterns that show lithologic units are explained in Figure 3. BRF: Border ranges Fault; ERF Eagle River fault; CF Contact Fault.

### Structural Setting

No identifiable basin lies within the area of the Kodiak island group. Although Cretaceous and Paleogene rocks onshore are presumed to be thick (no accurate measurement of their total thickness has yet

been made), this thickness results from folding and from structural repetition of stratigraphic units along thrust faults. Onshore Cretaceous and Paleogene rocks lie within the forearc area of a convergent margin, and abundant evidence suggests that their present structural aspect results




Figure 3. Explanation for Figure 2.

directly from a protracted history of convergent-margin tectonics. For example, paleomagnetic data obtained from Paleocene basalt and Eocene sedimentary rock exposed on Kodiak Island (Plumley et al., 1982, 1983; Moore et al., 1983) suggest that these rocks were deposited far south of their present position and migrated northward. The collision between the allochthonous rocks under Kodiak Island with what was then the continental margin of Alaska may have been responsible for the pervasive deformation of insular rocks by thrust faults and compressed folds. Microscopic examination of surface samples of the strongly deformed turbidite sequence indicates that the rocks apparently underwent ductile deformation at depths greater than about 10 km (Sample and Moore, 1987). Strong uplift and

erosion have since exposed these rocks at the surface.

Three tectonostratigraphic terranes, the Peninsular, the Chugach and the Prince William, underlie the Kodiak group of islands. Along the northwest coasts of the Kodiak islands, two tectonostratigraphic terranes lie sutured along the Border Ranges fault (Jones, et al., 1981; Fisher and von Huene, 1984; Figure 2). The Peninsular terrane extends northwestward from this suture zone, and the Chugach terrane extends southeastward from this fault to the Contact fault. The Prince-William terrane is represented by Paleogene rocks exposed in a narrow band along the southeast coast of the Kodiak Islands, southeast of the Contact fault.

### Lithology and Age of Rock Units

This brief description of the lithologies and ages of rocks exposed on the islands is derived from Moore (1967), Connelly and Moore (1979), Moore et al. (1983 and references therein) and Byrne (1984) as well as from other reports and maps listed in the bibliography. The geographic distribution of lithologic units exposed on the Kodiak islands (Figure 2) is controlled mainly by major thrust faults, which confine these units to four main belts. The first belt lies along the northwest coasts of the islands, northwest of the Border Ranges fault, and includes rocks of the Peninsular terrane, which are Triassic sandstone and greenstone as well as Early Jurassic plutonic rocks. This belt also contains metamorphic rocks that range from blueschist to amphibolite facies (Figure 2). The Chugach terrane includes the second and third fault-bounded lithologic belts. The second belt lies between the Border Ranges and Eagle River (or Uganik) thrust faults, and includes the Cretaceous melange of the Uyak Complex, which contains blocks of Paleozoic

limestone, chert and greenstone that are encased within a foliated claystone matrix. The youngest fossils recovered from the matrix of this melange indicate a middle Cretaceous age. The third fault-bounded belt lies between the Eagle River and Contact faults and includes strongly deformed, uppermost Cretaceous turbidite sequences (Kodiak Formation) that are rich in silty sandstone and siltstone. The fourth fault-bounded belt encompasses the Prince William terrane and lies southeast of the Contact fault. This belt includes strongly deformed turbidite sequences, melange and locally exposed basalt that are assigned to the Paleocene and Eocene Ghost Rocks Formation. The fourth belt also includes deformed turbidites assigned to the Eocene and Oligocene Sitkalidak Formation. Paleocene plutonic rocks intrude the Kodiak and Ghost Rocks Formations and show that the Chugach and Prince William terranes were together at the beginning of the Cenozoic.

Post-Eocene rocks outcrop in scattered patches, primarily on the small islands that lie south of Kodiak Island (Figure 2). These rocks unconformably overlie the Oligocene and older turbidite sequences and include upper Oligocene nonmarine and early Miocene marine sandstone as well as upper Miocene sandstone and conglomerate (T. Byrne, oral commun., 1987). In addition, one small outcrop of rocks of middle (macrofossils) or late (microfossils) Miocene age crops out on northern Kodiak Island.

### **Petroleum Geology**

Most rocks exposed on the Kodiak islands are so strongly deformed and highly indurated that few traps or reservoir rocks for hydrocarbons are present. This bleak assessment, however, does not pertain to Miocene and younger rocks, whose limited onshore exposure means that

they could contain significant hydrocarbons only in offshore areas.

Values of vitrinite reflectance and thermal alteration index (TAI) from surface samples suggest that Eocene and Oligocene rocks of the Sitkalidak Formation as well as Neogene rocks are either immature or just barely mature for generating hydrocarbons (Lyle et al., 1977, Fisher, 1980; Moore et al., 1983). These rocks generally contain less than 0.4% of organic carbon that is contained in woody and herbaceous kerogen. In contrast, Eocene and Paleocene rocks that are included within the Ghost Rocks Formation are overmature for hydrocarbons. Petrographic analysis of thin sections made from Upper Cretaceous rocks (Kodiak Formation) reveals prehnite, suggesting that these rocks have been thermally altered far beyond the thermal zone of hydrocarbon generation (Sample and Moore, 1987).

Scattered measurements of porosity and permeability indicate that, despite their generally low level of thermal alteration, post-Eocene rocks would make poor reservoirs for hydrocarbons. Maleable grains within these rocks have been deformed so that they occlude pore space. The pores in rocks of Paleocene and older age have been filled completely by diagenetic minerals. This diagenesis may have resulted from a high flux of fluids through the convergent margin where the rocks were deformed (Vrolijk, 1987).

### **PRINCIPLE PLAYS FOR PETROLEUM**

The foregoing discussion should have made clear that no conventional plays for petroleum exploration exist within rocks exposed on Kodiak Island or adjacent islands. In my opinion, onshore rocks make unpromising candidates for hydrocarbon exploration. Furthermore, rocks of Miocene and younger age that fill offshore basins thin markedly toward the islands, so that

within three miles of the shoreline, these rocks are thin and unlikely to contain hydrocarbons in economic quantities.

#### SELECTED REFERENCES

- Byrne, T., 1984, Early deformation in melange terranes of the Ghost Rocks Formation, Kodiak Islands, Alaska: in Raymond L. A., ed., *Melanges: their nature, origin, and significance*: Geological Society of America Special Paper 198, p. 21-52.
- Carden, J.R., Connelly, W., Forbes, R.B., and Turner, D.L., 1977, Blueschists of the Kodiak Islands, Alaska: An extension of the Seldovia schist terrane: *Geology*, v. 5, p. 529-533.
- Connelly, W., 1978, Uyak Complex, Kodiak Islands: A Cretaceous subduction complex: *Geological Society of America Bulletin*, v. 89, p. 755-769.
- Connelly, W., and Moore, J.C., 1979, Geologic map of the northwest side of the Kodiak Islands, Alaska, U.S. Geological Survey Miscellaneous Field Studies Map MF-1057, 2 sheets.
- Fisher, D.M., and Byrne, T., 1987, Structural evolution of underthrust sediments, Kodiak islands, Alaska: *Tectonics*, v. 6, p. 775-793.
- Fisher, M.A., 1980, Petroleum geology of the Kodiak shelf, Alaska: *American Association of Petroleum Geologists Bulletin*, v. 64, p. 1140-1157.
- Fisher, M.A., and von Huene, R., 1980, Structure of upper Cenozoic strata beneath Kodiak shelf, Alaska: *American Association of Petroleum Geologists Bulletin*, v. 64, p. 1014-1033.
- Fisher, M.A., and von Huene, R., 1984, Geophysical investigation of a suture zone: the Border Ranges fault of southern Alaska: *Journal of Geophysical Research*, v. 89, p. 11333-11351.
- Hill, M., Morris, J., and Whelan, J., 1981, Hybrid granodiorites intruding the accretionary prism, Kodiak, Shumagin, and Sanak Islands, southwest Alaska: *Journal of Geophysical Research*, v. 86, p. 10569-10590.
- Jones, D.L., Silberling, N.J., Berg, H.C., and Plafker, G., 1981, Map showing tectonostratigraphic terranes of Alaska, columnar sections, and summary description of terranes: U.S. Geological Survey Open File Report 81-792, 20 pp, 2 sheets.
- Keller, G., von Huene, R., McDougall, K., and Bruns, T., 198, Paleoclimactic evidence for Cenozoic migration of Alaskan terranes: *Tectonics*, v. 3, p. 473-495.
- Kienle, J., and Turner, D.L., 1976, The Shumagin-Kodiak batholith—A Paleocene magmatic arc?: Short notes on Alaskan Geology, Alaska Division of Geological and Geophysical Surveys Report 51, p. 9-15.
- Lyle, W., J. Morehouse, I.F. Palmer, J.G. Bolm, G.W. Moore and T.H. Nilsen, 1977, Tertiary formations in the Kodiak Island area, Alaska, and their petroleum reservoir and source-rock potential: Alaska Open File Report 114, 48 p.
- McClellan, P.H., Fisher, M.A., Barron, J.A., Arnal, R.E., Kling, S.A., and Moore, G.W., 1980, New biostratigraphic results of dredging and dart coring in the western Gulf of Alaska and their tectonic implications: U.S. Geological Survey Open-File Report 80-1237, 14 p.
- Moore, G.W., 1967, Preliminary geologic map of Kodiak Island and vicinity; U.S. Geological Survey Open file Report 271, 1 sheet.
- Moore, J.C., 1973, Cretaceous continental margin sedimentation southwestern

- Alaska: Geological Society of America Bulletin, v. 84, p 595-614.
- Moore, J.C., and Wheeler, R.L., 1978, Structural fabric of a melange Kodiak Islands, Alaska: American Journal of Science, v, 278, p. 739-765.
- Moore, J.C., Byrne, T., Plumley, P.W., Reid, M. Gibbons, H. and Coe, R.S., 1983 Paleogene evolution of the Kodiak Islands, Alaska: consequences of ridge-trench interaction in a more southerly latitude: Tectonics, v. 2, P. 265-293.
- Nilsen, T.H., and Moore, G.W., 1979, Reconnaissance study of Upper Cretaceous to Miocene stratigraphic units and sedimentary facies, Kodiak and adjacent islands, Alaska, with a section of sedimentary petrography by G.R. Winkler: U.S. Geological Survey Professional Paper 1093, 34 pp.
- Plumley, P.W., Coe, R.S., and Byrne, T., 1983, Paleomagnetism of the Ghost Rocks formation, Prince William terrane, Alaska: Tectonics, v. 2, p. 295-314.
- Plumley, P.W., Coe, R.S., Byrne, T., Reid, M.R., and Moore, J.C., 1982, Paleomagnetism of volcanic rocks of the Kodiak Islands indicates northward latitudinal displacement: Nature, v. 300, p. 50-52.
- Sample, J.C., and Moore, J.C., 1987, Structural style and kinematics of an underplated slate belt, Kodiak and Adjacent islands, Alaska: Geological Society of America Bulletin, v. 99, p 7-20.
- Turner, R. F., M.B. Lynch, T.A. Conner, P.J. Hallin, P.J. Hoose, G.C. Martin, D.L. Olson, J.A. Larson, T.O. Flett, K.W. Sherwood, and A.J. Adams, 1987, Geological and Operational summary, Kodiak shelf stratigraphic test wells, western Gulf of Alaska: OCS Report MMS 87-0109, Minerals Management Service, 340 p.
- Vrolijk, P., 1987, Tectonically driven fluid flow in the Kodiak accretionary complex, Alaska: Geology, v. 15, p. 466-469.