

DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

Reconnaissance Geology and Exploration Geochemistry
of King Cove, Alaska Peninsula

by

Gregory D. DuBois¹, Frederic H. Wilson¹,
Robert L. Detterman², and Roy T. Hopkins Jr.³

Open file report 89-350

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards (or with the North American Stratigraphic Code). Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

¹ 4200 University Dr., Anchorage, Alaska 99508-4667

² 345 Middlefield Road, Menlo Park, California 94025

³ Box 25046, Denver Federal Center, Denver, Colorado 80225

1989

Reconnaissance Geology and Exploration Geochemistry of
King Cove, Alaska Peninsula

by

Gregory D. DuBois, Frederic H. Wilson, Robert L. Detterman,
and Roy T. Hopkins Jr.

ABSTRACT

Volcanic sandstone and black coaly shale at Indian Head and northwest of Indian Head in Belkofski Bay, originally called (in part) the "arkose" unit and (in part) the Belkofski Tuff, and later assigned to the Belkofski Formation are herein reassigned to the Stepovak Formation. Volcanic and volcanoclastic sedimentary rocks on the west shore of Deer Island, previously designated as Tachini(?) Formation are herein assigned to the Unga Formation.

The Belkofski Formation consists of the same volcanic and volcanoclastic sedimentary rock lithologies as the Unga Formation but is contact metamorphosed. As mapped the Belkofski Formation occurs near plutons whereas the Unga Formation tends to be distant from plutons. The Unga and Belkofski Formations are not known to be in contact, however both overlie the Stepovak Formation. We suggest that the Belkofski Formation may be contact-metamorphosed Unga Formation.

The quartz diorite pluton in the vicinity of King Cove has a greater areal extent than previously mapped. The roof of the pluton contacts the Belkofski Formation at the high tide line of King Cove. Volcanic intrusive centers shown on ridge crests in the vicinity of King Cove may be components of the Belkofski Formation or may be equivalent to Miocene volcanic rocks to the northeast.

Geochemical analyses show low level Pb and other metal anomalies at the contact of the King Cove pluton and the Belkofski Formation.

INTRODUCTION

Five days of field work in the King Cove vicinity was done in July of 1988 as part of the Regional Alaska Mineral Resource Assessment Program (RAMRAP) for the Alaska Peninsula. King Cove is located near the southwestern end of the Alaska Peninsula, approximately 1000 km southwest of Anchorage, Alaska (fig. 1). Field work was accomplished primarily by road work and foot traverses along ridges and beaches, with limited helicopter support. Previous geologic mapping in the area was by Kennedy and Waldron (1955), Waldron (1961), Burk (1965), and McLean and others (1978).

We suggest several revisions to the above referenced geologic maps in the King Cove vicinity; these revisions are shown on the accompanying geologic map (fig. 2). The position and correlation of the Belkofski Formation is one of the outstanding problems in the Tertiary stratigraphy of the Alaska Peninsula. Data collected as part of this project provide some evidence to clarify the relation of the Belkofski to other stratigraphic units on the Alaska Peninsula.

Seventy three rock samples were collected for semi-quantitative geochemical analysis and the analytical results are included in this report (table 1).

GEOLOGY and DISCUSSION

Stepovak Formation

Strata that outcrop at Indian Head and along the shore northwest of Indian Head on Belkofski Bay (fig. 2, locality A) consist of cross-bedded volcanoclastic sandstone with pebble conglomerate and olive-gray siltstone with plant fragments. The sandstone is tuffaceous, has blotchy green and white spots characteristic of laumontite, and is composed of low-angle tabular cross-bed sets with pebble conglomerate layers between bed sets. Some bed sets are bioturbated and some are interbedded with dark brown to black coaly siltstone with plant fragments and leaves. We interpret the environment of deposition as a beach fronting a swamp. These units were originally designated (in part) the "arkose" unit and (in part) the Belkofski Tuff by Kennedy and Waldron (1947, 1955); Burk (1965) and McLean and others (1979) later mapped these rocks as Belkofski Formation. We tentatively reassign these rocks to the Stepovak Formation on the basis of their distinct difference from other rocks assigned to the Belkofski Formation and their similarity to sandstone of the Stepovak Formation. These outcrops are approximately 50 kilometers southwest of the previously mapped limit of the Stepovak Formation.

Unga Formation

Rocks on the west shore of Deer Island, south of West Cape (fig. 2, locality B), consist of lahars, volcanic conglomerate, breccia, sandstone, and mudstone cut by basalt dikes. The lahars and conglomerate are composed of a volcanic mud matrix with well rounded volcanic clasts up to 25 centimeters in diameter. The volcanic mudstone varies in color from dark brown to green with some carbonaceous layers and sandy layers in cut and fill channels. The sandstone is volcanic, dark brown, with interbedded conglomerate and carbonaceous layers. These strata were mapped as Tertiary volcanic flows and breccias by Burk (1965) and as Tachilni(?) Formation by McLean and others (1978). The rocks are distinctly unlike those found at the type locality of the Tachilni Formation on Cape Tachilni. Below are descriptions of the Unga and Tachilni Formations from their type localities, adapted from Detterman and others (in press) for comparison.

The Unga Formation consists of volcanoclastic sedimentary rocks: sandstone, conglomerate, siltstone, mudstone, shale, carbonaceous shale, debris flows and coal seams. The sandstone varies from light olive brown to yellowish gray, is fine- to coarse-grained and contains some pebble clasts. It is locally interbedded with siltstone, shale and carbonaceous shale. The lahars are massive, with volcanic mud matrices and angular volcanic clasts varying from pebble to boulder size. Thin marine sediments intertongued with thick non-marine sediments locally contain abundant molluscan fauna of early middle Miocene age (Detterman and others, in press).

The Tachilni Formation consists mainly of fossil-rich, poorly consolidated, marine, subgreywacke sandstone, interbedded with volcanic pebble conglomerate and siltstone. The sandstone is gray to brown, fine- to medium-grained, cross-bedded with pebble layers, and is composed of 30-35 percent angular quartz, 10-15 percent feldspar, 5 percent pyroxene and amphibole, and 30 percent volcanic rock fragments. The pebble conglomerate is thick-bedded to massive with a sand matrix. Both the sandstone and conglomerate contain mollusks of late Miocene age (Detterman and others, in press).

The Deer Island volcanoclastic sedimentary rocks consist of volcanic sandstone, carbonaceous shale, lahars and conglomerates with large volcanic clasts. The Unga Formation is composed of virtually the same materials, whereas the Tachilni Formation is fossil-rich and consists of a poorly consolidated subgreywacke sandstone with interbedded pebble conglomerate. We suggest assignment of these rocks to the Unga Formation on the basis of their lithologic similarity.

Belkofski Formation

Originally named the Belkofski Tuff by Kennedy and Waldron (1955), later called the Belkofski Formation by Burk (1965); this unit is characterized by beds of volcanic agglomerate, breccia, tuff, sandstone and conglomerate. It is commonly intensely silicified and sericitized where near intrusive contacts. Color ranges from gray to green and includes a distinctive purple. Rocks of the Belkofski Formation have been mapped from Cold Bay to Pavlof Bay (fig. 1). The age of the Belkofski Formation is uncertain. Fossils are rare and not age-diagnostic.

Detterman and others (in press) tentatively correlate the Belkofski Formation with either the Unga Formation or the Stepovak Formation on the basis of lithology. The rocks of the Belkofski Formation that Detterman and others (in press) tentatively correlate with the Stepovak Formation are the outcrops near Indian Head discussed above. The remaining rocks of the Belkofski Formation except for their alteration, have many lithologic similarities with the Unga Formation. We suggest in this report that the Belkofski Formation is hydrothermally altered and contact metamorphosed Unga Formation.

The Stepovak Formation underlies the Belkofski Formation (this report) and also unconformably underlies the Unga Formation. The Belkofski and Unga Formations are not seen in contact; however, the recognition of Unga Formation on Deer Island indicates that the Belkofski Formation lies entirely within the Unga depositional basin. The major outcrop areas of Belkofski Formation are all in near proximity to plutonic rocks and have been contact metamorphosed and hydrothermally altered, which is uncharacteristic of the Unga Formation. Contact metamorphism of the Unga Formation would yield rocks similar in character to the Belkofski Formation. Therefore, we suggest that the Belkofski Formation is altered Unga Formation.

Intrusive Rocks

A quartz diorite pluton intrudes the Belkofski Formation in the vicinity of King Cove and King Cove Lagoon. Its composition ranges from quartz diorite to hornblende quartz diorite, and it is locally hydrothermally altered. No age determinations are available from this pluton; however, the Moss Cape pluton to the east is a similar quartz diorite body intruding the Belkofski Formation; a potassium-argon age determination of 3.21 ± 0.14 Ma was obtained on biotite (DuBois and others, 1987) from this pluton. Another diorite body intruding the Belkofski Formation on Deer Island has a potassium-argon age of 6.00 ± 0.20 Ma on plagioclase (Hugh McLean, written commun., 1986). However, plagioclase is generally considered of marginal utility in dating of intrusive rocks, due to argon loss, hence the date on the Deer Island diorite may be considered a minimum age. The similar composition of these intrusive bodies and that they intrude the Belkofski Formation suggests a similar late Miocene or Pliocene age for the King Cove pluton. Plutons of similar age are well known on the Alaska Peninsula to the northeast (Wilson and others, 1983; F.H. Wilson and Nora Shew, unpublished data, 1989).

The King Cove pluton has greater areal extent than previously mapped. The contact of the quartz diorite pluton with the Belkofski Formation continues south of King Cove Lagoon along the high tide line of King Cove. Both the pluton and Belkofski Formation contain disseminated sulfides and minor sulfide-bearing quartz veins near their contact.

Mafic (andesitic?) dikes cut the Belkofski Formation but the exposures are not adequate to indicate whether the dikes cut the pluton. However, the mafic dikes are not mineralized as are the quartz diorite pluton and the Belkofski Formation, suggesting that the dikes are younger than the main intrusive phase.

Burk (1965) and McLean and others (1978) both show a number of volcanic intrusive centers within the Belkofski Formation on the ridges east and west of King Cove. Capping the ridge, east of King Cove is a volcanic unit mapped as a volcanic intrusive center by Burk (1965) and McLean and others (1978). However it does not appear to us to be an intrusive feature, but rather an eroded remnant of a volcanic flow. These volcanic rocks may be a part of the Belkofski Formation or equivalent to the Miocene volcanic rocks found to the east. Perhaps other volcanic intrusive centers in the King Cove area shown by Burk (1965) and McLean and others (1978) are this same type of volcanic unit. Other volcanic flows which occurred as valley fills but now, due to erosion and reversal of topography, form ridge crests, were mapped by Burk (1965) as volcanic intrusive centers elsewhere on the Alaska Peninsula (F.H. Wilson and others, unpublished data).

GEOCHEMICAL SAMPLING

Seventy-one rock samples from the King Cove vicinity were analyzed using 6-step semi-quantitative emission spectrography as described by Grimes and Marazino (1968), and by atomic absorption spectrophotometry (table 1). Sample data included coding for source, rock type, and other characteristics (table 2) to facilitate statistical analysis. The samples were analyzed for 35 elements (with lower level determination levels shown in table 3). No samples had detectable levels of gold (Au), antimony (Sb), germanium (Ge), thorium (Th), or tungsten (W) and these elements, though analyzed, are not shown in the table of analytical results (table 3).

Samples collected as part of this project were compared to thresholds calculated from data for 2708 samples from the adjacent Port Moller and Stepovak Bay quadrangles (Angeloni and others, 1985; Wilson and others, 1987). The ninetieth percentile for each metal for all rock types was used as the anomalous threshold for the rocks from King Cove. Sixteen samples had anomalous amounts of various metals (table 4). Eight of the anomalous samples are from near the contact between the pluton and the Belkofski Formation.

CONCLUSIONS

We have herein tentatively reassigned the bedrock exposures northwest of Indian Head to the Stepovak Formation of Burk (1965). This is the first recognition of rocks of this age and stratigraphic position in the Cold Bay quadrangle. The nearest other exposures of this formation occur 50 kilometers to the northwest.

The recognition herein of the Unga Formation on the west side of Deer Island indicates that the known outcrop area of the Belkofski Formation lies entirely within the Unga depositional basin. Nevertheless, we know of no place where the Belkofski and Unga are in contact. In all areas we have examined, the Belkofski Formation occurs in close proximity to large plutons, for example the King Cove pluton. Our preferred age for the Belkofski is the same as that for the Unga. We therefore suggest that the Belkofski Formation, though a distinct mappable unit, is hydrothermally altered and contact metamorphosed Unga Formation.

We have found the King Cove pluton to have a larger areal extent than has been previously shown. The nearly ubiquitous hydrothermal alteration, the development of sulfide veins, and occurrence of disseminated sulfides and low level Pb anomalies in its contact zone with the country rock indicates mineral potential.

REFERENCES

- Angeloni, L.M., Wilson, F.H., and Sutley, Stephen. 1985. Map and tables showing preliminary rock geochemical data, Port Moller, Stepovak Bay, and Simeonof Island quadrangles, Alaska: U.S. Geological Survey Open-File Report 85-470, about 70 p., scale 1:250,000, 1 map sheet.

- Burk, 1965, Geology of the Alaska Peninsula - Island arc and continental margin: Geological Society of America Memoir 99, pt.1, 250 p., scales 1:250,000 and 1:500,000, 3 sheets.
- Detterman, R.L., Case, J.E., Miller, J.W., Wilson, F.H., and Yount, M.E., 1989, Stratigraphic Framework of the Alaska Peninsula: U.S. Geological Survey Bulletin, in press.
- DuBois, G.D., Wilson, F.H., and Shew, N.B., 1987, Potassium-argon ages from the Port Moller and Stepovak Bay quadrangles, Alaska Peninsula: U.S. Geological Survey Open-File Report 87-191, 1 sheet, scale 1:250,000.
- Grimes, D.J., and Marranzino, A.P., 1968, Direct-current arc and alternating-current spark emission spectrographic field methods for the semiquantitative analysis of geologic materials: U.S. Geological Survey Circular 591, 6 p.
- Kennedy, G.C., and Waldron, H.H., 1947, Geology of Pavlof Volcano and vicinity, in U.S. Geological Survey Alaskan Volcano Investigations Report No. 2: Progress of investigations in 1946, p. 9-18.
- Kennedy, G.C., and Waldron, H.H., 1955, Geology of the Pavlof Volcano and vicinity, Alaska: U.S. Geological Survey Bulletin 1028-A, 19 p., scales 1:100,000.
- McLean, Hugh, 1979, Tertiary stratigraphy and petroleum potential of Cold Bay-False Pass area, Alaska Peninsula: American Association of Petroleum Geologists Bulletin, v. 63, no. 9, p. 1522-1526.
- McLean, Hugh, Engelhardt, C.L., and Howell, D.G., 1978, Reconnaissance geologic map of the Cold Bay and False Pass quadrangles, Alaska: U.S. Geological Survey Open-File Report 78-323, scale 1:250,000.
- Waldron, H.H., 1961, Geologic reconnaissance of Frosty Peak volcano and vicinity, Alaska: U.S. Geological Survey Bulletin 1028-T, p. 677-708.
- Wilson, F.H., Gaum, W.C., and Herzon, P.L., 1983, map and tables showing geochronology and whole-rock geochemistry of the Chignik and Sutwik Island quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1053-M, 3 sheets, scale 1:250,000.
- Wilson, F.H., Harris, E.E., and Bailey, E.A., 1987, Preliminary analytical results and sample locality map for rock samples collected in 1985 and 1986, Port Moller and Stepovak Bay quadrangles, Alaska: U.S. Geological Survey Open-File Report 87-497, 44 p., scale 1:250,000, 1 map sheet.

Table 1. Analytical results of rock geochemistry samples from King Cove, Alaska. Results are in ppm except for Fe, Mg, P, Ca, Ti, which are in percent. Analysts were R.T. Hopkins, Z.O. Brown and E.A. Bailey. Sample coding is shown in table 2, and the lower limits of determination for each element is shown in table 3. Elements analyzed by atomic absorption spectrophotometry methods are shown by adding -AA to the element symbols (i.e. Sb-AA, table 3).

| Sample number | Lat. N | Long. W | Sample type | Sample source | Rock class | Quad | FC3 | FC4 | Form | Rock type |
|---------------|--------|---------|-------------|---------------|------------|------|-----|-----|------|-----------|
| 88Dg1 | 550431 | 1621315 | A | A | D | A-1 | A | B | E | Q |
| 88Dg2a | 550444 | 1621227 | A | A | D | A-1 | A | A | B | M |
| 88Dg2b | 550444 | 1621227 | A | A | D | A-1 | A | B | B | M |
| 88Dg2c | 550444 | 1621227 | A | A | D | A-1 | A | B | E | Z |
| 88Dg3 | 550452 | 1621227 | A | A | D | A-1 | A | A | B | M |
| 88Dg4 | 550513 | 1621835 | A | A | D | A-1 | A | D | D | Q |
| 88Dg5 | 550510 | 1621838 | A | D | D | A-1 | A | D | D | Q |
| 88Dg6 | 550339 | 1621813 | A | A | D | A-1 | A | A | A | L |
| 88Dg7 | 550316 | 1621938 | A | A | D | A-1 | A | D | B | K |
| 88Dg7a | 550316 | 1621938 | A | A | D | A-1 | A | D | E | Q |
| 88Dg8 | 550306 | 1621943 | A | A | D | A-1 | A | B | D | N |
| 88Dg9 | 550259 | 1621946 | A | A | D | A-1 | A | B | E | Z |
| 88Dg9a | 550259 | 1621946 | A | A | D | A-1 | A | D | D | Q |
| 88Dg10 | 550250 | 1621946 | A | A | D | A-1 | A | B | E | L |
| 88Dg11 | 550337 | 1621807 | A | A | D | A-1 | A | A | F | L |
| 88Dg12 | 550335 | 1621756 | A | A | D | A-1 | A | A | F | L |
| 88Dg13 | 550332 | 1621750 | A | D | D | A-1 | A | D | F | L |
| 88Dg14 | 550332 | 1621747 | A | A | D | A-1 | A | D | B | M |
| 88Dg15 | 550337 | 1621744 | A | A | D | A-1 | A | A | B | Z |
| 88Dg16 | 550325 | 1621749 | A | A | D | A-1 | A | A | B | Z |
| 88Dg17 | 550316 | 1621756 | A | A | Y | A-1 | A | B | G | Z |
| 88Dt1 | 550601 | 1621510 | B | A | D | A-1 | A | A | A | K |
| 88Dt2 | 550558 | 1621531 | A | A | D | A-1 | A | A | A | K |
| 88Dt3 | 550602 | 1621455 | A | A | D | A-1 | A | A | A | M |
| 88Dt4 | 550606 | 1621430 | A | A | D | A-1 | A | A | A | M |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Lat. N | Long. W | Sample type | Sample source | Rock class | Quad | FC3 | FC4 | Form | Rock type |
|---------------|--------|---------|-------------|---------------|------------|------|-----|-----|------|-----------|
| 88Dt5 | 550615 | 1621341 | B | A | B | A-1 | A | A | A | C |
| 88Dt6 | 550603 | 1621331 | B | A | B | A-1 | A | A | A | A |
| 88Dt7 | 550548 | 1621305 | B | A | B | A-1 | A | A | A | B |
| 88Dt9a | 550312 | 1621821 | A | A | C | A-1 | A | D | A | Z |
| 88Dt9b | 550312 | 1621821 | A | A | D | A-1 | A | B | D | Q |
| 88Dt10c | 550238 | 1621805 | A | A | Y | A-1 | A | D | G | Z |
| 88Dt11 | 550238 | 1621805 | A | A | D | A-1 | A | A | D | Q |
| 88Dt12 | 550221 | 1621800 | A | A | D | A-1 | A | D | D | Z |
| 88Dt15 | 545506 | 1622515 | A | A | D | D-2 | A | A | A | Z |
| 88Dt16 | 545448 | 1622530 | A | A | B | D-2 | A | A | A | K |
| 88Dt17 | 545433 | 1622521 | A | A | B | D-2 | A | A | A | Z |
| 88Dt19 | 545348 | 1622459 | B | A | B | D-2 | A | A | A | C |
| 88Dt20 | 545828 | 1621645 | A | A | D | D-1 | A | A | A | K |
| 88Dt21 | 545837 | 1621649 | A | A | D | D-1 | A | A | D | Q |
| 88Ws1a | 550423 | 1621335 | A | A | C | A-1 | A | A | A | M |
| 88Ws1b | 550423 | 1621335 | A | A | D | A-1 | A | B | E | Q |
| 88Ws2a | 550426 | 1621326 | A | A | D | A-1 | A | A | F | M |
| 88Ws2b | 550426 | 1621326 | A | A | D | A-1 | A | B | F | M |
| 88Ws4a | 550520 | 1621201 | A | A | D | A-1 | A | A | F | L |
| 88Ws4b | 550520 | 1621201 | A | A | Y | A-1 | A | C | G | Z |
| 88Ws5a | 550514 | 1621840 | A | A | C | A-1 | A | A | A | Z |
| 88Ws5b | 550514 | 1621840 | A | A | D | A-1 | A | D | D | Q |
| 88Ws6a | 550511 | 1621840 | A | A | D | A-1 | A | A | D | Q |
| 88Ws6b | 550511 | 1621840 | A | A | D | A-1 | A | D | D | Q |
| 88Ws7 | 550654 | 1621739 | A | A | D | A-1 | A | A | F | M |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Lat. N | Long. W | Sample type | Sample source | Rock class | Quad | FC3 | FC4 | Form | Rock type |
|---------------|--------|---------|-------------|---------------|------------|------|-----|-----|------|-----------|
| 88Ws8a | 550654 | 1621730 | A | A | C | A-1 | A | B | A | Z |
| 88Ws8b | 550654 | 1621730 | A | A | D | A-1 | A | A | B | Q |
| 88Ws8c | 550654 | 1621730 | A | A | D | A-1 | A | B | F | L |
| 88Ws9 | 550654 | 1621705 | A | A | D | A-1 | A | A | F | L |
| 88Ws10a | 550647 | 1621654 | A | A | D | A-1 | A | A | D | N |
| 88Ws10b | 550647 | 1621654 | A | A | C | A-1 | D | A | A | Z |
| 88Ws10c | 550647 | 1621654 | A | A | C | A-1 | A | C | A | Z |
| 88Ws11 | 550640 | 1621708 | A | A | D | A-1 | A | A | A | K |
| 88Ws11b | 550640 | 1621708 | A | A | D | A-1 | A | A | A | K |
| 88Ws12 | 550413 | 1621813 | A | A | D | A-1 | A | A | D | N |
| 88Ws13 | 550402 | 1621937 | A | A | D | A-1 | A | D | E | L |
| 88Ws14 | 550359 | 1621938 | A | A | D | A-1 | A | A | D | N |
| 88Ws15 | 550353 | 1621939 | A | A | D | A-1 | A | A | D | N |
| 88Ws16 | 550350 | 1621938 | A | A | D | A-1 | A | A | D | N |
| 88Ws17 | 545522 | 1622519 | A | A | D | D-2 | A | A | B | L |
| 88Ws18 | 545453 | 1622527 | A | A | D | D-2 | A | C | E | J |
| 88Ws19 | 545712 | 1622510 | A | A | D | D-2 | A | A | B | J |
| 88Ws20a | 545509 | 1621529 | A | A | D | D-1 | A | B | B | L |
| 88Ws20b | 545509 | 1621529 | A | A | D | D-1 | A | B | F | L |
| 88Ws21a | 545436 | 1625040 | A | A | D | D-1 | A | A | F | K |
| 88Ws21b | 545436 | 1625040 | A | A | Y | D-1 | A | D | H | Z |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Fe | Mg | Na | P | Ca | Tl | Ag | As | As-AA | B | Ba | Be | Bi |
|---------------|------|------|------|-------|------|------|----|----|--------|--------|--------|-------|----|
| 88Dg1a | 2.00 | 0.70 | 2.00 | N | 1.00 | 0.30 | N | N | 60.00 | <10.00 | 700.00 | 1.00 | N |
| 88Dg2a | 3.00 | 1.50 | 2.00 | N | 1.00 | 0.50 | N | N | 10.00 | <10.00 | 500.00 | <1.00 | N |
| 88Dg2b | 5.00 | 1.00 | 2.00 | <0.20 | 0.70 | 0.50 | N | N | 10.00 | N | 500.00 | 1.00 | N |
| 88Dg2c | 5.00 | 1.50 | 2.00 | N | 2.00 | 0.50 | N | N | <10.00 | N | 500.00 | <1.00 | N |
| 88Dg3 | 3.00 | 1.00 | 2.00 | <0.20 | 1.50 | 0.50 | N | N | <10.00 | 10.00 | 500.00 | 1.00 | N |
| 88Dg4 | 5.00 | 1.50 | 1.50 | N | 2.00 | 0.50 | N | N | <10.00 | 50.00 | 500.00 | <1.00 | N |
| 88Dg5 | 5.00 | 1.50 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | 20.00 | 700.00 | <1.00 | N |
| 88Dg6 | 5.00 | 1.50 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | N | 150.00 | <1.00 | N |
| 88Dg7 | 5.00 | 1.50 | 2.00 | <0.20 | 2.00 | 0.50 | N | N | <10.00 | <10.00 | 200.00 | <1.00 | N |
| 88Dg7a | 5.00 | 1.50 | 1.50 | N | 3.00 | 0.50 | N | N | <10.00 | N | 300.00 | <1.00 | N |
| 88Dg8 | 3.00 | 1.50 | 1.50 | N | 3.00 | 0.50 | N | N | <10.00 | N | 700.00 | <1.00 | N |
| 88Dg9 | 3.00 | 1.50 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | <10.00 | 700.00 | <1.00 | N |
| 88Dg9a | 5.00 | 2.00 | 1.50 | N | 5.00 | 0.50 | N | N | <10.00 | N | 30.00 | N | N |
| 88Dg10 | 5.00 | 1.50 | 2.00 | N | 2.00 | 0.50 | N | N | <10.00 | N | 500.00 | N | N |
| 88Dg11 | 3.00 | 1.50 | 2.00 | N | 2.00 | 0.30 | N | N | <10.00 | <10.00 | 300.00 | <1.00 | N |
| 88Dg12 | 5.00 | 2.00 | 1.50 | N | 3.00 | 0.50 | N | N | <10.00 | <10.00 | 300.00 | <1.00 | N |
| 88Dg13 | 3.00 | 1.50 | 2.00 | N | 1.50 | 0.30 | N | N | <10.00 | 10.00 | 500.00 | <1.00 | N |
| 88Dg14 | 5.00 | 1.50 | 2.00 | N | 2.00 | 0.50 | N | N | <10.00 | 15.00 | 700.00 | <1.00 | N |
| 88Dg15 | 3.00 | 1.50 | 2.00 | N | 2.00 | 0.30 | N | N | <10.00 | N | 500.00 | <1.00 | N |
| 88Dg16 | 5.00 | 1.50 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | N | 300.00 | <1.00 | N |
| 88Dg17 | 3.00 | 0.70 | 1.50 | N | 3.00 | 0.50 | N | N | <10.00 | <10.00 | 300.00 | <1.00 | N |
| 88Dt1 | 5.00 | 1.00 | 2.00 | N | 1.50 | 0.50 | N | N | <10.00 | N | 200.00 | <1.00 | N |
| 88Dt2 | 7.00 | 2.00 | 3.00 | N | 2.00 | 0.50 | N | N | <10.00 | N | 150.00 | N | N |
| 88Dt3 | 5.00 | 1.00 | 1.50 | N | 3.00 | 0.70 | N | N | <10.00 | N | 300.00 | 1.50 | N |
| 88Dt4 | 5.00 | 1.50 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | <10.00 | 500.00 | 1.50 | N |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Fe | Mg | Na | P | Ca | Ti | Ag | As | As-AA | B | Ba | Be | Bi |
|---------------|------|-------|-------|-------|-------|------|------|----|--------|--------|--------|-------|-------|
| 88D15 | 7.00 | 2.00 | 3.00 | N | 3.00 | 1.00 | N | N | <10.00 | N | 100.00 | N | N |
| 88D16 | 5.00 | 1.00 | 2.00 | <0.20 | 1.50 | 0.50 | N | N | <10.00 | <10.00 | 500.00 | 1.00 | N |
| 88D17 | 5.00 | 1.50 | 2.00 | N | 1.00 | 0.30 | N | N | <10.00 | 10.00 | 700.00 | 1.00 | N |
| 88L19a | 5.00 | 2.00 | 1.50 | N | 1.50 | 0.5 | N | N | <10.00 | N | 300.00 | <1.00 | N |
| 88D19b | 5.00 | 1.50 | 2.00 | N | 5.00 | 0.5 | N | N | <10.00 | <10.00 | 200.00 | <1.00 | N |
| 88D10c | 7.00 | 2.00 | 2.00 | N | 5.00 | 0.7 | 0.30 | N | <10.00 | 10.00 | 300.00 | <1.00 | 10.00 |
| 88D11 | 5.00 | 1.50 | 2.00 | N | 5.00 | 0.50 | N | N | <10.00 | N | 300.00 | <1.00 | N |
| 88D12 | 5.00 | 0.70 | 1.00 | N | 3.00 | 0.50 | N | N | <10.00 | 20.00 | 150.00 | 1.50 | N |
| 88D15 | 5.00 | 2.00 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | N | 200.00 | <1.00 | N |
| 88D16 | 7.00 | 10.00 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | N | 150.00 | N | N |
| 88D17 | 7.00 | 1.50 | 3.00 | N | 3.00 | 1.00 | N | N | <10.00 | 10.00 | 150.00 | <1.00 | N |
| 88D19 | 7.00 | 1.50 | 3.00 | <0.20 | 3.00 | 0.70 | N | N | <10.00 | N | 150.00 | <1.00 | N |
| 88D20 | 3.00 | 1.00 | 2.00 | <0.20 | 1.00 | 0.50 | N | N | <10.00 | N | 200.00 | 1.50 | N |
| 88D21 | 7.00 | 2.00 | 1.50 | N | 7.00 | 0.30 | N | N | <10.00 | N | 200.00 | <1.00 | N |
| 88Ws1a | 5.00 | 0.15 | N | N | 7.00 | 0.30 | N | N | 60.00 | 10.00 | <20.00 | 2.00 | N |
| 88Ws1b | 5.00 | 1.50 | 1.50 | N | 2.00 | 0.30 | N | N | <10.00 | N | 500.00 | N | N |
| 88Ws2a | 7.00 | 1.50 | 3.00 | <0.20 | 2.00 | 1.00 | N | N | <10.00 | <10.00 | 500.00 | <1.00 | N |
| 88Ws2b | 3.00 | 0.50 | 1.00 | N | 3.00 | 0.30 | N | N | <10.00 | <10.00 | 70.00 | 1.00 | N |
| 88Ws4a | 7.00 | 1.50 | 3.00 | N | 1.50 | 0.50 | N | N | <10.00 | 10.00 | 500.00 | <1.00 | N |
| 88Ws4b | 0.70 | 0.30 | <0.20 | N | 15.00 | 0.07 | N | N | <10.00 | 10.00 | 300.00 | <1.00 | N |
| 88Ws5a | 5.00 | 2.00 | 1.00 | N | 7.00 | 0.07 | N | N | 20.00 | <10.00 | 500.00 | <1.00 | N |
| 88Ws5b | 5.00 | 1.50 | 1.50 | N | 3.00 | 0.70 | N | N | <10.00 | 10.00 | 500.00 | <1.00 | N |
| 88Ws6a | 5.00 | 1.50 | 2.00 | N | 3.00 | 0.70 | N | N | <10.00 | 10.00 | 500.00 | <1.00 | N |
| 88Ws6b | 3.00 | 1.50 | 1.50 | N | 3.00 | 0.30 | N | N | <10.00 | <10.00 | 500.00 | 1.00 | N |
| 88Ws7 | 7.00 | 1.50 | 5.00 | N | 0.70 | 0.70 | N | N | <10.00 | <10.00 | 70.00 | <1.00 | N |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Fe | Mg | Na | P | Ca | Ti | Ag | As | As-AA | B | Ba | Be | Bi |
|---------------|------|------|------|-------|------|------|----|----|--------|--------|---------|-------|----|
| 88Ws8a | 7.00 | 0.70 | 1.50 | N | 1.50 | 1.00 | N | N | <10.00 | 10.00 | 300.00 | <1.00 | N |
| 88Ws8b | 7.00 | 2.00 | 2.00 | N | 7.00 | 0.50 | N | N | <10.00 | <10.00 | 200.00 | N | N |
| 88Ws8c | 5.00 | 2.00 | 1.50 | N | 7.00 | 0.50 | N | N | <10.00 | <10.00 | 200.00 | N | N |
| 88Ws9 | 5.00 | 2.00 | 2.00 | N | 7.00 | 0.50 | N | N | <10.00 | <10.00 | 300.00 | N | N |
| 88Ws10a | 7.00 | 1.50 | 1.50 | N | 5.00 | 0.70 | N | N | <10.00 | <10.00 | 150.00 | N | N |
| 88Ws10b | 2.00 | 0.70 | 1.50 | N | 0.50 | 0.70 | N | N | <10.00 | 15.00 | 500.00 | 1.50 | N |
| 88Ws10c | 5.00 | 0.70 | 1.50 | <0.20 | 1.50 | 0.50 | N | N | <10.00 | <10.00 | 500.00 | 1.00 | N |
| 88Ws11 | 7.00 | 1.50 | 2.00 | N | 1.50 | 0.70 | N | N | <10.00 | 20.00 | 70.00 | <1.00 | N |
| 88Ws11b | 7.00 | 1.50 | 2.00 | N | 0.30 | 0.70 | N | N | 40.00 | 10.00 | 50.00 | <1.00 | N |
| 88Ws12 | 5.00 | 1.50 | 2.00 | N | 3.00 | 0.50 | N | N | <10.00 | 10.00 | 700.00 | <1.00 | N |
| 88Ws13 | 7.00 | 2.00 | 1.50 | N | 5.00 | 0.50 | N | N | 10.00 | <10.00 | 200.00 | <1.00 | N |
| 88Ws14 | 7.00 | 1.50 | 2.00 | N | 7.0 | 0.50 | N | N | <10.00 | N | 300.00 | <1.00 | N |
| 88Ws15 | 7.00 | 2.00 | 2.00 | N | 5.00 | 0.70 | N | N | 10.00 | <10.00 | 300.00 | N | N |
| 88Ws16 | 5.00 | 2.00 | 2.00 | N | 5.00 | 0.50 | N | N | <10.00 | <10.00 | 300.00 | N | N |
| 88Ws17 | 5.00 | 1.50 | 3.00 | N | 2.00 | 0.70 | N | N | <10.00 | <10.00 | 300.00 | <1.00 | N |
| 88Ws18 | 5.00 | 2.00 | 3.00 | N | 5.00 | 0.70 | N | N | <10.00 | N | 500.00 | <1.00 | N |
| 88Ws19 | 5.00 | 2.00 | 2.00 | <0.20 | 5.00 | 0.70 | N | N | <10.00 | <10.00 | 300.00 | <1.00 | N |
| 88Ws20a | 5.00 | 1.50 | 2.00 | <0.20 | 3.00 | 0.70 | N | N | <10.00 | <10.00 | 1000.00 | <1.00 | N |
| 88Ws20b | 5.00 | 3.00 | 2.00 | <0.20 | 7.00 | 0.50 | N | N | <10.00 | N | 150.00 | <1.00 | N |
| 88Ws21a | 5.00 | 1.50 | 2.00 | <0.20 | 1.50 | 0.50 | N | N | 10.00 | <10.00 | 500.00 | <1.00 | N |
| 88Ws21b | 3.00 | 0.30 | 2.00 | <0.20 | 0.30 | 0.50 | N | N | 30.00 | <10.00 | 300.00 | 1.50 | N |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Bi-AA | Cd | Cd-AA | Co | Cr | Cu | Ga | La | Mn | Mo | Nb | Ni |
|---------------|-------|----|-------|--------|--------|-------|-------|--------|---------|----|--------|-------|
| 88Dg1a | 1.00 | N | 1.00 | <10.00 | <10.00 | 20.00 | 20.00 | <50.00 | 700.00 | N | N | <5.00 |
| 88Dg2a | <1.00 | N | <0.10 | 10.00 | 50.00 | 20.00 | 20.00 | <50.00 | 1000.00 | N | N | 7.00 |
| 88Dg2b | <1.00 | N | <0.10 | 15.00 | 70.00 | 20.00 | 20.00 | <50.00 | 1000.00 | N | N | 7.00 |
| 88Dg2c | 1.00 | N | <0.10 | 20.00 | 50.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg3 | <1.00 | N | 0.10 | 20.00 | 50.00 | 50.00 | 30.00 | <50.00 | 1000.00 | N | N | 15.00 |
| 88Dg4 | <1.00 | N | 0.10 | 20.00 | 30.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg5 | 1.00 | N | 0.10 | 15.00 | 15.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg6 | 1.00 | N | <0.10 | 20.00 | 20.00 | 30.00 | 30.00 | <50.00 | 1500.00 | N | N | <5.00 |
| 88Dg7 | <1.00 | N | <0.10 | 15.00 | 30.00 | 70.00 | 20.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg7a | <1.00 | N | <0.10 | 15.00 | 50.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg8 | <1.00 | N | 0.20 | 15.00 | N | 20.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg9 | <1.00 | N | <0.10 | 20.00 | 50.00 | 20.00 | 30.00 | <50.00 | 1000.00 | N | N | 7.00 |
| 88Dg9a | <1.00 | N | <0.10 | 50.00 | 100.00 | 70.00 | 20.00 | N | 1000.00 | N | N | 20.00 |
| 88Dg10 | <1.00 | N | <0.10 | 20.00 | 10.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg11 | <1.00 | N | <0.10 | 20.00 | 10.00 | 20.00 | 20.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg12 | 1.00 | N | <0.10 | 30.00 | 100.00 | 50.00 | 20.00 | <50.00 | 1500.00 | N | N | 15.00 |
| 88Dg13 | <1.00 | N | <0.10 | 20.00 | 30.00 | <5.00 | 20.00 | <50.00 | 1000.00 | N | N | 5.00 |
| 88Dg14 | <1.00 | N | 0.10 | 20.00 | 30.00 | 5.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg15 | 1.00 | N | 0.10 | 15.00 | 10.00 | 20.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg16 | 1.00 | N | 0.10 | 20.00 | 30.00 | 30.00 | 20.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Dg17 | 1.00 | N | <0.10 | 10.00 | 20.00 | 30.00 | 20.00 | <50.00 | 500.00 | N | N | <5.00 |
| 88Dt1 | <1.00 | N | <0.10 | 15.00 | 20.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | <20.00 | <5.00 |
| 88Dt2 | <1.00 | N | <0.10 | 20.00 | 50.00 | 50.00 | 30.00 | N | 1000.00 | N | N | 5.00 |
| 88Dt3 | <1.00 | N | <0.10 | 10.00 | 50.00 | 20.00 | 30.00 | <50.00 | 1000.00 | N | <20.00 | <5.00 |
| 88Dt4 | <1.00 | N | 0.10 | 10.00 | 15.00 | 30.00 | 50.00 | <50.00 | 1000.00 | N | <20.00 | <5.00 |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Bi-AA | Cd | Cd-AA | Co | Cr | Cu | Ga | La | Mn | Mo | Nb | Ni |
|---------------|-------|----|-------|--------|---------|--------|-------|--------|---------|-------|--------|--------|
| 88Dt5 | <1.00 | N | <0.10 | 30.00 | <10.00 | 70.00 | 30.00 | <50.00 | 1500.00 | N | <20.00 | 5.00 |
| 88Dt6 | <1.00 | N | <0.10 | 15.00 | 50.00 | 15.00 | 30.00 | <50.00 | 1000.00 | N | <20.00 | 7.00 |
| 88Dt7 | <1.00 | N | <0.10 | 10.00 | 50.00 | 20.00 | 30.00 | N | 700.00 | N | N | 10.00 |
| 88Dt9a | <1.00 | N | 0.10 | 15.00 | 50.00 | 100.00 | 30.00 | N | 1500.00 | N | N | 10.00 |
| 88Dt9b | <1.00 | N | 0.10 | 20.00 | 30.00 | 30.00 | 50.00 | N | 1500.00 | N | N | <5.00 |
| 88Dt10c | 1.00 | N | 0.90 | 20.00 | 70.00 | 7.00 | 30.00 | N | 1000.00 | N | N | 20.00 |
| 88Dt11 | <1.00 | N | <0.10 | 15.00 | <10.00 | 30.00 | 30.00 | N | 1500.00 | N | N | <5.00 |
| 88Dt12 | 1.00 | N | 0.10 | 10.00 | 70.00 | 30.00 | 50.00 | <50.00 | 300.00 | N | <20.00 | 5.00 |
| 88Dt15 | <1.00 | N | <0.10 | 20.00 | 150.00 | 30.00 | 20.00 | <50.00 | 1000.00 | N | N | 20.00 |
| 88Dt16 | <1.00 | N | <0.10 | 30.00 | 1000.00 | 50.00 | 20.00 | N | 1000.00 | N | N | 150.00 |
| 88Dt17 | <1.00 | N | 0.20 | 20.00 | 30.00 | 50.00 | 30.00 | N | 1000.00 | N | <20.00 | 15.00 |
| 88Dt19 | <1.00 | N | 0.20 | 20.00 | 30.00 | 50.00 | 30.00 | N | 1000.00 | N | <20.00 | <5.00 |
| 88Dt20 | <1.00 | N | 0.10 | <10.00 | 20.00 | 20.00 | 30.00 | <50.00 | 1000.00 | N | <20.00 | N |
| 88Dt21 | <1.00 | N | <0.10 | 30.00 | 150.00 | 50.00 | 20.00 | <50.00 | 1000.00 | N | N | 50.00 |
| 88Ws1a | 3.00 | N | 1.80 | N | <10.00 | 10.00 | 50.00 | N | 2000.00 | <5.00 | N | 5.00 |
| 88Ws1b | 2.00 | N | 0.20 | 30.00 | 70.00 | 50.00 | 30.00 | N | 700.00 | N | N | 20.00 |
| 88Ws2a | <1.00 | N | 0.20 | 30.00 | 15.00 | 30.00 | 50.00 | N | 1000.00 | N | N | 15.00 |
| 88Ws2b | 1.00 | N | 2.10 | 10.00 | 30.00 | 50.00 | 20.00 | <50.00 | 1500.00 | 15.00 | N | 7.00 |
| 88Ws4a | 2.00 | N | 0.20 | 20.00 | 100.00 | 20.00 | 50.00 | N | 700.00 | N | N | 20.00 |
| 88Ws4b | <1.00 | N | <0.10 | N | 15.00 | 20.00 | N | <50.00 | 2000.00 | N | N | <5.00 |
| 88Ws5a | 1.00 | N | 0.40 | 30.00 | 70.00 | 50.00 | 20.00 | <50.00 | 1000.00 | N | N | 20.00 |
| 88Ws5b | 1.00 | N | 0.10 | 20.00 | 20.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Ws6a | 2.00 | N | 0.10 | 20.00 | 30.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | 5.00 |
| 88Ws6b | <1.00 | N | <0.10 | 15.00 | 30.00 | 30.00 | 30.00 | <50.00 | 700.00 | N | N | <5.00 |
| 88Ws7 | 3.00 | N | 0.30 | 30.00 | 50.00 | 50.00 | 50.00 | <50.00 | 1000.00 | N | N | 15.00 |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Bi-AA | Cd | Cd-AA | Co | Cr | Cu | Ga | La | Mn | Mo | Nb | Ni |
|---------------|-------|----|-------|--------|--------|-------|-------|--------|---------|----|--------|-------|
| 88Ws8a | 2.00 | N | 0.20 | 30.00 | 100.00 | 70.00 | 30.00 | <50.00 | 1000.00 | N | N | 20.00 |
| 88Ws8b | <1.00 | N | 0.10 | 30.00 | 20.00 | 30.00 | 50.00 | N | 1000.00 | N | N | 15.00 |
| 88Ws8c | 2.00 | N | <0.10 | 30.00 | 50.00 | 30.00 | 30.00 | N | 1000.00 | N | N | 15.00 |
| 88Ws9 | 2.00 | N | <0.10 | 30.00 | 100.00 | 30.00 | 30.00 | N | 1000.00 | N | N | 20.00 |
| 88Ws10a | 3.00 | N | 0.10 | 30.00 | 30.00 | 30.00 | 20.00 | N | 1000.00 | N | N | 7.00 |
| 88Ws10b | 1.00 | N | 0.10 | 15.00 | 30.00 | 30.00 | 30.00 | <50.00 | 700.00 | N | N | 7.00 |
| 88Ws10c | 1.00 | N | 0.10 | 15.00 | 30.00 | 20.00 | 30.00 | <50.00 | 1000.00 | N | N | 5.00 |
| 88Ws11 | <1.00 | N | 0.10 | 20.00 | 50.00 | 30.00 | 30.00 | N | 1000.00 | N | N | 15.00 |
| 88Ws11b | <1.00 | N | <0.10 | 30.00 | 70.00 | 50.00 | 30.00 | N | 1000.00 | N | N | 20.00 |
| 88Ws12 | 1.00 | N | <0.10 | 20.00 | 15.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | <5.00 |
| 88Ws13 | 2.00 | N | 0.40 | 30.00 | 70.00 | 30.00 | 20.00 | <50.00 | 1500.00 | N | N | 10.00 |
| 88Ws14 | 2.00 | N | 0.10 | 20.00 | N | 50.00 | 30.00 | <50.00 | 700.00 | N | N | <5.00 |
| 88Ws15 | <1.00 | N | 0.10 | 30.00 | 70.00 | 50.00 | 30.00 | N | 1000.00 | N | N | 10.00 |
| 88Ws16 | <1.00 | N | <0.10 | 30.00 | 50.00 | 30.00 | 30.00 | N | 1000.00 | N | N | 7.00 |
| 88Ws17 | 1.00 | N | 0.10 | 20.00 | <10.00 | 30.00 | 50.00 | <50.00 | 700.00 | N | N | N |
| 88Ws18 | <1.00 | N | 0.10 | 30.00 | 100.00 | 50.00 | 30.00 | <50.00 | 1000.00 | N | N | 30.00 |
| 88Ws19 | 1.00 | N | <0.10 | 30.00 | 100.00 | 30.00 | 30.00 | <50.00 | 1000.00 | N | N | 50.00 |
| 88Ws20a | 2.00 | N | 0.10 | 20.00 | 100.00 | 30.00 | 30.00 | <50.00 | 700.00 | N | N | 10.00 |
| 88Ws20b | <1.00 | N | 0.10 | 30.00 | 150.00 | 50.00 | 50.00 | <50.00 | 1000.00 | N | N | 30.00 |
| 88Ws21a | <1.00 | N | 0.10 | 20.00 | 30.00 | 20.00 | 30.00 | <50.00 | 1000.00 | N | N | 5.00 |
| 88Ws21b | <1.00 | N | 0.20 | <10.00 | <10.00 | 7.00 | 30.00 | <50.00 | 500.00 | N | <20.00 | N |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Pb | Sb-AA | Sc | Sn | Sr | V | Y | Zn | Zn-AA | Zr |
|---------------|-------|-------|-------|-------|--------|--------|-------|---------|-------|--------|
| 88Dg1a | 20.00 | <2.00 | 5.00 | N | 300.00 | 70.00 | 20.00 | N | 85.00 | 150.00 |
| 88Dg2a | 30.00 | <2.00 | 15.00 | N | 300.00 | 100.00 | 20.00 | <200.00 | 95.00 | 150.00 |
| 88Dg2b | 15.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 20.00 | <200.00 | 90.00 | 150.00 |
| 88Dg2c | 20.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | N | 80.00 | 150.00 |
| 88Dg3 | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 90.00 | |
| 88Dg4 | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | N | 40.00 | 150.00 |
| 88Dg5 | 30.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 75.00 | 150.00 |
| 88Dg6 | 30.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 40.00 | 150.00 |
| 88Dg7 | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 75.00 | 150.00 |
| 88Dg7a | 20.00 | <2.00 | 20.00 | 20.00 | 300.00 | 150.00 | 20.00 | <200.00 | 45.00 | 100.00 |
| 88Dg8 | 20.00 | <2.00 | 15.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 50.00 | 100.00 |
| 88Dg9 | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 35.00 | 150.00 |
| 88Dg9a | 20.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | <200.00 | 30.00 | 70.00 |
| 88Dg10 | 20.00 | <2.00 | 20.00 | N | 700.00 | 200.00 | 30.00 | N | 55.00 | 100.00 |
| 88Dg11 | 20.00 | <2.00 | 15.00 | N | 700.00 | 150.00 | 30.00 | <200.00 | 30.00 | 100.00 |
| 88Dg12 | 15.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 20.00 | N | 35.00 | 70.00 |
| 88Dg13 | 20.00 | <2.00 | 15.00 | 50.00 | 300.00 | 150.00 | 20.00 | <200.00 | 60.00 | 100.00 |
| 88Dg14 | 20.00 | <2.00 | 15.00 | N | 300.00 | 150.00 | 30.00 | N | 80.00 | 150.00 |
| 88Dg15 | 30.00 | <2.00 | 15.00 | N | 300.00 | 150.00 | 30.00 | N | 50.00 | 150.00 |
| 88Dg16 | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | N | 25.00 | 150.00 |
| 88Dg17 | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 35.00 | 150.00 |
| 88Dt1 | 15.00 | <2.00 | 15.00 | N | 300.00 | 100.00 | 20.00 | N | 30.00 | 150.00 |
| 88Dt2 | 15.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 20.00 | N | 80.00 | 70.00 |
| 88Dt3 | 20.00 | <2.00 | 15.00 | N | 500.00 | 100.00 | 30.00 | N | 80.00 | 200.00 |
| 88Dt4 | 20.00 | <2.00 | 15.00 | N | 500.00 | 100.00 | 30.00 | N | 20.00 | 200.00 |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Pb | Sb-AA | Sc | Sn | Sr | V | Y | Zn | Zn-AA | Zr |
|---------------|--------|-------|-------|----|---------|--------|-------|---------|--------|--------|
| 88Dt5 | <10.00 | <2.00 | 20.00 | N | N | 300.00 | 20.00 | <200.00 | 85.00 | 100.00 |
| 88Dt6 | 20.00 | <2.00 | 15.00 | N | <100.00 | 150.00 | 20.00 | N | 80.00 | 100.00 |
| 88Dt7 | 20.00 | <2.00 | 15.00 | N | 200.00 | 100.00 | 20.00 | N | 70.00 | 70.00 |
| 88Dt9a | 15.00 | <2.00 | 15.00 | N | 300.00 | 150.00 | 20.00 | N | 30.00 | 70.00 |
| 88Dt9b | 15.00 | <2.00 | 15.00 | N | 500.00 | 200.00 | 20.00 | N | 40.00 | 150.00 |
| 88Dt10c | 200.00 | <2.00 | 20.00 | N | 200.00 | 150.00 | 20.00 | <200.00 | 110.00 | 100.00 |
| 88Dt11 | 50.00 | <2.00 | 15.00 | N | 300.00 | 200.00 | 20.00 | <200.00 | 65.00 | 100.00 |
| 88Dt12 | 10.00 | <2.00 | 15.00 | N | 300.00 | 70.00 | 30.00 | N | 5.00 | 150.00 |
| 88Dt15 | 15.00 | <2.00 | 20.00 | N | 500.00 | 150.00 | 20.00 | N | 60.00 | 100.00 |
| 88Dt16 | 10.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 15.00 | N | 50.00 | 70.00 |
| 88Dt17 | 15.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | N | 75.00 | 100.00 |
| 88Dt19 | 15.00 | <2.00 | 20.00 | N | 300.00 | 200.00 | 30.00 | N | 80.00 | 100.00 |
| 88Dt20 | 20.00 | <2.00 | 15.00 | N | 300.00 | 100.00 | 30.00 | N | 70.00 | 150.00 |
| 88Dt21 | 20.00 | <2.00 | 20.00 | N | 700.00 | 150.00 | 15.00 | N | 30.00 | 70.00 |
| 88Ws1a | 200.00 | 2.00 | 15.00 | N | 700.00 | 500.00 | 15.00 | <200.00 | 140.00 | 70.00 |
| 88Ws1b | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 20.00 | N | 55.00 | 150.00 |
| 88Ws2a | 20.00 | 2.00 | 20.00 | N | 500.00 | 150.00 | 30.00 | N | 90.00 | 150.00 |
| 88Ws2b | 50.00 | 2.00 | 15.00 | N | 300.00 | 150.00 | 30.00 | N | 60.00 | 200.00 |
| 88Ws4a | 30.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | <200.00 | 110.00 | 200.00 |
| 88Ws4b | 15.00 | <2.00 | <5.00 | N | 500.00 | 30.00 | 15.00 | N | 5.00 | 15.00 |
| 88Ws5a | 30.00 | 2.00 | 30.00 | N | 700.00 | 200.00 | 20.00 | <200.00 | 130.00 | 150.00 |
| 88Ws5b | 30.00 | <2.00 | 20.00 | N | 500.00 | 150.00 | 30.00 | N | 50.00 | 200.00 |
| 88Ws6a | 20.00 | 2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | N | 30.00 | 200.00 |
| 88Ws6b | 30.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 20.00 | N | 30.00 | 150.00 |
| 88Ws7 | 20.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | <200.00 | 95.00 | 150.00 |

Table 1 (cont.). Analytical results of rock geochemistry samples from King Cove, Alaska.

| Sample number | Pb | Sb-AA | Sc | Sn | Sr | V | Y | Zn | Zn-AA | Zr |
|---------------|-------|-------|-------|----|--------|--------|-------|---------|-------|--------|
| 88Ws8a | 20.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | <200.00 | 50.00 | 150.00 |
| 88Ws8b | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 20.00 | N | 55.00 | 70.00 |
| 88Ws8c | 15.00 | <2.00 | 30.00 | N | 300.00 | 200.00 | 30.00 | N | 45.00 | 100.00 |
| 88Ws9 | 20.00 | <2.00 | 30.00 | N | 500.00 | 200.00 | 30.00 | N | 25.00 | 150.00 |
| 88Ws10a | 15.00 | <2.00 | 20.00 | N | 300.00 | 200.00 | 30.00 | N | 35.00 | 100.00 |
| 88Ws10b | 30.00 | <2.00 | 20.00 | N | 200.00 | 200.00 | 30.00 | N | 20.00 | 150.00 |
| 88Ws10c | 20.00 | <2.00 | 15.00 | N | 500.00 | 100.00 | 30.00 | N | 70.00 | 200.00 |
| 88Ws11 | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | N | 70.00 | 100.00 |
| 88Ws11b | 15.00 | 4.00 | 20.00 | N | 200.00 | 200.00 | 30.00 | N | 80.00 | 150.00 |
| 88Ws12 | 20.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | N | 25.00 | 150.00 |
| 88Ws13 | 30.00 | <2.00 | 20.00 | N | 300.00 | 200.00 | 30.00 | <200.00 | 90.00 | 100.00 |
| 88Ws14 | 20.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | N | 25.00 | 150.00 |
| 88Ws15 | 20.00 | <2.00 | 30.00 | N | 500.00 | 300.00 | 30.00 | N | 55.00 | 70.00 |
| 88Ws16 | 20.00 | <2.00 | 30.00 | N | 500.00 | 300.00 | 20.00 | <200.00 | 40.00 | 70.00 |
| 88Ws17 | 15.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | N | 65.00 | 100.00 |
| 88Ws18 | 20.00 | <2.00 | 30.00 | N | 500.00 | 200.00 | 30.00 | N | 70.00 | 150.00 |
| 88Ws19 | 10.00 | <2.00 | 20.00 | N | 500.00 | 200.00 | 30.00 | N | 45.00 | 100.00 |
| 88Ws20a | 20.00 | <2.00 | 20.00 | N | 300.00 | 150.00 | 30.00 | N | 45.00 | 200.00 |
| 88Ws20b | 15.00 | <2.00 | 20.00 | N | 500.00 | 150.00 | 30.00 | <200.00 | 75.00 | 150.00 |
| 88Ws21a | 20.00 | <2.00 | 15.00 | N | 200.00 | 150.00 | 30.00 | N | 75.00 | 200.00 |
| 88Ws21b | 20.00 | <2.00 | 15.00 | N | 300.00 | 50.00 | 30.00 | N | 30.00 | 200.00 |

Table 2 Sample coding for rock samples.

| Sample type | Sample source | Rock class |
|--|---|--|
| A Grab sample B Composite | A Outcrop or rubble B Mine C Dump or prospect D Float G Other | B Sedimentary C Metamorphic D Igneous E Unconsolidated |
| Quad | FC3 (Type of collection) | FC4 (Class of sample): |
| A to D and 1 to 6 depending on 1:63,360 sheet. | A Primary sampling B Resampling C Replicate sample D Reanalysis | A Background (major) B Background (minor) C Atypical (very minor) D Atypical (mineralized) |
| Form | Rock type | |
| A Bed B Mass C Foliated D Pluton E Dike or sill F Extrusive G Vein H Fault I Other | A Feldspathic sandstone B Lithic sandstone C Sandstone D Shale or mudstone E Siltstone F Conglomerate G Chert H Argillite I Limestone | J basalt K Volcanic breccia L Andesite M Tuff N Quartz diorite O Granodiorite P Quartz monzonite Q Diorite Z Other |

Table 3 Lower limit of determination for respective elements (All analyses are by emission spectrography unless otherwise noted; AA indicates atomic absorption spectrophotometry)

| Element | Limit | Element | Limit | Element | Limit | Element | Limit |
|---------|-------|---------|-------|---------|-------|---------|-------|
| Fe | 0.05 | B | 10 | Ga | 5 | Sc | 5 |
| Mg | 0.02 | Ba | 20 | Ge | 10 | Sn | 10 |
| Na | 0.2 | Be | 1 | La | 50 | Sr | 100 |
| P | 0.02 | Bi | 10 | Mn | 10 | Th | 100 |
| Ca | 0.05 | Bi-AA | 1 | Mo | 5 | V | 10 |
| Ti | 0.002 | Cd | 20 | Nb | 20 | W | 20 |
| Ag | 0.5 | Cd-AA | 0.1 | Ni | 5 | Y | 10 |
| As | 200 | Co | 10 | Pb | 10 | Zn | 200 |
| As-AA | 10 | Cr | 10 | Sb | 100 | Zn-AA | 5 |
| Au | 10 | Cu | 5 | Sb-AA | 2 | Zr | 10 |

Table 4. Samples that equal or exceed threshold values in ppm for lead (Pb), copper (Cu), zinc (Zn), or molybdenum (Mo). Threshold values for the anomalous metals are: Pb = 30 ppm, Cu = 100 ppm, Zn = 100 ppm, Mo = 5 ppm.

| | | | |
|-----------|-------|-----------|-------|
| 88ADg 2a | Pb | 88AWs 1a | Pb,Zn |
| 88ADg 5 | Pb | 88AWs 2b | Pb,Mo |
| 88ADg 6 | Pb | 88AWs 4a | Pb,Zn |
| 88ADg 15 | Pb | 88AWs 5a | Pb,Zn |
| 88ADt 9a | Cu | 88AWs 5b | Pb |
| 88ADt 10 | Pb | 88AWs 6b | Pb |
| 88ADt 10c | Pb,Zn | 88AWs 10b | Pb |
| 88ADt 11 | Pb | 88AWs 13 | Pb |





Figure 1. Index map of study area.



EXPLANATION

- Alluvium (Quaternary)
 - Quartz diorite (Pliocene to late Miocene)
 - Belkofski(?) Formation (middle Miocene to late Oligocene)
 - Volcanic Intrusive centers
 - Volcanic rocks, undivided
 - Belkofski Formation (middle Miocene to late Oligocene)
 - Unga Formation (middle Miocene to late Oligocene)
 - Stepovak Formation of Burk (1965) (Oligocene and Eocene)
- Contact
 Dashed where approximately located
 dotted where concealed

Figure 2. Geologic Map of King Cove area. Geology revised from McLean and others, 1978.




Figure 3. Station location map.