

U. S. Department of the Interior
U. S. Geological Survey

Devonian-Mississippian Carbonate Sequence in the Maiyumerak Mountains, Western Brooks
Range, Alaska
by

Julie A. Dumoulin¹ and Anita G. Harris²

1992

Open File Report 92-3

This report is preliminary and has not been reviewed for conformity with Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U. S. Government.

¹Anchorage, Alaska

²Reston, Virginia




TABLE OF CONTENTS

ABSTRACT	1
ACKNOWLEDGEMENTS	1
INTRODUCTION	1
GENERAL GEOLOGY	3
STRATIGRAPHIC NOMENCLATURE	4
PREVIOUS WORK	4
METHODS	5
SEDIMENTARY SUCCESSION IN THE MAIYUMERAK MOUNTAINS	5
BAIRD GROUP	8
Lithologies	8
Laminated Mudstone	8
Bioclastic Wackestone	12
Peloidal Grainstone	15
Argillaceous Dolosiltstone	19
Distribution	19
Age and Sedimentation Rate	19
Depositional Environment	21
ELI LIMESTONE	23
Lithologies	23
Dolosiltstone	23
Nodular Limestone	23
Skeletal Wackestone/Packstone	27
Echinoderm-bearing Crystalline Limestone	27
Distribution	30
Age and Sedimentation Rate	30
Depositional Environment	31
UTUKOK FORMATION	33
Lithologies	33
Siliciclastic Rocks	35
Quartz Arenite	35
Quartzose Calcarenite	35
Siltstone and Shale	39
Skeletal Grainstone	39
Skeletal Wackestone/Packstone	39
Distribution	41
Age and Sedimentation Rate	41
Depositional Environment	41
KOGRUK FORMATION	43
Lithologies	43
Skeletal Wackestone	44
Echinoderm-bearing Crystalline Limestone	44
Spiculitic Limestone	44
Distribution	49
Age and Sedimentation Rate	49
Depositional Environment	49
SUMMARY OF THE MAIYUMERAK MOUNTAINS CARBONATE SEQUENCE	51
REGIONAL RELATIONSHIPS	52
DEFINITIONS OF SEQUENCES AND ALLOCHTHONS	57
CORRELATION	57

Baird Group	57
De Long Mountains Allochthon Belt	58
Northwestern Area	58
Southeastern Area	59
Schwatka Mountains Structural Province	59
Summary	59
Eli Limestone	59
De Long Mountains Allochthon Belt	60
Argillaceous Carbonate Rocks	60
Clean Carbonate Rocks	60
Sandy Carbonate Rocks	60
Summary	61
Utukok Formation	61
De Long Mountains Allochthon Belt	61
Impure Carbonate Rocks	61
Black Shale	62
Summary	63
Kogruk Formation	63
De Long Mountains Allochthon Belt	63
Lisburne Hills	64
Schwatka Mountains Structural Province	64
Summary	65
Correlation Summary and Discussion	65
PALEOGEOGRAPHY	67
Paleogeographic Model	67
Paleogeographic Implications of Maiyumerak Sequence	67
Relationship Between Paleozoic Carbonate Platforms	69
Source of Clastic Detritus in the Maiyumerak Sequence	70
Relationship Between Carbonate Platform and Endicott	
Clastic Wedge	71
Demise of Kelly Platform	74
CONCLUSIONS	75
REFERENCES CITED	77

LIST OF ILLUSTRATIONS

Figure 1. Bedrock geology of the Maiyumerak Mountains area	2
Figure 2. Locations of measured sections and lithologic and fossil collections.	6
Figure 3. Devonian-Mississippian carbonate succession in the Maiyumerak Mountains quadrangle.	7
Figure 4. Lithologic and paleontologic symbols used in this report.	9
Figure 5. Generalized section, Baird Group and Eli Limestone.	10
Figure 6. Sedimentary features of laminated mudstone lithology, Baird Group.	11
Figure 7. Sedimentary features of laminated mudstone lithology, Baird Group.	13
Figure 8. Stromatoporoid wackestone, Baird Group.	14
Figure 9. Photomicrographs of peloidal grainstone lithology, Baird Group.	16
Figure 10. Photomicrographs of dolomitized peloidal grainstone lithology, Baird Group.	18
Figure 11. Laminated mudstone overlying peloidal grainstone, Baird Group.	20
Figure 12. Bioclastic limestone overlying argillaceous dolostone, Eli Limestone.	24
Figure 13. Microtextures of dolosiltstone, Eli Limestone.	25
Figure 14. Nodular limestone, Eli Limestone.	26
Figure 15. Microtextures of Eli Limestone.	28
Figure 16. Photomicrographs of recrystallized limestone, Eli Limestone.	29
Figure 17. Generalized section, Utukok and Kogruk Formations.	34
Figure 18. Sedimentary features of Utukok Formation.	36
Figure 19. Photomicrographs of Utukok Formation.	37
Figure 20. Skeletal wackestone/packstone, Utukok Formation.	40
Figure 21. Black chert nodules in bioclastic limestone, Kogruk Formation.	45
Figure 22. Photomicrographs of skeletal wackestone, Kogruk Formation.	46
Figure 23. Sedimentary features, Kogruk Formation.	47
Figure 24. Photomicrograph of spiculitic limestone, Kogruk Formation.	48
Figure 25. Central and northern Alaska, showing area of western Brooks Range discussed in this chapter.	53
Figure 26. Distribution of allochthons and sedimentary sequences in De Long Mountains allochthon belt.	54
Figure 27. Age, lithology, stratigraphy, and thickness of Devonian-Mississippian rocks in northwestern Alaska.	55
Figure 28. Schematic paleogeographic map of northwestern Alaska in Late Devonian-Early Mississippian time.	68
Figure 29. Alternate paleogeographic reconstructions of northwestern Alaska in Late Devonian-Early Mississippian time.	73

Handwritten notes, possibly bleed-through from the reverse side of the page. The text is illegible due to blurriness and low contrast.

Vertical text on the right side of the page, likely bleed-through from the reverse side. The text is illegible.



Vertical text on the left side of the page, likely bleed-through from the reverse side. The text is illegible.

Vertical text on the right side of the page, likely bleed-through from the reverse side. The text is illegible.

Small handwritten mark or signature at the bottom right corner.



ABSTRACT

Virtually continuous, dominantly carbonate sedimentation occurred from at least Middle Devonian through early Late Mississippian time in the area that is now the Maiyumerak Mountains, western Brooks Range. Conodonts and megafossils throughout the Maiyumerak sequence indicate that any hiatus is less than a stage in duration, and there is no apparent physical evidence of an unconformity within the succession. The sequence is best exposed northwest of the Eli River, where Eifelian-Givetian dolostone (Baird Group) is conformably overlain by Givetian-Famennian argillaceous limestone (Eli Limestone). These rocks, in turn, are overlain conformably by Kinderhookian-Osagean quartzose limestone (Utukok Formation) and Osagean-Meramecian cherty dolomitic limestone (Kogruk Formation) of the Lisburne Group. Sedimentary structures and conodont species assemblages indicate deposition in a range of shallow-water, middle to inner shelf environments; the shallowest and most restricted depositional regimes prevailed during late Middle Devonian time.

The Maiyumerak sequence is in striking contrast to Paleozoic successions in adjacent fault slices in northwestern Alaska, in the eastern Brooks Range, and in the subsurface across northern Alaska. Other successions contain thick intervals of Upper Devonian-Mississippian siliciclastic rocks, condensed basinal Carboniferous facies, or an unconformity which spans much or all of Devonian and Early Mississippian time. Sedimentologic and biostratigraphic data support correlation of the Maiyumerak sequence with sequences of the Kelly River allochthon to the north and west, and suggest that these sequences were deposited contiguously on a single geographic high, the Kelly platform. Terrigenous material in the Maiyumerak sequence matches the Endicott Group in age and composition, and probably derived from a common source. This conclusion implies that the Kelly platform and the Endicott delta were at least partly adjacent during Late Devonian-Early Mississippian time, and offers a new constraint for paleogeographic reconstructions of the western Brooks Range. Inundation of the Kelly platform in the late Meramecian was probably due largely to tectonic factors, such as northward drift of northern Alaska, abetted by short-term eustatic shifts.

ACKNOWLEDGMENTS

This study was part of a survey of the geology and mineral resources of the Baird Mountains 1 x 3 degree quadrangle, conducted by the U.S. Geological Survey under the auspices of the Alaska Mineral Resource Assessment Program (AMRAP). We gratefully acknowledge the assistance of William Thompson and Inyo Eilersieck in the field, and thank all our AMRAP colleagues for logistical support and useful geologic discussions, particularly Susan M. Karl, Jeanine M. Schmidt, and Allison B. Till.

INTRODUCTION

The Maiyumerak Mountains are in the southwestern part of the Brooks Range in northwestern Alaska, and are made up of a diverse assemblage of Paleozoic and Mesozoic sedimentary and igneous rocks (Fig. 1; Karl and others, 1989). The Devonian-Mississippian carbonate rocks described in this report comprise a northeast-trending, fault-bounded sequence that is exposed for at least 75 km along the southeast flank of the Maiyumerak Mountains; this study concentrates on the northeastern exposures of this sequence in the Baird Mountains quadrangle.

In order to facilitate correlation with previous geologic studies of northern Alaska, the term "sequence" is used here in the sense of Mayfield and others (1988) and not as it has been employed in seismic stratigraphy. Mayfield and others (1988, p. 146) considered the thrust sheets of northwestern Alaska to be made up of "structurally overlapping sequences....composed of sedimentary,



EXPLANATION




Figure 1. Bedrock geology of the Maiyumerak Mountains area, Baird Mountains quadrangle, northwestern Alaska, generalized and slightly modified from Karl and others (1989).

and Harris, 1987) did not include rocks of the Maiyumerak Mountains.

METHODS

The carbonate rocks described in this report were observed and sampled in a series of partial measured sections along the cutbanks of Ahaliknak Creek and in the mountains to the north and south; this area of detailed study extends about 11 km and lies within a single fault block (Figs. 1, 2, and 3). Less detailed lithologic studies and sampling were also carried out further north and south along the trend of the Maiyumerak Mountains carbonate sequence within the Baird Mountains quadrangle.


Petrographic descriptions are based on field studies of lithology and sedimentary structures as well as examination of polished slabs and more than 200 thin sections. Sections were measured using Jacob's staff, Brunton compass, and tape; identification of calcite and dolomite was made using the Alizarin Red-S and potassium ferricyanide staining technique of Dickson (1966). Carbonate rocks are classified following Dunham (1962); when descriptive modifiers are used, they are listed in order of increasing abundance. Biostratigraphy relies largely on conodont faunas from our measured sections and spot samples, but data from previous U.S. Geological Survey collections of foraminifers, brachiopods, corals, and stromatoporoids that were made within the study area have been included where appropriate. Interpretations of depositional environments are based on depositional models for carbonate rocks outlined in Wilson (1975), and on a version of these models designed specifically for the Lisburne Group by Armstrong and Mamet (1977, 1978). The environmental implications of conodont assemblages, as outlined by Sandberg (1976), Sandberg and Gutschick (1984), and Sandberg and Ziegler (1979), were also used to constrain our interpretations.

SEDIMENTARY SUCCESSION IN THE MAIYUMERAK MOUNTAINS

The original configuration of the basin in which Paleozoic carbonate rocks in northwestern Alaska were deposited has been obscured by subsequent regional metamorphism and tectonic dismemberment, and is at present poorly understood. Most carbonate rocks throughout the Brooks Range occur in discrete, thrust-bounded sequences; thus, the initial relationships between these sequences, the nature of the basement on which they accumulated, and the type of shelf-to-basin transitions they once formed is uncertain. No encompassing depositional model has yet been presented for Devonian and older carbonate rocks of northwestern Alaska; the biostratigraphy and lithofacies of these rocks are only beginning to be unraveled (e.g., Dumoulin and Harris, 1987).

The Carboniferous Lisburne Group has been more studied, and general depositional patterns across the Brooks Range have been described in a series of publications by Armstrong and others (e.g., Armstrong, 1974; Armstrong and Bird, 1976; Armstrong and Mamet, 1978). These studies suggest that much of the Lisburne Group was deposited on a carbonate platform with a low-gradient, south-facing ramp margin that developed after initial northward transgression across an irregular surface. However, this model is based largely on sections in the northeastern Brooks Range; Carboniferous paleogeography to the west may have been more complex and involved a number of small (rift?) basins (Mayfield and others, 1988).

Devonian-Mississippian carbonate rocks in the Maiyumerak Mountains appear to have formed through relatively continuous deposition in a range of shallow-water environments. In the following sections, the general terms "carbonate platform" and "shelf" will be used to describe the depositional setting of these rocks; speculations concerning the shape of this platform and its relationships to other geographic elements of northwestern Alaska will be further discussed below.



EXPLANATION

MDI Limestone (Mississippian and (or) Devonian)

Maiyumerak carbonate sequence

Mk Kogruk Formation (Mississippian)

MDue Utukok Formation and Eli Limestone
(Mississippian and Devonian)

Db Baird Group (Devonian)

Other symbols

∠ Bedding attitude

/ Contact

— Fault

↘ Thrust fault, sawteeth
on upper plate

↗ Normal fault, ball-and-stick
on down-dropped plate

/ (A) Measured section locality
referred to in text

■ (B) Fossil collection locality
referred to in text

Figure 2. Locations of measured sections and lithologic and fossil collections, Maiyumerak Mountains area, Baird Mountains quadrangle.




Figure 3. Devonian-Mississippian carbonate succession in the Maiyumerak Mountains.

BAIRD GROUP

Medium- to dark-gray, medium- to light-gray-weathering dolostone and lesser dolomitic limestone and limestone forms the base of the carbonate sequence in the Maiyumerak Mountains and is here assigned to the Baird Group. Exposures include massive cliffs, but consist largely of blocky rubble. The unit may be as much as 1000 m thick in this area (Tailleur and others, 1967); the base is truncated by a thrust fault which places carbonate rocks above siliciclastic rocks of the Endicott Group. The upper part of the Baird Group includes a few thin beds of argillaceous dolosiltstone similar to the overlying Eli Limestone; the contact is placed above the highest light-gray-weathering dolomitic bed, and is marked by a sharp change in overall weathering color, from light-gray below to brown above.

The Baird Group was most closely studied in an incomplete measured section 440 m thick on the north side of Ahaliknak Creek (Fig. 2, locality A; Figs. 4 and 5); the measured section includes only the upper part of the unit and at least several hundred meters of additional section are exposed at this locality. Reconnaissance studies through the entire Baird Group were made along a series of traverses in the mountains to the north and south.

Lithologies

Most of the Baird Group in the Maiyumerak Mountains has been thoroughly dolomitized but retains considerable relict primary texture that is visible both megascopically and microscopically. Some beds, primarily towards the top of the section and in outcrops in the southern part of the study area, have been only slightly to partially dolomitized; these intervals provide additional information on the composition of the Baird Group prior to dolomitization. Lithologies discerned in the Baird Group and described below are: laminated mudstone, bioclastic wackestone, peloidal grainstone, and argillaceous dolosiltstone.

Laminated Mudstone

Light- to dark-gray, fine-grained dolostone and lesser limestone that is generally laminated and contains fenestral fabric, local sheet cracks, desiccation cracks, and evaporite laths occur throughout the Baird Group. Relict microtextures indicate that most of these rocks originated as carbonate mudstone; they are associated with thin beds of grainstone and flat-pebble conglomerate.

Fine lamination is the most notable megascopic feature of these mudstones. Laminae are 0.5-5 mm thick, crinkled and irregular, form small-scale hummocks with several mm of relief, and are interpreted as cryptalgal in origin (Fig. 6A). Fenestral fabric is well-developed in these rocks (Fig. 6B) and occurs in intervals a few cm to 20 m thick. Fenestrae range from horizontal, laminar forms that are 0.5-1 mm X 3-10 mm in size, to more irregular shapes as much as 5 mm across. Relatively coarse, sparry calcite or coarsely crystalline dolomite fills the fenestrae. Sheet cracks occur as horizontal zones a few cm thick and several cm long that contain finely laminated, red-weathering sediment. In some cases, a layer of coarse dolomite or calcite spar overlies (and rarely, underlies) the red sediment, coating the upper and lower surfaces of the crack (Fig. 6C). V-shaped, vertical cracks as much as 0.5 cm wide at the top and 1.5 cm deep are interpreted as desiccation cracks; they are filled with red sediment like that found in the sheet cracks (Figs. 6C and D).

Most of the rocks that contain the distinctive sedimentary features described above are dolostone, and consist largely of euhedral to subhedral dolomite crystals 8 to 80 μ m in size. Such saccharoidal dolomite typically replaces carbonate mudstone (Armstrong, 1970b), and the fine-grained laminated dolostone in the Baird Group is thought to have originated in this way. This interpretation is

EXPLANATION FOR STRATIGRAPHIC COLUMNS




Figure 4. Lithologic and paleontologic symbols used in this report.




Figure 6. Sedimentary features of laminated mudstone lithology, Baird Group. Scale bar equals 1 cm in A-C, 1 mm in D. A. Whole thin-section photograph of cryptalgal layering in dolomitic mudstone, locality G, Fig. 2. B. Whole thin-section photograph of well-developed fenestral fabric in peloidal lime mudstone, locality D, Fig. 2. C. Whole thin-section photograph of sheet crack (S) and desiccation crack (D) in laminated dolomitic mudstone, locality G. D. Photomicrograph of anhydrite laths in sediment filling the desiccation crack shown in Fig. 6C.

strengthened by the occurrence of locally undolomitized or partly dolomitized intervals of carbonate mudstone that are interbedded with the laminated dolostone and possess identical megascopic features. For example, lime mudstone interbedded with dolostone at locality B (Fig. 2) contains cryptalgal lamination, well-developed fenestral fabric, and excellently preserved original microtexture. Fenestrae in these rocks are surrounded by lime mud or fine-grained peloids; the peloids have locally coalesced into a flocculent, clotty mass (*structure grumeleuse*) in which individual grain outlines are difficult to discern (Fig. 7A). The peloids are made of micrite (crystals 2-4 μm), have maximum diameters of 75 to 150 μm , and are quite uniformly ellipsoidal in shape.

Laminae in these rocks generally reflect subtle variations in dolomite color or crystal size or in the amount of intercrystalline opaque material, but some are formed by alternations of lime micrite and finely crystalline dolomite.

Several percent bioclasts and bioclastic debris occur disseminated throughout these fine-grained rocks. Bioclasts are best preserved in the undolomitized rocks but also occur as remnant forms in dolostone. Whole bioclasts are dominantly calcispheres that range from 60-300 μm in diameter and include thick-walled radiosphaerid types.

The red sediment in the sheet and desiccation cracks consists mostly of silt-sized dolomite euhedra and lesser bioclastic and quartz silt in a clay-sized matrix that appears dark brownish black in thin section. Laths of anhydrite a few mm long occur within the red sediment in both horizontal and vertical cracks, but are especially large and abundant at the base of the vertical cracks. The laths are generally randomly oriented within the mud matrix (Fig. 6D).

Lighter colored layers of relatively coarse grained peloidal and (or) bioclastic grainstone and packstone are locally interbedded with laminated mudstone (Fig. 7B). These layers are generally undolomitized or partly dolomitized, 0.5 to 5 cm thick, and some are distinctly graded. Peloids are irregular to ovoid in form, 0.1-0.5 mm in diameter, and very similar to those in the peloidal grainstone lithology described below. Recognizable bioclasts include calcispheres, foraminifers, and algae.

Rare flat-pebble conglomerate occurs in beds a few cm to 12 cm thick. Clasts are elongate (typical dimensions are 0.5-1.0 cm thick and 5-10 cm long), imbricated, and consist mostly of dolomitic laminated mudstone and subordinate peloidal grainstone.

Bioclastic Wackestone

Another major Baird Group lithology is bioclastic wackestone and lesser packstone; these rocks are mostly dolostone or slightly calcitic dolostone. Bioclasts are dominantly stromatoporoids, primarily the small, tubular form *Amphipora*, and more rarely, massive laminar forms 1-2 cm thick and 3-6 cm long (Fig. 8). Locally, amphiporids make up 30-70 percent of these rocks and form biostromes 0.5 to 2.5 m thick and several tens of m in lateral extent. Elsewhere, amphiporid material is less concentrated (10-30 percent of the rock) and occurs in even to undulatory beds 2-10 cm thick. The amphiporid tubes are 1-3 mm in maximum diameter and 1-2 cm long; details of internal structure are poorly preserved, as most tubes have been replaced by dolomite euhedra or rarely, chert. The matrix of the amphiporid rocks is variably grain-rich, fine-grained bioclastic wackestone; small bioclasts in the matrix are generally broken and not specifically identifiable, but whole forms observed include ostracodes and calcispheres. Bioclasts are surrounded by fine-grained saccharoidal dolomite; this dolomite is interpreted as a replacement of carbonate mudstone as described above.

Megafossils other than stromatoporoids are rare in the Baird Group in the Malyymerak




Figure 7. Sedimentary features of laminated mudstone lithology, Baird Group. A. Photomicrograph of *structure grummeuse* in peloidal limestone, locality D, Fig. 2. Small spheres (C) are calcispheres; calcite spar-filled voids (F) are fenestrae. Scale bar is 0.5 mm. B. Whole thin-section photograph of thinly interlayered bioclastic peloidal grainstone (light) and laminated mudstone (dark), locality G, Fig. 2. Scale bar is 1 cm.



CM

Figure 8. Stromatoporoid wackestone, Baird Group, Malyumerak Mountains. Laminar form on far left; stick-shaped forms mostly *Amphipora* sp. Field locality no. 86 AD 33.

Mountains. Corals, including *Thamnopora* sp., *Alveolites?* sp., *Cladopora ?*sp., *Dendrostella?* sp., and indeterminate disseminated rugose corals, are reported, generally associated with *Amphipora* sp. (written commun., W. A. Oliver, Jr., 1974). Like the amphiporids, the corals are typically poorly preserved and occur in dolomitized wackestone or packstone.

Two other types of bioclastic wackestone are observed in the Baird Group. Relatively grain-rich fine-grained wackestone forms blocky rubble and contains brachiopod, echinoderm, and bryozoan debris 0.1 mm to 1.0 cm in size. More grain-poor wackestone occurs in irregular bioturbated beds as much as 30 cm thick; bioclasts are generally less than 30 percent of the rock, smaller than 0.5 cm, and include ostracodes, gastropods such as *Macrocheilus* sp. (written commun., E. L. Yochelson, 1972), calcispheres, and rare sponge? spicules.

Peloidal Grainstone

The most widely distributed Baird Group lithology is grainstone and (or) packstone made up of peloids and lesser skeletal grains. Excellent exposures of this lithology that are undolomitized to only slightly dolomitized occur north of the Eli River (Fig. 2, locality C). Here, beds are even to slightly irregular, 2 to 80 cm thick, and consist of moderately well-sorted, fine- to medium-grained peloidal grainstone. The peloids are ellipsoidal to spherical to irregular in form, 0.1 to 0.5 mm in maximum dimension (average 0.2-0.3 mm), and are notably larger and more irregular in shape than those in the pelleted mudstones described above. Most consist entirely of micrite (crystals 2-3 μ m), but rare peloids are composed of skeletal fragments and micrite (Figs. 9A and B). Peloids constitute more than 95 percent of the grains in the samples examined; other grains are bioclasts, mainly echinoderm, bryozoan, and brachiopod fragments. The grainstone is closely packed (locally, overly closely packed with interpenetrative grain contacts) and has a matrix of fine crystalline sparry calcite.

Rocks similar to these peloidal grainstones but moderately to completely dolomitized make up much of the Baird Group in the Maiyumerak Mountains. Partially dolomitized rocks retain a sparry calcite matrix, but peloids have been replaced by dolomite euhedra that are 50 μ m or less in diameter and choked with opaque (probably organic) material (Fig. 9C). Completely dolomitized rocks consist of anhedral to subhedral dolomite crystals that range from 50 μ m to 1.5 mm in size (average 0.2-0.4 mm). Within the dolomite crystal mosaic, a second texture is clearly visible: a closely packed mass of brown ellipsoidal to irregular forms, 0.1-0.5 mm in maximum dimension (Fig. 9D). Individual ellipsoids clearly cross-cut dolomite crystal boundaries, and their size and shape are identical to those of the peloids in the limestones described above; however, they consist not of micrite but of a dark-brownish residue (organic matter?) enclosed within one or more dolomite crystals. These rocks also contain some spherical to irregular, relatively coarse single grains of dolomite that resemble echinoderm fragments (Figs. 10A and B).

It seems probable that the dolostone with brown ellipsoids was once lime grainstone and(or) packstone that has been completely dolomitized but has retained a palimpsest of original texture. The brown ellipsoids are ghosts of the original peloids, and the coarse, spherical to irregular dolomite crystals are echinoderm debris that has been pseudomorphically replaced. In some cases, both echinoderm fragments and associated calcite rim-cement overgrowths have been replaced; the relict contact between the more turbid echinoderm grain and the clearer rim cement is clearly discernable within the dolomite crystal (Fig. 10A). The tendency of dolomite to pseudomorphically replace crinoid fragments and rim cements was noted by Lucia (1962); similar textures were described by Armstrong (1970b) in the Lisburne Group north of the study area.

Figure 9. Photomicrographs of peloidal grainstone, Baird Group. A-C, locality C, Fig. 2; D, locality A, Fig. 2. Scale bars equals 1 mm in A, C, and D, 0.5 mm in B. A. and B. Undolomitized sample; grains are lime micrite peloids and rare skeletal fragments in a matrix of sparry calcite cement. Some peloids contain small skeletal fragments in addition to micrite. C. Partially dolomitized sample; peloids in lower right have been replaced by dolomite. D. Completely dolomitized sample; peloids (P), skeletal grains such as crinoid fragments (C), and sparry cement have been replaced by dolomite.

Argillaceous Dolosiltstone

The final lithology distinguished in the Baird Group of the Maiyumerak Mountains is brown- to red-weathering argillaceous dolosiltstone; it occurs only in the uppermost part of the unit in beds a few cm thick. These rocks are parallel- to cross-laminated and consist of dominantly silt-sized dolomite euhedra intermixed with argillaceous material and as much as 10 percent quartz and calcite silt. Calcareous grains include peloids and spicules. Laminae are straight and laterally continuous, not irregular and locally domed like the cryptalgal laminae in the laminated mudstone lithology, and are formed mostly by alternations of clay and silt. Similar rocks occur in the overlying Eli Limestone, and are more fully described below.

Distribution

Most of the Baird Group in the Maiyumerak Mountains consists of dolostone with relict peloidal grainstone or amphiporid wackestone texture. Wackestone containing corals, brachiopods, echinoderms, and (or) bryozoans is rare and was observed mainly in the lower half of the unit and in the uppermost beds. The upper half of the unit is characterized by alternations of the first three lithologies described above; some intervals are distinctly cyclic. A typical cycle is 2-5 m thick and consists of 1-2 m of peloidal grainstone or packstone overlain by a similar thickness of amphiporid wackestone. The wackestone grades up into a thinner interval of laminated mudstone with fenestral fabric, which is in turn overlain by another bed of peloidal grainstone (Fig. 11). The cycles appear to have formed under shallowing-upward (regressive) conditions, because transitions from peloidal grainstone up into amphiporoid wackestone and from amphiporoid wackestone up into laminated mudstone are generally gradational, whereas those from laminated mudstone up into peloidal grainstone are abrupt (cf. otherwise similar cycles described by Fischer, 1975, interpreted as deepening-upward deposits). Bioclastic wackestone containing ostracodes, gastropods, calcispheres, and (or) spicules is locally associated with (and grades into) amphiporid wackestone and laminated mudstone. Sheet cracks, desiccation cracks, and evaporite laths were noted in laminated mudstone in the uppermost third of the unit (Fig. 5). At the very top of the section, argillaceous dolosiltstone that is lithologically similar to beds in the overlying Eli Limestone occurs interbedded with gray calcitic dolostone that contains corals and brachiopods. The base of the Eli Limestone is placed at the top of the highest gray dolostone bed.

Age and Sedimentation Rate

The Baird Group is the oldest unit known in the Maiyumerak Mountains; the base is not exposed and is consistently fault-bounded. Conodont faunas provide the most precise information currently available on the age of the Baird Group in the study area. The oldest biostratigraphically restricted conodont fauna obtained contains *Polygnathus* aff. *P. robusticostatus*, indicating an early Middle Devonian (earliest Eifelian through earliest Givetian) age (USGS colln. 11385-SD); this fauna occurs in a sample of bioclastic peloidal grainstone collected near the base of the section (Fig. 2, locality D). The youngest biostratigraphically restricted samples contain *Polygnathus xylus* and *P. cf. P. pseudofoliatus*, (USGS collns. 11137-SD and 11138-SD) which denote a middle(?) Givetian (possibly Lower to Middle *varcus* Subzone) age; these faunas come from argillaceous dolosiltstone and interbedded coralline dolostone that directly underlies the Eli Limestone (Fig. 2, locality A). All other samples from the Baird Group contain conodonts that are longer ranging within the late Early through Middle Devonian (Emsian through Givetian). Poorly preserved coral and stromatoporoid faunas from the Baird Group in the Maiyumerak Mountains yield long-ranging ages of Middle Devonian or Devonian (written commun., W. A. Oliver, Jr., 1974).

Because the age and total thickness of the Baird Group in the Maiyumerak Mountains is not




Figure 11. Laminated mudstone overlying medium- to thick-bedded peloidal grainstone, Baird Group, Malyumerak Mountains; field locality no. 86 AD 43.




Figure 13. Microtextures of dolosiltstone, Eli Limestone, locality A, Fig. 2. Whole thin-section photograph (A) and microphotograph (B) of cross-laminated dolosiltstone; laminae consist of alternations of lime peloids (P) and dolomite rhombs (D). Scale bar equals 1 cm in A, 1 mm in B.




Figure 14. Nodular limestone, Eli Limestone, locality A, Fig. 2.

nodules separated by seams of orange-weathering argillaceous dolostone. The nodules are a few mm to several cm long, cylindrical to ovoidal to irregular in shape, and locally form sedimentary boudinage or ball-and-flow structures (Fig. 15A); they consist of skeletal wackestone to packstone, locally peloidal (Fig. 15B). The dolostone seams consist of brownish euhedral dolomite crystals, 15-100 μm (average 40-50 μm) in diameter, with considerable (10-30 percent) dark-brown intercrystalline argillaceous material and as much as 5 percent disseminated quartz silt. Skeletal fragments also float within the dolostone seams, but are generally more abundant in the limestone nodules (Fig. 15C). Identifiable fossils found in both nodules and seams include coral and stromatoporoid fragments, brachiopods, gastropods, and calcispheres. Shelter porosity, now filled with calcite spar or geopetal layers of carbonate sediment and sparry cement, is locally well developed beneath and within larger shell fragments in the packstones; such umbrella effects indicate that the skeletal fragments were deposited in grain support (Dunham, 1962).

The nodular limestone lithology is interpreted as a combination of two discrete sediment types, skeletal wackestone to packstone and argillaceous carbonate mud, that were originally interbedded but have been partially mixed by bioturbation; the argillaceous carbonate mud has been thoroughly dolomitized. Locally, the light-gray limestone fills vertical and horizontal burrows a few mm to several cm in diameter, but discrete burrow forms are relatively rare. Both dissolution seams and fitted fabric, as defined by Bathurst (1987, p. 763), occur locally within this lithology; the former are "smooth, undulose seam(s) of insoluble (noncarbonate) residue which lack the sutures of stylolites" and the latter is "a framework of interpenetrant grains."

Skeletal Wackestone/Packstone

Subordinate intervals, generally less than 2 m thick, of thin-bedded skeletal wackestone and lesser packstone that contain little argillaceous material occur throughout the Eli Limestone. Beds are 2-10 cm thick and generally nodular, but locally more even, planar beds predominate. Most intervals are nodular bedded, relatively grain-rich wackestone to packstone that is thoroughly bioturbated; bioclasts include fenestrate bryozoans, brachiopods, echinoderms, ostracodes, and foraminifers, as well as unidentified calcite spheres and spicules (Fig. 15D). Shelter porosity, now filled with calcite spar, is locally well-developed under larger shelly fragments such as brachiopods in packstone (Fig. 15D). More evenly bedded intervals of this lithology contain a similar range of relatively well-preserved bioclasts, but also include up to 30 percent peloids and 10 percent quartz and subordinate feldspar silt and fine sand (Fig. 15E). Some samples contain a few percent disseminated dolomite rhombs.

Echinoderm-bearing Crystalline Limestone

Intervals of echinoderm-bearing crystalline limestone less than a meter to several tens of meters thick occur throughout the Eli Limestone but are especially abundant in the upper part of the unit. Some intervals contain virtually no argillaceous material, weather light gray, and are crudely bedded on a scale of 10-40 cm. Other intervals weather orange and consist of nodular beds 2-10 cm thick separated by irregular, undulatory partings of argillaceous dolostone. Locally, beds are so nodular and argillaceous material is so abundant that this lithology grades into the nodular limestone lithology; most of the crystalline limestone, however, contains little (0-5 percent) argillaceous material. Lenses and nodules of black to gray chert a few cm thick occur rarely. The limestone consists mostly of relatively coarse and inequigranular calcite crystals, dominantly 0.5-5.0 mm in diameter. Much of the calcite is anhedral but relict echinoderm fragments (0.2-2.5 mm) are abundant and readily identifiable (Fig. 16A). In some intervals, saccharoidal dolomite and (or) opaque material forms irregular wisps and patches between the calcite crystals (Fig. 16A). Some samples contain a few percent disseminated quartz (and rarely feldspar) silt.




Figure 15. Microtextures of Eli Limestone. A-C, locality A, Fig. 2; D-E, locality G, Fig. 2. A and C are whole thin-section photographs; B, D, and E are photomicrographs. Scale bar equals 1 cm in A and C, 1 mm in D, and 0.5 mm in B and E. A. Ball-and-flow structure in nodular limestone; nodules of light-colored limestone in darker dolomitic, argillaceous matrix. B. Close-up of A, showing peloids in limey nodule (above) and dolomite rhombs in argillaceous matrix (below). C. Stick-like tabulate corals, stomatoporoids, and nodules of peloidal limestone in dolomitic, argillaceous matrix. D. Skeletal packstone containing bryozoa (B), echinoderm fragments (C), and brachiopod shells (Br); shelter porosity under brachiopod shells is filled with calcite spar (S). E. Peloidal bioclastic packstone with minor quartz silt (Q); recognizable skeletal grains include echinoderm fragments (C).

Conodonts are relatively sparse in the Eli Limestone; interpretations of the depositional environment suggested by conodont biofacies agree with those based on sedimentary structure outlined above. Conodont faunas from nodular limestone, skeletal wackestone, and echinoderm-bearing recrystallized limestone are similar and consist mostly of polygnathids with lesser apatognathids and minor icriodids and "ozarkodinids". This species association represents the polygnathid biofacies and suggests a shallow to moderate depth, normal marine depositional environment (Sandberg and Ziegler, 1976; Sandberg, 1976). The laminated dolosiltstone lithology has produced only Dvorakias(?) and poorly preserved polygnathids that appear to have undergone hydraulic sorting and abrasion; as previously discussed, these assemblages may have been reworked.

Clay-rich bedding surfaces in the middle shelf environment typically represent diastems; hiatuses occur mostly during periods of terrigenous sedimentation that separate spurts of more rapid and continuous carbonate deposition (Wilson and Jordan, 1983). Numerous, brief episodes of non-deposition could account for the overall low sedimentation rate of the Eli Limestone; no discrete erosional surfaces were found in the sections examined.

In summary, sedimentologic and paleontologic evidence suggest that the lower (Frasnian) part of the Eli Limestone was deposited in a somewhat restricted inner shelf setting, whereas the upper (Famennian) part of the section accumulated in a more open middle shelf environment. Argillaceous material occurs throughout the formation but is more abundant in the lower beds; grainstone shoals are developed primarily in the upper part of the unit. No major erosional surfaces were observed within the Eli Limestone, but numerous diastems may occur throughout the section.

UTUKOK FORMATION

The Utukok Formation in the Maiyumerak Mountains consists of gray to brown limestone, sandy limestone, quartzose to calcareous sandstone, and subordinate siltstone and shale; it is at least 180 m thick and exposed in a series of resistant limestone and sandstone ridges separated by recessive intervals of siltstone and shale. Dolomite and chert occur only locally in carbonate rocks of this unit, and constitute less than 15 percent of most samples studied. Like the Eli Limestone, the Utukok Formation is variably recrystallized within the study area; our analyses concentrated on sections with relatively well-preserved original textures. The unit was most closely studied in a measured section along the ridgetop due south of the type section of the Eli Limestone (Locality I, Fig. 2), and in several other ridgetop sections to the north and south (Localities J and K, Fig. 2). The contact with the underlying Eli Limestone is drawn at the base of the lowest calcarenite bed; the upper contact with the Kogruk Formation is placed at the top of the highest interval of siltstone and shale.

Lithologies

Siliciclastic rocks, mostly fine-grained sandstone, lesser siltstone, and minor shale, make up about two-fifths of the Utukok Formation in the Maiyumerak Mountains and increase in abundance upward within the section (Fig. 17). The clastic rocks are plane- to cross-laminated and range from relatively pure quartz arenite to quartzose calcarenite; they are interbedded with skeletal lime grainstone, packstone, and wackestone that contain little siliciclastic material. No completely satisfactory nomenclatural scheme has yet been proposed for mixed calcareous-quartz sediments (Zuffa, 1980; Mount, 1984). In the descriptions below, quartz arenite is used to describe sandstone that contains more than 80 percent quartz (not including chert) and grainstone is used for rocks that consist mostly of carbonate clasts and include less than 10 percent noncarbonate detrital grains. Sandstone of intermediate composition is called calcarenite.

Siltclastic Rocks

Quartz Arenite Relatively pure quartz arenite occurs in the upper part of the Utukok Formation, most notably about 20 m below the top of the unit in a distinctive, white-weathering horizon that is 10 m thick and can be traced along strike for at least 12 km. Within this horizon, quartz arenite is interbedded with subordinate quartzose calcarenite and brachiopod coquina. The quartz arenite forms even beds 4-30 cm thick that are plane- and cross-laminated; cross laminae are 1-5 cm thick.

A combination of textural immaturity and compositional maturity characterizes this lithology. Samples are fine- to very fine-grained, sorting is good, but grains are primarily subangular to subrounded (Fig. 18A). Silica cement has overgrown most grains but original grain outlines can still be seen because inclusions have been trapped at the boundary between detrital and authigenic quartz. Later pressure solution has produced local serrate grain contacts and wispy stylolites. A few samples (those richest in calcareous grains) contain patchy, locally poikilotopic calcite cement.

Quartz grains constitute 80-95 percent of the samples studied and are dominantly monocrystalline with straight to slightly undulose extinction. Other grains are mostly sedimentary lithic clasts and include chert, shale and mudstone, limestone, calcareous bioclasts such as crinoid fragments, dolomite rhombs, white mica, opaque grains, tourmaline, and zircon (Fig. 18A). The dolomite rhombs appear to be detrital and not authigenic, as shown by presence of rounded rhomb corners, equivalence of dolomite crystal size and sediment grain size, and a lack of replacement textural relations.

Quartzose Calcarenite Quartzose calcarenite occurs in intervals 10-25 m thick throughout the Utukok Formation. Beds are thin to medium (1-20 cm) and mostly even to slightly irregular; local beds are lenticular and have distinctly channeled bases. The sediment is plane- or cross-laminated fine- to very fine-grained sandstone and lesser siltstone; cross beds are low angle and form sets up to 5 cm thick. Local vertical burrows a few mm to a cm in diameter cut some sedimentary laminae. Differences in the amount of quartz and carbonate in these sediments are very evident in outcrop weathering profile and color: slightly calcareous quartz arenite is resistant and white to orange or pink, quartzose calcarenite is more recessive and tan to light brown, and calcarenite with subordinate quartz is distinctly recessive and light gray (Fig. 18B). Relatively quartz-rich sandstone and more calcareous sandstone are intercalated throughout the Utukok Formation on a scale ranging from several cm to tens of meters.

Carbonate grains make up 30-85 percent of most samples and are mostly bioclasts, peloids, calcareous lithic clasts, and dolomite rhombs. Bioclasts are chiefly echinoderm and lesser brachiopod fragments, but include ostracodes, foraminifers, and calcispheres; some bioclasts have well-developed micritic rims. Peloids are chiefly ovoid (less commonly spherical to irregular) in form, and 0.2 mm or less in maximum dimension. Micritic intraclasts as much as 3 mm in diameter have very irregular shapes and appear to have been deposited while still soft. Dolomite rhombs typically have iron-rich reddish-brown rims and clear centers, are the same size as associated quartz grains, and appear to be detrital as described above. Some patchy dolomite may be cement, but sparry calcite is the predominant cement in the calcarenite samples studied.

Monocrystalline quartz grains identical in size and shape to those in the quartz arenite comprise 30 to 70 percent of most quartzose calcarenite. Other noncarbonate grains consist chiefly of chert and lesser mudstone, tourmaline, and opaques.

Size sorting in these mixed carbonate-quartz sandstones is notably poorer than in the quartz arenites; for example, crinoid columnals to 2 mm occur within very fine grained quartz sand (Fig. 19A). This disparity probably does not indicate reduced current action but instead reflects the hydraulic




Figure 18. Sedimentary features of Utukok Formation. A. Photomicrograph (crossed nicols) of fine-grained quartz arenite, locality K, Fig. 2. Grains are subangular to subrounded and consist mostly of quartz (Q) and lesser chert (C). Scale bar is 0.5 mm. B. Interbedded quartzose calcarenite (Q) and less resistant calcarenite (C).

level fall. Recently published eustatic curves for the late Paleozoic (Ross and Ross, 1988) indicate generally rising long-term sea level throughout the Early Mississippian, punctuated by several short-term falls in the Osagean. Quartz sand is most abundant in the upper (Osagean) part of the Utukok Formation, but paleontologic control in these strata is not precise enough to postulate correlation of individual sand layers with specific global sea level events.

Conodont faunas from the Utukok Formation support the conclusion that most of the section accumulated in normal marine, moderate energy conditions in water of shallow to moderate depth. Faunas from the lower third of the unit are typical of relatively shallow-water, moderate-energy environments. A fauna representative of the hindeodid biofacies of Sandberg and Gutschick (1984), which indicates a shallow-water, possibly restricted depositional environment, was collected from peloidal wackestone in the middle of the section. The uppermost beds of the formation yield conodonts characteristic of a normal-marine, outer platform or shelf setting (Sandberg and Gutschick, 1984).

The Utukok Formation thus appears to have been deposited in a middle to inner shelf environment characterized by generally normal marine salinity and moderate energy and water depths; some faunal evidence for restricted circulation is noted about the middle of the section. Calcareous and (or) quartz sandstone formed in local shoals and was introduced into muddier environments by storms; sandy lithologies and quartz are most abundant in the upper (Osagean) part of the formation. No paleontological evidence of hiatuses and no physical evidence of exposure and (or) erosional surfaces have been found within the Utukok Formation, but the low sedimentation rate, like that in the underlying Eli Formation, suggests diastems occur within the section.

KOGRUK FORMATION

The Kogruk Formation in the Maiyumerak Mountains consists of light- to medium-gray-weathering, medium- to dark-gray, fine- to coarse-grained limestone and dolostone. Chert, black or less commonly gray or white, forms layers or nodules a few cm thick that constitute 10-50 percent of most sections; the amount of chert varies markedly both laterally and vertically. The unit typically crops out in terraced hillsides made up of resistant low cliffs of skeletal grainstone or packstone and less resistant rubbly intervals of wackestone or mudstone. The top of the Kogruk Formation in the Maiyumerak Mountains is an erosional surface; the lower contact is gradational into the underlying Utukok Formation and is placed above the highest bed of sandstone or shale more than 1 m thick. At locality I (Fig. 2) in the central part of the study area, the Kogruk Formation is 163 m thick; further north, more than 300 m of section may be exposed (Karl and others, 1989). Our studies concentrated on the measured section at locality I, another section 6 km north (locality M), and the spiculitic limestone belt north of Kivivik Creek (localities N and O).

Lithologies

Recrystallization, dolomitization, and silicification have locally obscured original texture and composition of the Kogruk Formation, but three major lithologies can be recognized. Skeletal wackestone, typically dolomitic and (or) cherty and rich in echinoderms and bryozoans, makes up most of the unit, and is intercalated with subordinate intervals of echinoderm-bearing crystalline limestone (Fig. 17). A third lithology, spiculitic limestone, has been found only in the uppermost part of the Kogruk Formation in the structurally complex area north of Kivivik Creek (Figs. 1, 2). These spiculitic rocks are similar to rocks assigned to the Tupik Formation (Sable and Dutro, 1961) in the Misheguk Mountain quadrangle.

Skeletal Wackestone

Bioclastic limestone relatively rich in lime mud and partly dolomitized and (or) silicified makes up most of the Kogruk Formation (Fig. 21). These rocks are dark gray, weather light to medium gray, and may have shaley partings. Most beds are wavy, irregular, or nodular; thickness varies but is generally 5-30 cm. Many beds are mottled and appear thoroughly bioturbated. Wackestone texture predominates, but packstone and mudstone occur locally. Peloids, generally ovoid and less than 200 μm in diameter, are locally abundant (20-30 percent) but everywhere subordinate to skeletal grains.

Bioclasts in most intervals consist primarily of echinoderm debris (crinoid columnals) but include notable bryozoan fragments as well as brachiopods and ostracodes (Fig. 22A). Large bryozoan fronds and articulated crinoid stems occur on some bedding planes. Foraminifers, calcareous algae, and calcispheres are locally common, particularly in peloid-rich intervals (Fig. 22B). Corals, chiefly lithostrotionids, occur sporadically in this lithology and may be as much several tens of cm across, but biostromal accumulations have not been recognized. Corals are most abundant in skeletal packstone interbedded with echinoderm-bearing crystalline limestone described below.

Much of the muddy matrix of this lithology has been altered to fine-grained, saccharoidal dolomite, but only rare, disseminated dolomite rhombs occur within bioclasts. Scattered quartz silt (a few percent) occurs locally. Black to gray chert, in lenses, nodules, and irregular masses a few to several tens of cm thick and 0.3 to more than 5 m long, constitutes 10-20 percent (rarely, 40 percent) of this lithology. Some discontinuous and irregular chert blebs appear to follow burrows (Fig. 23A).

Echinoderm-bearing Crystalline Limestone

Crystalline limestone forms intervals a few to tens of meters thick in the Kogruk Formation. In contrast to the rocks just described, this lithology is slightly lighter colored and contains little or no dolomite or chert. Beds are variable in thickness and somewhat irregular in form; most are 3-40 cm thick and slightly nodular. Most samples consist of coarse-grained sparry limestone in which abundant echinoderm fragments can be recognized; some samples also contain foraminifers and fragments of corals and bryozoans. Bioclasts are typically partly or completely micritized. These rocks are similar to crystalline limestone in the Eli Limestone, and are interpreted as recrystallized skeletal grainstone.

Spiculitic Limestone

Spiculitic limestone makes up most of the upper part of the Kogruk Formation north of Kivivik Creek. A pink or reddish weathering tint, a dark gray to black fresh surface, and rhythmic bedding are characteristic of these rocks and distinguish them from other Kogruk lithologies. Beds are generally thin (0.5-6 cm), regular, internally laminated, and separated by shaley partings (Fig. 23B). Chert content is variable but reaches 50 percent in some outcrops; it generally occurs as even layers a few cm thick that are more regular in shape, consistent in thickness, and persistent laterally than chert elsewhere in the Kogruk Formation.

Most rocks are fine-grained packstone or lesser wackestone and are made up chiefly of calcareous and siliceous sponge spicules and subordinate peloids and other bioclasts (Fig. 24). The original composition of spicules in these samples is uncertain, as siliceous and calcareous spicules are readily replaced by calcite and silica, respectively (Horowitz and Potter, 1971). Some spicules exhibit an internal canal in cross section; most appear to consist of a single ray. Peloids are small (40-100 μm); other bioclasts include poorly preserved radiolarians?, echinoderm debris, and calcispheres.



Figure 21. Black chert nodules in bioclastic limestone, Kogruk Formation, Maiyumerak Mountains, locality 1, Fig. 2.




Figure 22. Photomicrographs of skeletal wackestone, Kogruk Formation; scale bars equal 0.5 mm. A. Echinoderm fragment (C) and bryozoa (B), locality I, Fig. 2. B. Foraminifer (F), locality M, Fig. 2.

of the Kogruk Formation indicates that a deeper water environment prevailed by late Meramecian or early Chesterian time; this facies change coincides with a regional drowning of carbonate platforms that will be further discussed below.

SUMMARY OF THE MAIYUMERAK MOUNTAINS CARBONATE SEQUENCE

The sedimentary sequence that is exposed along the west side of the Maiyumerak Mountains consists of dominantly shallow-water carbonate rocks of Middle and Late Devonian and Early and Late Mississippian age. Diagnostic faunas from these rocks are mostly conodonts but include corals, stromatoporoids, and brachiopods, and are of Givetian, Frasnian, Famennian, Kinderhookian, Osagean, and Meramecian age; Eifelian (and older?) and Chesterian rocks may also be present. Any hiatus in this succession is less than a stage in duration.

The Maiyumerak sequence accumulated in a range of open middle shelf to more restricted inner shelf environments. Shallow-water conditions existed throughout deposition of the Baird Group in the Middle Devonian; sedimentary structures characteristic of intertidal and supratidal settings and faunas suggestive of restricted circulation are particularly well-developed in the uppermost (Givetian) beds. A transgression occurred in latest Givetian-early Frasnian time and resulted in deposition of the lower, more argillaceous part of the Eli Limestone; open circulation, moderate to shallow water, and periodic input of dominantly clay-sized clastic material persisted during formation of the upper (Famennian) part of this unit.

Relatively open, moderate- to shallow-water conditions also prevailed throughout deposition of the Utukok and Kogruk Formations in Kinderhookian, Osagean, and Meramecian time; well-washed skeletal grainstone (and arenite in the Utukok Formation) formed in local high-energy shoals. Siliciclastic detritus, largely fine-grained quartz and subordinate chert sand grains, occurs sporadically throughout the Utukok Formation and is most abundant in the upper (Osagean) part of the unit, but is exceedingly rare in the overlying Kogruk Formation. Possible sources of this siliciclastic input are discussed below. Spiculitic limestone in the uppermost (Meramecian-Chesterian?) part of the Kogruk Formation suggests an outer shelf to basinal setting; this is a deeper water depositional environment than that indicated for any other part of the Maiyumerak Mountains carbonate sequence, and marks the demise of the Maiyumerak carbonate platform.

Several recent models relate the development of specific constellations of carbonate facies to global sea-level changes, generally within the context of seismic stratigraphy (e.g., Sarg, 1988; Bond and others, 1989). The complex present-day tectonic setting of northwestern Alaska (foreland fold-and-thrust belt) and the virtual absence of seismic data from this region preclude direct application of these models to the Maiyumerak carbonate succession. However, possible eustatic influences on carbonate facies within the Maiyumerak succession can be noted and are summarized here.

The most recently published eustatic curves for the Paleozoic (Ross and Ross, 1988) indicate long-term sea level generally rose from the Middle Devonian through the Early Mississippian, then declined gradually through most of the rest of the Mississippian. Thus, the Baird Group was deposited during a period of relatively low sea level, the Eli Limestone and Utukok Formation during a time of generally rising sea level, and the Kogruk Formation during a period of overall sea level decline. Short-term pulses of eustatic change are superimposed on these long-term trends, however, and include several pronounced cycles of sea-level fall and rise in the Osagean and Meramecian.

In general, carbonate facies in the Maiyumerak sequence are consistent with this suggested eustatic curve. A short-term sea-level fall inferred for the early and middle Givetian correlates with the upper part of the Baird Group, which includes the shallowest depositional environments (supratidal

and intertidal) that have been documented within the Maiyumerak sequence. Facies in the Eli Limestone and Utukok Formation, deposited during a period of generally rising sea level, indicate deeper water conditions. These units include abundant intervals of nodular to thin-bedded argillaceous limestone; such lithologies, rich in carbonate and (or) siliciclastic mud and relatively slowly deposited, are common on incipiently drowned platforms (Read, 1985) and have been described as "catch-up" carbonate systems by Sarg (1988). Particularly fast pulses of eustatic rise appear to have occurred during the Late Devonian (Kendall and Schlager, 1981) and would be expected to inhibit carbonate platform growth; carbonate production on the Maiyumerak platform may have been further suppressed by turbidity due to clastic input from the Endicott delta (discussed below).

Deposition of the upper part of the Utukok Formation and most of the Kogruk Formation coincide with a period of overall eustatic decline. Mixed carbonate and siliciclastic strata, notable in the upper part of the Utukok Formation, typically form during the early part of a global sea level fall (Sarg, 1988). The final inundation of the Maiyumerak platform appears to correlate with a short-term eustatic rise, but also reflects tectonic factors, such as the northward drift of Alaska (see below).

REGIONAL RELATIONSHIPS

The Maiyumerak Mountains lie within a fold and thrust belt that extends across northern Alaska, and are part of the De Long Mountains allochthon belt of Mayfield and others (1988) (Figs. 25, 26). These authors have proposed the most detailed paleogeographic and tectonic reconstruction presently available for northwestern Alaska. In their model, thrust sheets that contain similar lithologic sequences and occur at approximately the same structural level are grouped into structural units called allochthons. These allochthons, although structurally complex internally, are thought to overlie one another in a regionally consistent pattern of broad synclines parallel to the mountain front. Seven allochthons are formally named, sequentially numbered (allochthon 1 is structurally lowest), and considered to consist of fifteen sedimentary and two igneous sequences, which have also been formally named. As noted above, these sequences are not equivalent to seismic stratigraphic sequences, as defined by Mitchum (1977). Rather, the sedimentary sequences of Mayfield and others (1988, p. 146) are "distinctive column(s) of sedimentary rocks that have slightly different lithologic facies compared to coeval rocks of other sequences" that are found in other thrust sheets.

The Schwatka Mountains structural province to the south of the De Long Mountains allochthon belt (Fig. 26) consists of metasedimentary and metaigneous rocks thought by Mayfield and others (1988) to be autochthonous or parautochthonous relative to the allochthons of the De Long Mountains belt. Other workers (e.g., Mull and others, 1987; Till and others, 1988) have shown that these metamorphic rocks, like rocks of the allochthon belt, were folded and thrust during the Brooks Range orogeny and are thus not autochthonous, but the degree of displacement of the metamorphic rocks has not been established. In the following discussion, rocks of the Schwatka province are considered allochthonous, but recognition of two separate structural provinces (De Long Mountains and Schwatka Mountains) is maintained.

Carbonate rocks of the Maiyumerak Mountains are included by Mayfield and others (1988) in the Eli sequence of the Kelly River allochthon. Figure 26 shows the generalized regional extent of allochthons and sequences in northwestern Alaska as proposed by Mayfield and others (1988), and Figure 27 presents simplified stratigraphic columns for sedimentary sequences within the allochthons. Although the sequences include rocks of Devonian through Cretaceous age, only the Devonian-Pennsylvanian parts of the sequences are shown and will be considered in this report. In the discussion below, the sequences and allochthons of Mayfield and others (1988) are briefly defined, units of the Maiyumerak Mountains carbonate sequence are compared to correlative units in the allochthon belt and the Schwatka province, the position of the Maiyumerak sequence within the tectonostratigraphic




Figure 25. Central and northern Alaska, showing area of western Brooks Range discussed in this chapter. Letters identify 1:250,000 quadrangles outlined by dashed lines and referred to in text: PH, Point Hope; DM, De Long Mountains; MM, Misheguk Mountain; HP, Howard Pass; N, Noatak; BM, Baird Mountains; AR, Ambler River.




Figure 26. Distribution of allochthons and sedimentary sequences in De Long Mountains allochthon belt, northwestern Alaska, from Mayfield and others (1988). Rocks of Schwaike Mountains structural province occur in southeast quadrant of map.

Figure 27. Age, lithology, stratigraphy, and thickness of Devonian-Mississippian rocks in the western part of the Schwatka Mountains province (leftmost column) and in the sedimentary sequences of Mayfield and others (1988) in the De Long Mountains allochthon belt (all other columns). Numbers indicate allochthons of Mayfield and others (1988). Letters indicate data sources: A, Karl and others (1989); B, Armstrong (1970a, 1972) and Armstrong and others (1971); C, Campbell (1967); D, Dutro (1987); E, Mayfield and others (1988); F, Curtis and others (1983), Eilersieck and others (1983), and Mayfield and others (1983); G, Curtis and others (1984), Eilersieck and others (1984), and Mayfield and others (1984); H, Mayfield and others (1987); I, Holdsworth and Murchey (1988). Stratigraphic units: PMc, black chert, shale, limestone, and (or) dolostone; PMk, Kuna Formation; PMn, Nuka Formation; Mk, Kayak Shale; Mko, Kogruk Formation; Mkoc, cherty member of Kogruk Formation; MI, Lisburne Group, undifferentiated; Mis, Lisburne Group, shale and micritic limestone; Mml, Lisburne Group, micritic limestone; Mn, Nasorak Formation; Mu, Utukok Formation; Mt, Tupik Formation; MDe, Endicott Group; MDI, limestone (in part equivalent to Kugururok Formation in Bastille sequence); Db, Baird Group, undifferentiated; Dbd, Baird Group, dolomite; Dbl, Baird Group, limestone (includes Eli Limestone in Eli sequence); Dbs, Baird Group, limestone and shale; DI, limestone and dolomite (in part equivalent to Kugururok Formation in Bastille sequence); Dn, carbonate and clastic rocks of the Nakolik River; Dsl, sandstone, shale, and limestone (in part equivalent to Kugururok Formation in Bastille sequence); Pzqs, quartz metaconglomerate, metasandstone, and siliceous phyllite. Lithologic units that have yielded no fossils indicated by *; age estimates for these units based on age of similar units in adjacent sequences.

framework of Mayfield and others (1988) is described, and the implications of the Maiyumerak sequence for paleogeographic reconstruction of the western Brooks Range are considered.

DEFINITIONS OF SEQUENCES AND ALLOCHTHONS

Each sedimentary sequence of Mayfield and others (1988) is characterized by a distinctive succession of Devonian-Cretaceous sedimentary rocks; the sequences are believed to have been deposited contiguously within a broad epicontinental basin. A few sequences are distinguished by a single unique lithostratigraphic unit, but most are differentiated on the basis of more subtle variations in coeval lithofacies or stratigraphic successions. Facies differences between sequences within the same allochthon are less pronounced than facies differences between sequences in different allochthons; overall, facies contrasts are greatest in rocks of Mississippian age.

The Brooks Range allochthon (1) is the most extensively exposed and structurally lowest of the allochthons. Sequences of this allochthon generally include a thick interval of Upper Devonian-Lower Mississippian clastic rocks (Endicott Group); three sequences (Ivotuk, Key Creek, and Lisburne Hills) are recognized (Figs. 26 and 27). The Picnic Creek allochthon (2) includes four sedimentary sequences (Amaruk, Nigu, Picnic, and Wulik); all contain notable bedded black chert of Late Mississippian age. The Kelly River allochthon (3) is exposed only in the western part of the De Long Mountains allochthon belt. Three sequences have been recognized (Amphitheater, Eli, and Kelly); all include thick intervals of Devonian-Mississippian carbonate rocks (Baird Group, Kogruk Formation) and lack Mississippian black shale (Kayak Shale). The Ipnavik River allochthon (4) contains three sequences (Ipnavik, Nachralik Pass, and Puzzle Creek) which are similar to those of allochthon 2 but distinguished by the presence of rare to common diabase sills. The Nuka Ridge allochthon (5) is structurally highest of the allochthons composed primarily of sedimentary rocks. Two sequences (Bastille and Bogle) are recognized and contain Devonian-Carboniferous arkosic limestone (Kugurok and Nuka Formations). The Copter Peak and Misheguk Mountain allochthons (6 and 7) consist of mafic igneous rocks.

Information on rocks of the allochthon belt is available primarily in a series of mile:inch geologic maps of parts of the Misheguk Mountain, De Long Mountains, and Noatak quadrangles (Curtis and others, 1983, 1984; Eilersieck and others, 1983, 1984; Mayfield and others, 1983, 1984, 1987); these maps include summaries of fossil collections and lithologic data. Unless otherwise stated, all data below on rocks of the allochthon belt north and west of the Baird Mountains quadrangle are taken from these maps. Endothyrid foraminifer zones referred to are those of Mamet and have been described in a number of publications (e.g., Mamet and Skipp, 1970; Armstrong and Mamet, 1977).

Metasedimentary rocks of the Schwatka structural province are treated as a single stratigraphic succession, the Schwatka sequence, by Mayfield and others (1988), but it is likely that several distinct sedimentary sequences exist within this poorly understood province (Dumoulin and Harris, 1987; Dillon and others, 1987). In the discussion below, remarks on stratigraphy of the Schwatka province refer chiefly to rocks in the westernmost part of this province (the northeastern Baird Mountains quadrangle), as relatively detailed stratigraphic and structural information (Dumoulin and Harris, 1987; Karl and others, 1989; Till and others, 1988) is available from this area.

CORRELATION

Baird Group

Rocks lithologically and biostratigraphically correlative with the Baird Group in the Maiyumerak Mountains occur both northwest and southeast of the study area within the De Long Mountains allochthon belt, but have not been found in the western Schwatka Mountains province. As discussed above,

stratigraphic nomenclature of Devonian carbonate rocks in northwestern Alaska remains somewhat unsettled. The name Baird Group has been applied to a variety of carbonate rocks of Devonian and older age that occur in both the De Long Mountains and Schwatka Mountains structural provinces, but few formations or members have yet been delineated within this group. Baird Group rocks, however, show differences in internal stratigraphy and detailed sedimentology from one area and (or) structural package to another, and these differences may help constrain regional paleogeographic and tectonic reconstructions. In order to highlight this diversity, and to facilitate detailed comparisons with rocks of the Maiyumerak Mountains, specific Baird Group sequences will be treated separately below; mile:inch map unit designations and geographic references will be used as necessary to distinguish various sequences.

De Long Mountains Allochthon Belt

Rocks correlative with the Baird Group in the Maiyumerak Mountains occur in the Eli and Kelly sequences, exposed northwest of the Maiyumerak Mountains, and in a sequence of uncertain tectonic affinity that crops out through the central Baird Mountains quadrangle southeast of the Maiyumerak Mountains (Agashashok sequence of Eilersieck, 1985).

Northwestern Area Middle and (or) Upper Devonian carbonate rocks occur in all sequences of allochthon 3 and two sequences of allochthon 4 and have been assigned to the Baird Group by Mayfield and others (1988), but only the Eli and Kelly sequences of allochthon 3 contain clean carbonate rocks of probable Middle Devonian or older age. In these sequences, the base of the Baird Group is a thrust fault, and dominantly dolomitic Middle Devonian carbonate rocks are conformably overlain by locally argillaceous Upper Devonian carbonate rocks. In other sequences, rocks assigned to the Baird Group cannot, on the basis of presently available paleontologic evidence, be shown to be older than Late Devonian.

The Eli sequence in the Misheguk Mountain quadrangle contains carbonate rocks that, on the basis of published descriptions and fossil collections, are most like the Middle Devonian Baird Group in the Maiyumerak Mountains. These rocks are mapped as Db3e (Baird Group undivided) in most areas, but in the southeastern part of the quadrangle are subdivided into a lower unit, Dbd3e (dolomite), and an upper unit, Dbl3e (unnamed limestone and Eli (?) Formation) (Curtis and others, 1984). Dbd3e consists of 200 m or less of massive to thick-bedded, amphiporid-bearing dolostone, and contains corals and tentaculitids of Middle to Late Devonian age. This unit is exposed due north of the Maiyumerak Mountains along the east side of the Avan Hills, and is overlain by argillaceous carbonate rocks similar to the Eli Formation (see below). At least parts of Db3e undivided are also coeval with the Baird Group in the Maiyumerak Mountains; Middle Devonian corals occur in outcrops of Db3e along the west side of Misheguk Mountain.

The Kelly sequence in the Misheguk Mountain and De Long Mountains quadrangles also contains rocks assigned to the Baird Group that may correlate with the Middle Devonian carbonate rocks in the Maiyumerak Mountains. In the Misheguk Mountain quadrangle, map unit Db3 of Mayfield and others (1984) consists of about 100 m of medium- to thick-bedded limestone and dolostone. The lower part of this unit contains corals and stromatoporoids of probable Middle Devonian age as well as corals, stromatoporoids, and brachiopods of latest Silurian-Middle Devonian age. In the De Long Mountains quadrangle, map unit Dbl3 consists of medium- to coarse-grained limestone and is a maximum of 70 m thick; the lower part of this unit contains late Middle Devonian brachiopods as well as imprecisely dated Silurian-Devonian corals, stromatoporoids, and brachiopods (Curtis and others, 1983; Mayfield and others, 1983).

this sequence are not available, but compositional similarity to sandstone of the Endicott Group is suggested by Curtis and others (1984).

The Kugururok Formation of the Bastille sequence also correlates with the Eli Limestone, but contains a more diverse clastic component than that of the units just described. In the Baird Mountains, the Kugururok Formation consists of as much as several hundred m of interbedded limestone, dolostone, and calcareous sandstone, siltstone and shale; it is Middle (probably late Middle) to Late Devonian in age, based on conodonts, brachiopods, and foraminifers (Karl and others, 1989). Sandstone composition is variable from bed to bed, but generally consists mostly of quartz, calcareous bioclasts, and sedimentary lithic clasts; feldspar (including potassium feldspar and microcline) and volcanic lithic clasts make up 5-20 percent of some samples (Dumoulin, unpublished data).

In the Misheguk Mountain quadrangle, rocks originally designated the Kugururok Formation by Sable and Dutro (1961) are included in map units Dsl5, DI5, and MDI5 of Curtis and others (1984) and Ellersieck and others (1984). Dsl5 is equivalent to the lower part of the Kugururok Formation; it consists of 150 m or less of interbedded shale, limestone, calcareous siltstone, sandstone, and minor conglomerate and contains conodonts of Middle Devonian age. DI5 and MDI5 interfinger laterally and are equivalent to the upper part of the Kugururok Formation. DI5 is at least 300 m thick, consists of massive to thin-bedded limestone and dolostone, and contains Givetian brachiopods, corals and stromatoporoids and early or middle Late Devonian brachiopods. MDI5 is interbedded limestone and sandstone which may contain up to 15 percent potassium feldspar; it contains brachiopods and corals of Frasnian or older age and foraminifers of Famennian (zone 5) age.

Summary

Upper Devonian carbonate rocks are widely distributed in the De Long Mountains allochthon belt and are generally relatively impure. The coarsest noncarbonate detritus occurs in Upper Devonian carbonate rocks in the structurally highest and lowest sequences. Sandy limestone in the highest sequence (allochthon 5) contains quartz, feldspar, and volcanic and sedimentary lithic clasts; noncarbonate material in limestone of the lowest sequences (allochthons 1 and 2) is primarily quartz. Argillaceous carbonate rocks like the Eli Limestone in the Malyumerak Mountains, as well as clean carbonate rocks lacking a significant terrigenous component, occur in sequences of intermediate structural level; purer carbonate rocks crop out mainly in the southwestern part of the allochthon belt (western Noatak quadrangle).

Utukok Formation

Impure carbonate rocks correlative with the Lower Mississippian Utukok Formation in the Malyumerak Mountains occur in six of the fifteen sedimentary sequences in the De Long Mountains allochthon belt (Mayfield and others, 1988). Lower Mississippian rocks in most of the other sequences consist of a thin interval of black shale and subordinate limestone and sandstone, the Kayak Shale. Impure carbonate rocks or black shale of Early Mississippian age have not been recognized in western exposures of the Schwatka province (Baird Mountains quadrangle).

De Long Mountains Allochthon Belt

Impure Carbonate Rocks The Utukok Formation is widely exposed throughout the Misheguk Mountain, De Long Mountains and Noatak quadrangles in the western part of the De Long Mountains allochthon belt, but has not been recognized east of the western Howard Pass quadrangle. The formation occurs in sequences of allochthons 1 (Key Creek sequence), 2 (Amaruk and Wulik sequences), 3 (Eli and

Kelly sequences), and 4 (Puzzle Creek sequence), but it is thickest and most widely distributed in the Kelly sequence of allochthon 3. It is discontinuously present and less than 50 m thick in most sequences of allochthons 1 and 2, and is at most 200 m thick in allochthon 4. In contrast, thicknesses of 1000 m are reported for the Utukok Formation in the Kelly sequence, both at the type area in the Misheguk Mountain quadrangle (Sable and Dutro, 1961), and to the west in the De Long Mountains quadrangle (Mayfield and others, 1983). However, the unit appears to be less than 100 m thick in most of the Eli sequence and may be locally absent (Mayfield and others, 1988).

In addition to striking intersequence differences in thickness, the Utukok Formation displays notable vertical and lateral variations in clastic content. The unit includes more shale and less sand in structurally higher sequences and in the more westerly exposures of all sequences. For example, in the De Long Mountains quadrangle, estimates of shale content in the Utukok Formation range from 25 percent in the Key Creek sequence to 30-40 percent in the Amaruk sequence to 50 percent in parts of the Kelly sequence (Mayfield and others, 1983), and the Utukok Formation in all sequences in the De Long Mountains and Noatak quadrangles is shalier than in the Misheguk Mountain and Baird Mountain quadrangles. In contrast, interbeds of clean quartz sandstone in the Utukok Formation are reported mostly from structurally lower sequences and more easterly exposures.

Most fossil collections from the Utukok Formation are of Early Mississippian age; some may be as old as latest Devonian (Famennian) or as young as early Late Mississippian (Meramecian). The unit contains a diverse megafauna of crinoids, brachiopods, corals, gastropods, bryozoans, and pelecypods, but the tightest age determinations are based on microfossil assemblages of foraminifers and conodonts. Foraminifers representative of zones 7, 8, 9, 10 or 11, 11, and 12 indicate ages equivalent to late Kinderhookian through early Meramecian; a more limited number of conodont assemblages are of Famennian-early Kinderhookian, late Kinderhookian, and early-middle Osagean ages. Data at present are insufficient to prove regional or intersequence variations in the age of the Utukok Formation, but the youngest tightly diagnostic faunas, foraminifers of early Meramecian (zones 11 and 12) age, have been obtained from the Kelly sequence in all areas, and from westernmost exposures of the Key Creek and Eli sequences in the De Long Mountains and Noatak quadrangles, respectively.

Black Shale The other widely distributed unit of Early Mississippian age in the western Brooks Range is the Kayak Shale, uppermost formation of the Endicott Group. The formation was defined in the central Brooks Range by Bowsher and Dutro (1957), but has been recognized throughout the Brooks Range and in the subsurface in northern Alaska (e.g., Armstrong and Bird, 1976). In the sequences of the De Long Mountains allochthon belt, the Kayak Shale is generally less than 100 m thick and consists mostly of black to dark-gray shale; it contains beds of sandstone and siltstone in its lower part and thin layers of bioclastic argillaceous limestone in its upper part (Mayfield and others, 1988).

The Kayak Shale is part of the transgressive suite of Armstrong and Bird (1976). Sections of Kayak Shale in the western Brooks Range appear to have been deposited in progressively deepening water; the basal sandstone beds have sedimentary features indicative of tidal-flat deposits, but thin graded sandstone beds higher in the unit are turbidites and (or) storm deposits, and rare limestone beds in the upper part of the formation are interpreted as debris flows (Nilsen, 1981; Moore and Nilsen, 1984). Shale in this unit is characteristically black and is generally described as carbonaceous, but little geochemical data is available so total organic carbon content is uncertain (Bird and Jordan, 1977; Magoon and Bird, 1988).

The age of the Kayak Shale in the western Brooks Range is relatively poorly constrained but appears to be chiefly Early Mississippian. Many sections have not been dated or yield only long-ranging Mississippian or Early Mississippian forms (e.g., Mayfield and others, 1983, 1984); a few widely-

scattered collections of goniatites and conodonts are of definitive Kinderhookian and Osagean age (Dutro, 1987; Karl and others, 1989). The unit may not be coeval in all allochthons and geographic areas; Meramecian fossils have been reported from sections in the Noatak quadrangle (Mayfield and others, 1987).

In general, sequences in the allochthon belt that contain the Utukok Formation do not contain Kayak Shale. Where both formations are recognized in the same sequence, the Kayak Shale either overlies the Utukok Formation, as in the Key Creek and Wulik sequences, or interfingers with it, as in the Amaruk and Puzzle Creek sequences. The depositional relationship between these two units is further discussed below.

Summary

Light-colored sandy limestone assigned to the Utukok Formation is widely distributed in the De Long Mountains allochthon belt; it occurs in some sequences of allochthons 1, 2, 3, and 4. The unit is thicker and contains more terrigenous detritus in the Kelly sequence and in more westerly exposures of all sequences. Lower Mississippian rocks in sequences that do not contain the Utukok Formation generally consist of thin intervals of black shale and subordinate sandstone and limestone (Kayak Shale).

Kogruk Formation

The Kogruk Formation has been recognized in sequences of allochthons 1, 2, and 3 in the De Long Mountains allochthon belt (Mayfield and others, 1988), and in the Schwatka province in the northeastern Baird Mountains quadrangle (Karl and others, 1989). Most of these rocks correlate well both biostratigraphically and lithologically with rocks referred to the Kogruk Formation in the Malyumerak Mountains.

De Long Mountains Allochthon Belt

Sequences of allochthons 1 (Key Creek and Lisburne Hills), 2 (Amaruk), and 3 (Amphitheater, Eli, and Kelly) all contain carbonate rocks assigned to the Kogruk Formation (Mayfield and others, 1988); occurrences of this unit in the Lisburne Hills, however, are considerably younger than sections of this formation elsewhere, and will be discussed separately below. Like the Utukok Formation, the Kogruk Formation occurs only in the western part of the allochthon belt and displays pronounced intersequence and intrasequence variations in thickness. It is less than 30 m thick and only discontinuously present in sequences of allochthons 1 and 2 (exclusive of the Lisburne Hills sequence), but as much as 700 m thick in some sections of allochthon 3. Within individual sequences, the unit is thicker to the west and perhaps northwest. For example, the Kogruk Formation in the Kelly sequence is more than 300 m thick in the southwest part of the Misheguk Mountain quadrangle, but less than 30 m thick 60 km to the east (Curtis and others, 1984; Mayfield and others 1984). Greatest thicknesses of the formation in all sequences are reported from the De Long Mountains quadrangle (Mayfield and others, 1983).

Intersequence and intrasequence lithologic trends are also apparent. Corals (particularly lithostrotionids) are abundant in the Kogruk Formation only in the sequences of allochthon 3. Layers and nodules of black chert are most abundant in the Kogruk Formation in the sequences of allochthons 1 and 2 and in western occurrences of all sequences.

Only a few sections of the Kogruk Formation, in the Kelly and Eli sequences, have been studied in detail (Sable and Dutro, 1961; Armstrong, 1970a). These sections consist mainly of bryozoan-

echinoderm packstone and wackestone with subordinate intervals of lime mudstone (composed of dotted pellets) and ooid and bioclastic grainstone and packstone. Dolomite occurs mainly as small disseminated rhombs that make up 1-15 percent of the lime mudstone intervals. Nodular to lenticular chert is found in all sections but varies markedly in abundance both vertically and laterally; it is most abundant in mudstone and wackestone. Armstrong (1970a) concluded that the Kogruk Formation was deposited in an open-marine environment in which carbonate deposition and subsidence were near equilibrium; shoal conditions were locally achieved but sedimentary features indicative of intertidal and supratidal environments are absent.

Rocks similar to the thin-bedded spiculitic limestone in the upper part of the Kogruk Formation in the Maiyumerak Mountains have been described from all three sequences of allochthon 3. In the Misheguk Mountain quadrangle, these rocks occur in the Kelly and Eli sequences and have been referred to the Tupik Formation (Mayfield and others, 1984); in the De Long Mountains and Noatak quadrangles, they occur in the Kelly and Amphitheater sequences respectively, and are assigned to map unit Mkoc3 (Mayfield and others, 1983, 1987). All occurrences are less than 75 m thick, form the upper part of or overlie the Kogruk Formation, and consist of dark-gray to black limestone thinly interbedded with as much as 50 percent chert.

Fossils are relatively abundant in the Kogruk Formation, and the age of this unit in the allochthon belt is well constrained by conodonts, corals, and foraminifers. The base of the unit in most sections is early Meramecian and contains foraminifers representative of zones 11 or 12, but in some parts of the Eli sequence (for example, the Noatak quadrangle), late Osagean ages (zone 10) are reported (Mayfield and others, 1987). The youngest definitive ages, late Meramecian (zone 14 or 15), have been obtained from the Kelly sequence in all areas, and the Eli sequence in the Misheguk Mountain quadrangle. The Tupik Formation and equivalent cherty limestone units are less well-dated; sparse fossil collections include foraminifers of late Meramecian (zone 14 or 15) age, and nautiloids (Gordon, 1957) considered by Dutro (1987) to indicate an early Chester age.

Lisburne Hills In the Point Hope quadrangle on the Lisburne Peninsula, rocks referred to the Kogruk Formation were studied by Armstrong (1970a, 1972) and Armstrong and others (1971) and rocks designated Kogruk? Formation were described by Campbell (1967). The structural position of these rocks is uncertain; they have been included in the Lisburne Hills sequence of allochthon 1, but Carboniferous stratigraphy in the area is most like that in the sequences of allochthon 3 (Mayfield and others, 1988).

Several partial sections of Kogruk Formation, 70 to 600 m thick, were examined by Armstrong (1970a, 1972) and Armstrong and others (1971) in both northern and southern parts of the peninsula. These sections consist mostly of bryozoan-echinoid packstone to wackestone, locally cherty and (or) dolomitized; some intervals include shallowing-upward cycles 6-40 m thick of bioclastic grainstone or packstone capped by cherty dolomitic mudstone. The rocks contain foraminifers representative of Meramecian to Chesterian age (zones 13 through 18). Campbell (1967) describes several thoroughly dolomitized sections as much as 1400 m thick from the southern seacliffs of the peninsula that consist mostly of variously fossil-rich biomicrites and are of Chesterian age. The uppermost part of the Kogruk Formation on the Lisburne Peninsula contains large brachiopods that may be of Early Pennsylvanian (Morrowan) age (Dutro, 1987). Thus, the Kogruk Formation on the Lisburne Peninsula appears to be similar in lithology but notably younger than the Kogruk Formation elsewhere within the allochthon belt.

Schwatka Mountains Structural Province

Rocks referred to the Kogruk Formation in the Schwatka province crop out in the northeastern

Baird Mountains quadrangle; these rocks have been described by Karl and others (1989) and were further analyzed for the present study. They are similar in age and general lithology to the Kogruk Formation in the Maiyumerak Mountains, but occur in a distinctly different stratigraphic context and include sedimentary features suggestive of very shallow water (supratidal or intertidal) depositional environments. The section is 30 m thick and overlies undated siliceous metaclastic rocks that may be metamorphosed equivalents of the Endicott Group; these rocks in turn disconformably(?) overlie massive metacarbonate rocks that contain conodonts that are no younger than late Early Devonian (Emsian) and may be as old as Middle Silurian. All lithologic contacts in this area have probably been somewhat structurally disrupted, but the general stratigraphic succession at this locality appears to be relatively intact.

The Kogruk Formation in the northeastern Baird Mountains quadrangle consists of dolostone and lesser limestone with locally abundant black chert nodules and lenses. Some original textures are well preserved and indicate that much of the section was deposited as bryozoan-echinoderm wackestone and packstone; lithostrotionid corals occur locally. Several intervals are rich in sponge spicules; these spiculitic rocks contain laminar to hummocky cryptalgal layering and appear to have formed in very shallow, restricted environments. The section contains the conodont *Hindeodus scitulus* (USGS colln. 28976-PC) indicative of late Osagean to early Chesterian age and a shallow-water setting, and the coral *Lithostrotionella* aff. *L. birdi*, of middle Meramecian to early Chesterian age (equivalent to foraminifer zones 13-16) (written commun., A. K. Armstrong, 1978).

Summary

Carbonate rocks lithologically and biostratigraphically correlative with the Kogruk Formation in the Maiyumerak Mountains occur in sequences of allochthons 1, 2, and 3 in the De Long Mountains allochthon belt, and in the western part of the Schwatka province. In most sequences of the allochthon belt, with the exception of the Lisburne Hills sequence, the Kogruk Formation consists chiefly of bryozoan-echinoderm wackestone and packstone, locally dolomitic and (or) cherty, deposited in open marine conditions. Sections are thicker to the west and northwest and are late Osagean or early Meramecian to late Meramecian in age. Most sections are capped by thin-bedded cherty limestone of late Meramecian or early Chesterian age deposited in a deeper water setting. The Kogruk Formation in the Lisburne Hills is locally completely dolomitized, includes distinct shallowing-upward cycles, and is Meramecian to Chesterian or Morrowan in age. In the western part of the Schwatka province, the Kogruk Formation was deposited in shallow to very shallow water and is imprecisely dated; the base may be as old as late Osagean and the top may be as young as early Chesterian.

Correlation Summary and Discussion

As is evident from the discussion above and the generalized sections in Fig. 27, a number of distinct sedimentary facies coexisted during middle to late Paleozoic time in what is now northwestern Alaska. This lithologic diversity is thought to reflect a complex Devonian-Carboniferous paleogeography of shallow platforms and intracratonic basins in which allochthonous quartzose and quartzofeldspathic material and autochthonous carbonate sediment accumulated (Mayfield and others, 1988). Lithologic and stratigraphic features of sedimentary sequences in the De Long Mountains allochthon belt and in the western Schwatka province are summarized below, and compared to those of carbonate rocks in the Maiyumerak Mountains.

The Schwatka province and 11 of the 15 sequences in the allochthon belt contain rocks of confirmed or probable Devonian age (Mayfield and others, 1988) (Fig. 27). No rocks definitively older than late Middle Devonian (Givetian) have been found in most of the allochthon belt; in contrast, the

western part of the Schwatka province contains no Devonian rocks known to be younger than late Early Devonian (Karl and others, 1989). All Devonian rocks in the allochthon belt were deposited in relatively shallow water and can be grouped into three broad categories: carbonate rocks with little or no sand-sized siliciclastic component (Baird Group, Eli Limestone), siliciclastic rocks with little carbonate component (Endicott Group), and quartzofeldspathic calcareous rocks (Kugururok Formation); facies transitional between the first two categories have also been recognized. Devonian rocks in sequences of allochthon 1 and 2 are siliciclastic rocks (Amaruk, Picnic Creek sequences) or mixed carbonate and siliciclastic rocks (Key Creek, Wulik, Nigu sequences). Devonian rocks in all sequences of allochthon 3 and 4 are relatively pure limestone; argillaceous limestone occurs in some parts of the Eli, Kelly, and Puzzle Creek sequences. Clastic rocks that contain greater than 10 percent feldspar grains have been recognized only in allochthon 5 (Bastille sequence).

The Schwatka province and all sequences of the allochthon belt with the exception of the Bastille sequence contain rocks of confirmed Mississippian age. Both Lower and Upper Mississippian rocks comprise several distinct facies. The major Lower Mississippian units are dark shale and lesser sandstone and limestone (Kayak Shale, uppermost unit of the Endicott Group) and limestone, sandy limestone, sandstone, and (or) shale (Utukok Formation of the Lisburne Group). In some sequences, most notably the Key Creek sequence, coarse siliciclastic rocks of the Endicott Group (Noatak Sandstone and (or) Kanayut Conglomerate) are of Early Mississippian age (Mayfield and others, 1984). Most sequences of the allochthon belt contain the Kayak Shale; the Utukok Formation occurs primarily in the sequences of allochthon 3. Endicott Group undivided is recognized in the Lisburne Hills and Nigu sequences; the Amphitheater sequence contains an interval of interbedded shale and limestone assigned to the Lisburne Group but not to the Utukok Formation (Fig. 27).

Upper Mississippian rocks in sequences of the allochthon belt consist of thick accumulations of light-colored carbonate rocks formed in relatively shallow water environments (Kogruk Formation, Lisburne Group undivided in the Ivotuk sequence), and (or) generally thinner intervals of dark limestone, shale and (or) chert deposited in deeper water settings (Tupik and Kuna Formations, Akmalik Chert, and various units informally referred to as "black Lisburne") (Fig. 27). "Kogruk-type" neritic carbonate rocks comprise relatively thick accumulations in all sequences of allochthon 3 and most sequences of allochthon 1, and thinner deposits in the Key Creek and Amaruk sequences and in at least parts of the Schwatka province. These rocks are overlain in most sequences by a thin interval of "black Lisburne" facies, generally the Tupik Formation. Upper Mississippian rocks in most sequences of allochthons 2 and 4, on the other hand, consist only of one or more of the "black Lisburne" facies, and in the Bogie sequence of allochthon 5 consist of feldspathic carbonate rocks (Nuka Formation).

The complete succession of Devonian-Mississippian carbonate rocks in the Maiyumerak Mountains is most similar to Devonian-Mississippian sections in the sequences of allochthon 3 (Kelly River allochthon) that are found throughout the Noatak, De Long Mountains, and Misheguk Mountain quadrangles to the north and west (Fig. 26). Thus, on sedimentologic and biostratigraphic grounds, the inclusion of the Maiyumerak sequence with sedimentary sequences of allochthon 3 proposed by Mayfield and others (1988) seems warranted.

Other sequences in the De Long Mountains allochthon belt include deeper water basinal deposits and (or) siliciclastic rocks of Devonian-Mississippian age. Only the sequences of allochthon 3 preserve a record of relatively continuous shallow-water carbonate deposition from at least Middle Devonian into Late Mississippian time, and even these sequences contain evidence of some clastic input during Late Devonian and Early Mississippian time. The lithologic and biostratigraphic similarities between Paleozoic rocks in all the sequences of the Kelly allochthon suggest that these sequences were once contiguous, and formed originally as part of a single carbonate platform. This hypothetical paleogeographic feature, which includes carbonate rocks of the Maiyumerak sequence, is here called the

Kelly platform. In the following section, implications of the Maiyumerak Mountains sequence for paleogeographic models of the northwestern Brooks Range are explored.

PALEOGEOGRAPHY

Paleogeographic Model

The prevailing model for the Paleozoic paleogeography of the northwestern Brooks Range is that of Mayfield and others (1988). Other authors (e.g., Churkin and others, 1979) have outlined alternate paleogeographic and tectonic reconstructions of northwestern Alaska, but the model of Mayfield and others (1988) is the only one presented in sufficient detail to permit precise stratigraphic and sedimentologic inferences to be drawn and tested. These authors propose a palinspastic synthesis in which the relative movement of all thrust sheets is north under south. Rocks of the Schwatka province, considered autochthonous or parautochthonous, are restored farthest north, igneous rocks of allochthons 6 and 7 are restored farthest south, and the sedimentary sequences of allochthons 1 through 5 are restored sequentially between them.

The Devonian-Mississippian paleogeography that results from this restoration is a complex series of platforms and intracratonic basins interpreted by Mayfield and others (1988) as follows (Fig. 28). Rocks of the Schwatka province are thought to have been an emergent land mass during much of middle Paleozoic time, and another continental mass ("Nukaland") is postulated to have existed south of the sequences of allochthon 5. Sequences of allochthon 1, which include thick sections of siliciclastic rocks of the Endicott Group, were deposited on a shallow shelf flanking the Schwatka continental mass and were derived from a source to the northwest. Sequences of allochthon 5, also shallow-water deposits, contain feldspar and volcanic lithics derived from Nukaland. Sequences of allochthon 3 are envisioned to have formed a topographic high (Kelly platform of this report) that was flanked by basins to the north and south; deposits of these basins are represented by sequences of allochthons 2 and 4, respectively.

It is beyond the scope of this paper to present a detailed structural critique of the model of Mayfield and others (1988), but some questions may be raised concerning the original position of rocks in the Schwatka province. As noted above, rocks of this province have been folded and thrust during the Brooks Range orogeny, and thus are not autochthonous (Till and others, 1988). In addition, if metamorphic rocks of the Schwatka province are restored farthest north, and igneous rocks of allochthons 6 and 7 are restored farthest south, present-day juxtaposition of these rocks (Fig. 26) is difficult to explain. Finally, metamorphism of rocks in the Schwatka province is attributed to underthrusting beneath more than 10 km of thrust sheets and is postulated to have occurred during the later stages of the underthrusting event. However, the high-pressure metamorphism of rocks in the Schwatka province took place during late Jurassic-Early Cretaceous time and is coeval with the oldest well-dated flysch deposits produced during thrusting (Armstrong and others, 1986; Till and others, 1988).

Paleogeographic Implications of Maiyumerak Sequence

Comparison of the Devonian-Mississippian carbonate sequence in the Maiyumerak Mountains to correlative sequences in the allochthon belt and within the Schwatka province constrains the paleogeographic model of Mayfield and others (1988) described above. Specific problems which the Maiyumerak data illuminate include the relationship between Paleozoic carbonate platforms in the allochthon belt and the Schwatka province, the source of the clastic material in the Maiyumerak sequence, the relationship between Paleozoic carbonate platforms and the Endicott clastic wedge, and the timing and cause of the demise of the Kelly platform.




Figure 28. Schematic paleogeographic map of northwestern Alaska in Late Devonian-Early Mississippian time, based on palinspastic reconstruction of Mayfield and others (1988). Reconstruction restores each lower-numbered allochthon northward from under the adjacent allochthon to the south. Thrusting direction is assumed to have followed an arc with the pole of rotation centered in the Mackenzie River Delta area in northwestern Canada. Allochthons as in Fig. 26; geographic features from Fig. 26 are S, Schwatka Mountains, M, Maiyumerak Mountains, and N, Nuka Ridge.

Relationship Between Paleozoic Carbonate Platforms

Relatively extensive sequences of early and middle Paleozoic metacarbonate rocks occur throughout the Schwatka structural province and appear to represent the deformed remains of a Paleozoic carbonate platform or series of platforms that may extend more than 800 km through the Brooks Range (Dumoulin and Harris, 1987). The relationship of these older carbonate platform sequences to the oldest carbonate rocks of the allochthon belt is uncertain (Mayfield and others, 1988). Thus, a comparison of the Baird Group in the Maiyumerak Mountains with adjacent sequences of Devonian carbonate rocks in the Baird Mountains quadrangle provides some insight into middle Paleozoic paleogeography.

The oldest well-dated part of the Baird Group in the Maiyumerak Mountains is at least as old as early Givetian and could be as old as early Eifelian. The youngest part of the Baird Group in the central Baird Mountains quadrangle (Agashashok sequence of Ellersieck, 1985) is at least as young as Eifelian and could be, in part, Givetian. Middle Devonian rocks in both sequences consist of dolomitic bioclastic limestone containing local coral and stromatoporoid biostromes. Thus, these successions overlap in lithology and probably in age. This correspondence suggests that the Baird Group in the Maiyumerak Mountains was once depositionally contiguous with the Baird Group in the central Baird Mountains quadrangle but has since been structurally dismembered, and that on sedimentologic and stratigraphic grounds, the Agashashok sequence should be grouped with other sequences of allochthon 3. If this correlation is correct, the Kelly carbonate platform existed from at least Early Ordovician (Dumoulin and Harris, 1987) through early Late Mississippian time.

However, Middle-Upper Devonian carbonate and clastic rocks (limestone of the Nakolik River, unit Dnl of Karl and others, 1989) lie directly north of but structurally below Ordovician-Devonian carbonate rocks of the Agashashok sequence (Karl and others, 1989) (Fig. 1). Dnl is gradationally overlain by rocks of the Endicott Group and both units are included by Mayfield and others (1988) in the Key Creek sequence of allochthon 1. Karl and others (1989) suggest that a stratigraphic contact originally existed between Dnl and rocks of the Agashashok sequence. If this hypothesis is correct, the Agashashok sequence is actually part of allochthon 1, and Upper Devonian rocks deposited above the Baird Group in the central Baird Mountains quadrangle are much richer in siliciclastic detritus than are coeval rocks in the Maiyumerak Mountains. Such discordance in Upper Devonian facies suggests that the Agashashok and Maiyumerak sequences were not directly contiguous during middle Paleozoic time, and that the Agashashok sequence is not part of the Kelly carbonate platform.

The Baird Group in the Maiyumerak sequence cannot easily be compared with carbonate sequences in the western part of the Schwatka province (northeastern Baird Mountains quadrangle), because no rocks as young as Middle Devonian have been recognized in these sequences. Middle and Upper Devonian rocks may have been eroded or structurally removed from the Schwatka province, or may never have been deposited. It is possible that at least parts of the Schwatka province were emergent through Middle and (or) Late Devonian time. Much of northern Alaska (i.e., parautochthonous sequences in the northeastern Brooks Range and autochthonous sequences in the subsurface north of the Brooks Range) shows evidence of Devonian uplift and (or) regression; in these areas, Mississippian clastic rocks unconformably overlie Proterozoic or early Paleozoic sedimentary and metasedimentary rocks (e.g., Brosgé and Tailleux, 1970). This unconformity has been attributed to regional tectonism, the Ellesmerian orogeny (Lerand, 1973; Smith, 1987) but could at least in part reflect eustatic changes; evidence of Givetian regression has been noted in western Canada (Fischbuch, 1968), and Johnson and others (1985) and Ross and Ross (1988) proposed a worldwide fall in sea level during the early-middle Givetian.

Although there is no physical evidence of unconformity within the Maiyumerak carbonate

sequence, and fossils indicate that any hiatus in deposition of this sequence was less than a stage in duration, very shallow water depositional environments are interpreted for the middle to late Givetian part of the sequence. Sedimentologic features of the upper part of the Baird Group in the Maiyumerak Mountains indicate that intertidal and supratidal environments prevailed across at least the Maiyumerak part of the Kelly platform near the end of Middle Devonian time; these are the shallowest water conditions that have been documented for any interval within the Maiyumerak section.

Source of Clastic Detritus in the Maiyumerak Sequence

The Endicott Group consists of as much as 4300 m of Upper Devonian-Lower Mississippian fluviodeltaic deposits that crop out for at least 950 km along the crest of the Brooks Range (Nilsen, 1981; Moore and Nilsen, 1984). Petrographic data and stratigraphic and spatial relationships suggest that most of the clay- to sand-sized noncarbonate material in the Maiyumerak sequence was derived from the same source that produced the Endicott Group; the present-day location of this source is uncertain.

In northwestern Alaska, the Endicott Group is exposed in a series of thrust plates and comprises four formations. The lowest unit, the Hunt Fork Shale, consists largely of fine detritus and is thought to have been deposited in a marine slope to outer shelf setting. The overlying Noatak Sandstone and Kanayut Conglomerate contain coarser siliciclastic material and formed as shallow-marine and fluvial deposits respectively. The uppermost part of the group, the Kayak Shale discussed above, is dominantly fine-grained and was deposited in a progressively deepening marine environment.

The Endicott Group in the western Brooks Range is roughly coeval with the most clastic-rich part of the Maiyumerak sequence. The Hunt Fork Shale contains brachiopods, cephalopods, and corals of Late Devonian age in the Misheguk Mountain quadrangle (Ellersieck and others, 1984), and brachiopods of late Frasnian or early Famennian age in the Baird Mountains quadrangle (Karl and others, 1989). Sandier parts of the Endicott Group are less well-dated; particularly poorly constrained is the age of the uppermost coarse clastic rocks in the Kanayut Conglomerate. In the Misheguk Mountain quadrangle, the Noatak Sandstone and Kanayut Conglomerate (undivided) contain brachiopods of Late Devonian to Early Mississippian age; in the Howard Pass quadrangle, Kanayut Conglomerate contains Mississippian plant fossils (Mayfield and others, 1984). Fossils from the overlying Kayak Shale chiefly indicate a broad Early Mississippian or Mississippian age and thus do not tightly limit the age of the top of the Kanayut.

Little detailed information is available on the composition of shale in the Endicott Group, but the coarser siliciclastic rocks have been broadly characterized. Conglomerate and sandstone in the Kanayut Conglomerate in the central and eastern Brooks Range are compositionally mature. Sandstone is well-sorted, subrounded, and consists of 30-75 percent quartz, 10-50 percent chert, less than 20 percent lithic grains, including argillite, quartzite, granite, and gneiss, and minor tourmaline and opaque grains; feldspar is generally absent (Moore and Nilsen, 1984; Brosgé and others, 1988). Karl and others (1989) report 90 percent quartz and chert and 10 percent argillite clasts in metasandstone of the Kanayut Conglomerate in the Baird Mountains quadrangle. Little variation in composition with respect to stratigraphic or geographic position has been noted in the Kanayut Conglomerate and the unit is thought to have been derived from a single major source terrane (Brosgé and others, 1988).

Paleocurrent and clast-size data indicate that the Endicott Group was derived primarily from the east and north (Nilsen, 1981); clast size maxima suggest that the depositional basin was fed by two major trunk streams in the east-central Brooks Range and a smaller stream in northwest Alaska. In the western Brooks Range, clast size in Kanayut Conglomerate declines southward from a maximum along the northwestern edge of the outcrop belt in the De Long Mountains quadrangle, and paleocurrent measurements west of 159° indicate transport to the southwest (Moore and Nilsen, 1984).

Determination of the source of the Endicott Group is complicated by the allochthonous nature of the unit in the central and western Brooks Range and by uncertainty concerning the proper palinspastic restoration of northern Alaska with respect to the rest of North America. Nilsen (1981) noted that the Endicott Group more closely resembles Devonian redbeds of the Franklinian orogen in northern Canada than Upper Devonian flysch of the adjacent Cordilleran orogen to the south. Anderson (1987) analyzed composition of sandstone in the Endicott Group in the central Brooks Range, and proposed a recycled orogenic provenance for the unit. Mayfield and others (1988) suggested that the Endicott Group was derived from rocks of the Schwatka province and (or) rocks presently under the Colville Basin in the subsurface of northwestern Alaska; these areas seem unlikely to have generated a huge wedge of dominantly quartzose sediment because known pre-Middle Devonian lithologies include abundant carbonate rocks (Schwatka province) and argillite (Colville basin). However, the compositional maturity of the Endicott Group could reflect intense chemical and physical abrasion during erosion and transport; if so, its present composition may not be fully indicative of its provenance (Brosigé and others, 1988). No completely satisfactory source for the Endicott Group has yet been proposed.

The composition and age of noncarbonate detritus in the Maiyumerak sequence matches the Endicott Group. Detritus in the Maiyumerak sequence coarsens upward, from clay in the Upper Devonian Eli Limestone to sand in the Lower Mississippian Utukok Formation. The Eli Limestone correlates with the fine-grained lower part of the Endicott Group, the Upper Devonian Hunt Fork Shale, and the Utukok Formation is at least partly coeval with the coarser grained upper parts of the Endicott Group, the Upper Devonian-Lower Mississippian Noatak Sandstone and Kanayut Conglomerate. Sand-sized noncarbonate material in the Utukok Formation, like sandstone in the Endicott Group, is subrounded, compositionally mature, and consists largely of quartz, chert, subordinate muddy lithic clasts, and minor tourmaline and opaque grains. Thus, we suggest that siliciclastic detritus in the Maiyumerak carbonate sequence and in the Endicott Group have a common provenance.

One possible complication in this scenario concerns the extent of overlap in age between the sand-rich part of the Endicott Group and the Utukok Formation. As discussed above, the age of the youngest thick sandstone beds in the Endicott Group (upper Kanayut Conglomerate and lowest Kayak Shale) is poorly constrained in the western Brooks Range; it may be as young as Osagean or even Meramecian or as old as Kinderhookian. The Utukok Formation is Kinderhookian and Osagean. If the uppermost sandstone beds in the Endicott Group were indeed deposited in the Kinderhookian, quartz could have been provided to uppermost (Osagean) parts of the Utukok Formation by erosion and reworking of previously deposited Endicott Group strata.

Relationship Between Paleozoic Carbonate Platforms and Endicott Clastic Wedge

Both the sandy middle part of the Endicott Group (Noatak Sandstone and Kanayut Conglomerate) and the upper black shale unit (Kayak Shale) are at least partly coeval with the Utukok Formation in the western Brooks Range. Paleogeographic and depositional implications of this correlation are explored below.

As noted previously, several intervals of interbedded quartz sandstone and bioclastic limestone in the De Long Mountains allochthon belt have been interpreted as deposits formed by intertonguing of the Endicott clastic wedge with one or more carbonate platforms (Curtis and others, 1984; Karl and others, 1989); these intervals occur in sequences of allochthons 1 and 2. If siliciclastic material in the Maiyumerak sequence of allochthon 3 was derived from the same source as the Endicott Group, some constraints on Devonian-Mississippian paleogeographic reconstructions are implied.

In the model of Mayfield and others (1988) (see also Fig. 27 of Moore and others, in press), a

marginal sea or epicontinental basin 50 to 100 km wide is thought to have separated the Kelly carbonate platform (now preserved in the sequences of allochthon 3) and the Endicott fluviodeltaic system (preserved mostly in the sequences of allochthon 1) during Late Devonian-Mississippian time. Deposits of this intervening basin are represented by the sequences of allochthon 2 (Fig. 28).

A modern analog for this proposed paleogeography is the northeast continental margin of Australia. Here, carbonate platforms such as the Eastern and Queensland plateaus are separated from the Australian continental shelf by deep troughs which formed as rift basins (Davies and others, 1988, 1989). Terrigenous and carbonate sediments are juxtaposed and intercalated along the Australian margin; siliciclastic material derived from the continent is distributed along the inner continental shelf and carbonate reefs grow along the outer shelf and on the marginal plateaus. The outer shelf is a mixed carbonate/siliciclastic province; reefs are enclosed within terrigenous fluviodeltaic deposits and carbonate sediment occurs interbedded and intermixed with quartz sand and terrigenous mud. The plateaus, however, are relatively pure carbonate provinces; terrigenous detritus is not transported across the intervening troughs.

If the quartz sand in the Utukok Formation and in the Endicott Group was derived from the same source, a basin completely separating the Kelly platform from the Endicott delta seems unlikely; three solutions are possible. First, prevailing tectonic reconstructions may be in error, perhaps as a result of unrecognized out-of-sequence thrusting (Fig. 29A). Perhaps sequences containing carbonate platform deposits (allochthon 3) should be restored directly south of sequences containing fluviodeltaic deposits (allochthon 1); sections containing basinal deposits (allochthon 2) would then be restored south of the sequences of allochthon 3. Such a restoration has been proposed by previous workers (e.g., C. G. Mull, Alaska Department of Geological and Geophysical Surveys, written commun., 1990).

Alternatively, the basin between the Kelly platform and the Endicott delta may not have formed until the Late Mississippian, after the Utukok Formation was deposited. In this scheme, Late Devonian-Early Mississippian paleogeography of northwestern Alaska was like that shown in Figure 29A, but Late Mississippian paleogeography was like that in Figure 28. Presently available fossil data is permissive of this hypothesis but does not prove it. The oldest well-dated facies of basinal aspect in the sequences of allochthon 2 is bedded gray chert in the Nigu sequence which contains radiolarians of late Osagean to Meramecian age (Holdsworth and Murchey, 1988). However, this chert is underlain by at least 90 m of black to gray interbedded chert, shale, and limestone that also appears to have formed in a deep-water depositional environment; these rocks have yielded no diagnostic fossils, but contain poorly preserved tentaculitids? that suggest these strata are no younger than Devonian (Holdsworth and Murchey, 1988). Basinal facies in other sequences of allochthon 2 contain no definitive fossil assemblages older than Late Mississippian.

A third possibility is that the "basin" between the Kelly platform and Endicott delta was actually an embayment, open on only one end (Fig. 29B). Support for this third hypothesis is provided by geographic gradients in thickness and lithology of Devonian-Mississippian rocks in allochthons 2 and 3. These gradients suggest that, if present-day west-to-east disposition of facies is representative of their original configuration, the Kelly platform and Endicott delta were connected to the west and any embayment between them deepened to the east. Mississippian rocks in the eastern sequences of allochthon 2, the Nigu and Picnic Creek sequences, consist mostly of chert and black shale, whereas Mississippian sections in the Amarak and Wulik sequences to the west contain some platform carbonate rocks and siliciclastic detritus. Within the sequences of allochthon 3, the Utukok Formation both thickens notably and contains more clastic detritus to the west.

Thus, if siliciclastic detritus in the Maiyumerak sequence was derived from the same source as the Endicott Group, the paleogeographic model of Mayfield and others (1988) must be modified. Present




Figure 29. Alternate paleogeographic reconstructions of northwestern Alaska in Late Devonian-Early Mississippian time. Allochthons, geographic features, and symbols as in Fig. 28. **A.** Reconstruction based on ideas of C. G. Mull (written commun., 1990); sequences of allochthon 3 are restored directly south of sequences of allochthon 1. **B.** "Basin" between Kelly platform and Endicott delta is an embayment, open to the east.

data suggest that the Kelly platform was not completely cut off from the Endicott delta by a marginal sea, but only partially separated by an embayment that deepened to the east and (or) may not have formed until late Late Mississippian time. Further testing of these hypotheses requires tighter age control on cherts in the sequences of allochthon 2, and paleocurrent data from the Utukok Formation and other mixed carbonate and clastic deposits in the western Brooks Range.

The relationship between the Utukok Formation and the Kayak Shale in the western Brooks Range also has paleogeographic implications. The Utukok Formation is as much as 1000 m thick, generally light colored, and contains an abundant fauna indicative of shallow-water depositional environments. The Kayak Shale, in contrast, is typically less than 100 m thick, black to dark gray, and contains little indigenous fauna; fossils are broken, abraded, hydraulically sorted, and occur concentrated in rare limestone beds that appear to have been redeposited. Black shales like the Kayak Shale often occur near the base of transgressive sequences that blanket areas of irregular topography; these shales form at a variety of depths controlled by regional topographic restrictions on bottom-water circulation and are commonly deposited in local areas where sea floor subsidence is relatively rapid (Hallam and Bradshaw, 1979). The Kayak Shale in most sequences in the allochthon belt is succeeded by black chert and (or) siliceous shale interpreted as basinal deposits, whereas the Utukok Formation is generally overlain by fossiliferous carbonate rocks formed in shallow water. Thus, the Utukok Formation probably accumulated on topographic highs within the overall northwestern Alaska depositional basin, whereas Kayak Shale was deposited primarily in adjacent lows, subbasins, or embayments.

Demise of Kelly Platform

Thin-bedded spiculitic limestone forms the upper part of the Kogruk Formation in the Maiyumerak Mountains and appears to have been deposited in an outer shelf or basinal setting; this is a deeper water depositional environment than that indicated for any other part of the Maiyumerak sequence. Similar deeper water facies (Tupik Formation, unnamed "black Lisburne" units) overlie Upper Mississippian platform carbonate rocks in all sequences of allochthon 3 and are in turn overlain by condensed basinal deposits (chert and shale of the Pennsylvanian-Jurassic Etivluk Group) (Mayfield and others, 1988). The spiculitic limestone in the Maiyumerak Mountains is not tightly dated but is probably late Meramecian or early Chesterian in age. In other sequences of allochthon 3, the transition from shallow-water dolomitic limestone to spiculitic cherty limestone occurs in the latest Meramecian, about the level of foraminifer zone 15.

Few platform carbonate rocks younger than Meramecian have been found anywhere in the De Long Mountains allochthon belt; the exceptions are the Kogruk? Formation in the Lisburne Hills sequence (allochthon 1?) and the Nuka Formation in the Bogie sequence (allochthon 5). Both these units include neritic carbonate rocks which are at least as young as Chesterian and may be in part Pennsylvanian, and these younger carbonates differ in other ways from the Kogruk Formation in the sequences of allochthon 3. For example, the Kogruk? Formation in the Lisburne Hills contains distinct shallowing-upward cycles not found elsewhere in the Kogruk Formation, and the Nuka Formation is unique among Carboniferous carbonate rocks in northwestern Alaska in containing abundant detrital feldspar. These stratigraphic and lithologic differences suggest that carbonate rocks in the Lisburne Hills and Bogie sequences formed on discrete carbonate platforms that were not connected to the Kelly platform and that experienced separate histories of subsidence and clastic input.

Inundation of the Kelly platform probably reflects a variety of factors, both tectonic, ecologic, and eustatic. Long-term tectonic processes, such as thermal subsidence of passive margins, are an order of magnitude slower than typical sedimentation rates of neritic carbonates and thus are not usually sufficient in themselves to drown healthy carbonate platforms (Kendall and Schlager, 1981). However,

ecologic shifts may depress the growth potential of carbonate benthos to levels that can be outpaced by long-term subsidence (e.g., Davies and others, 1989). Plate motion to higher latitudes, and out of the realm most favorable to carbonate production, is one such adverse environmental change that may have affected the Kelly platform. Recent paleogeographic reconstructions (Crane, 1987; Smith, 1987) suggest that northern Alaska lay at about 40° north latitude at the start of Carboniferous time and moved progressively farther north throughout the late Paleozoic.

Paleontologic evidence indicates that Carboniferous rocks of northern Alaska were deposited in relatively cool water and supports the notion of increasingly adverse climatic conditions. Foraminifers of the Lisburne Group are characteristic of the Alaska-Taimyr faunal subrealm and are interpreted as temperate-water faunas on the basis of their low diversity, high genus/species ratios, and high number of cosmopolitan forms (Armstrong and Marnett, 1970). The Lisburne Group lacks features such as dasycladacian algal banks that are common in Carboniferous strata deposited in warm-water environments. As noted above, sedimentation rates in the Utukok and Kogruk Formations in the Maiyumerak sequence are low compared to other ancient neritic carbonate successions, and are consistent with a depositional environment only marginally conducive to carbonate accumulation. Deteriorating climate may have been responsible for the otherwise unexplained extinction, documented by Armstrong (1972), of the Lisburne Group lithostrotionid coral fauna near the end of Meramecian time but somewhat prior to cessation of neritic carbonate production. Limited persistence of latest Mississippian carbonate platform sedimentation in northwestern Alaska may reflect vicissitudes of paleolatitudinal position; although the exact paleogeographic position of the Lisburne Hills sequence is uncertain, the palinspastic model of Mayfield and others (1988) restores the Nuka sequence farther south than any other sedimentary sequence in the De Long Mountains belt.

Eustatic changes may also have played a role in the demise of the Kelly platform. Long-term global sea level fell throughout the Late Mississippian (e.g., Vail and others, 1977), but this general decline was punctuated by a number of short-term sea level rises (Ross and Ross, 1988). One such rise took place during the late Meramecian (upper part of Marnett foraminifer zone 15); this is precisely the age of the transition from shallow to deeper water carbonate facies on the Kelly platform where this transition has been most tightly dated. Rapid pulses of relative sea level rise can outpace even optimally producing carbonate platforms, but will be even more likely to inundate platforms whose growth potential is already in decline as a result of other factors, i.e., drift to higher latitudes; demise of carbonate platforms along the Atlantic margin of North America during the Early Cretaceous has been ascribed to just such a superposition of effects (Kendall and Schlager, 1981).

Thus, the Kelly platform appears to have drowned about the end of the Meramecian, and deposition of neritic carbonate rocks was never reestablished in this region. Carbonate platform sedimentation persisted into the late Late Mississippian only in limited areas of northwestern Alaska, such as adjacent to Nukaland and on the Lisburne Peninsula. The inundation of the Kelly platform at the end of Meramecian time is probably primarily attributable to northward drift of northern Alaska, although eustatic changes may also have been a factor.

CONCLUSIONS

Middle Devonian through Upper Mississippian carbonate rocks of the Maiyumerak Mountains represent essentially continuous deposition in a variety of open middle shelf to more restricted inner shelf environments. Episodic terrigenous input occurred throughout Late Devonian-Early Mississippian time and consisted mostly of clay in the Devonian and quartz and lesser chert sand in the Mississippian. Fossil assemblages, chiefly conodonts, indicate that any hiatus in this sequence is less than a stage in duration. Detailed lithologic and biostratigraphic comparisons of the Maiyumerak sequence with coeval

rocks throughout the western Brooks Range confirm that the Maiyumerak rocks are most similar to Devonian-Mississippian rocks in sequences of allochthon 3, as proposed by Mayfield and others (1988), and suggest that these rocks accumulated on a single carbonate platform, the Kelly platform.

Middle Devonian carbonate rocks in the Maiyumerak Mountains correlate well with coeval rocks in the central Baird Mountains quadrangle, but Upper Devonian rocks in these areas display contrasting facies and suggest that these rocks were not contiguous during deposition. No rocks correlative with the Devonian part of the Maiyumerak sequence are known to occur within the Schwatka structural province in the northeastern Baird Mountains quadrangle; at least some parts of the Schwatka province may have been subaerially exposed throughout the Middle and (or) Late Devonian. No evidence of unconformity has been found within the Maiyumerak sequence, but intertidal and supratidal environments prevailed towards the end of the Middle Devonian. These are the shallowest depositional environments that have been documented in the Maiyumerak sequence, and correspond to a period of uplift and (or) regression throughout northern Alaska.

Terrigenous material in the Maiyumerak carbonate sequence matches the Endicott Group in age and composition and was probably derived from the same source. This conclusion necessitates certain modifications of the paleogeographic model proposed by Mayfield and others (1988) for the western Brooks Range. Their model implies a Devonian-Mississippian paleogeography similar to that of present-day northeastern Australia, with the Endicott delta separated from the Kelly carbonate platform by a trough 50-100 km wide. A common provenance for the Endicott Group and terrigenous detritus reaching the Kelly carbonate platform suggests that any trough between the two was only an embayment, and (or) did not form until Late Mississippian time. The Kelly platform was drowned in late Meramecian time, and neritic carbonate deposition was never reestablished in this part of northwestern Alaska; demise of the carbonate platform most likely reflects long-term tectonic factors, such as northward drift of northern Alaska, in conjunction with a short-term eustatic rise.

REFERENCES CITED

- Adams, J. E., and Rhodes, M. L., 1960, Dolomitization by seepage refluxion: American Association of Petroleum Geologists Bulletin, v. 44, p. 1912-1920.
- Anderson, A. V., 1987, Provenance and petrofacies of the Endicott and Hammond terranes, Phillip Smith Mountains and Arctic quadrangles, Brooks Range, Alaska (abs.): Geological Society of America Abstracts with Programs, p. 354.
- Armstrong, A. K., 1970a, Carbonate facies and the lithostrotionid corals of the Mississippian Kogruek Formation, DeLong Mountains, Northwestern Alaska: U.S. Geological Survey Professional Paper 664, 38 p.
- Armstrong, A. K., 1970b, Mississippian dolomites from Lisburne Group, Killik River, Mount Bupto Region, Alaska: American Association of Petroleum Geologists Bulletin, v. 54, p. 251-264.
- Armstrong, A. K., 1972, Biostratigraphy of Mississippian lithostrotionoid corals, Lisburne Group, Arctic Alaska: U.S. Geological Survey Professional Paper 743-A, p. A1-A28.
- Armstrong, A. K., 1974, Carboniferous carbonate depositional models, preliminary lithofacies and paleotectonic maps, Arctic Alaska: American Association of Petroleum Geologists Bulletin, v. 58, p. 621-645.
- Armstrong, A. K., and Bird, K. J., 1976, Carboniferous environments of deposition and facies, Arctic Alaska, in Miller, T. P., ed., Symposium on Recent and Ancient Sedimentary Environments in Alaska: Alaska Geological Society Symposium Proceedings, p. A1-A16.
- Armstrong, A. K., and Mamet, B. L., 1970, Biostratigraphy and dolomite porosity trends of the Lisburne Group, in Adkison, W. L., and Brosgé, M. M., eds., Proceedings of the geological seminar on the North Slope of Alaska: American Association of Petroleum Geologists Pacific Section Meeting, Los Angeles, p. N1-N16.
- Armstrong, A. K., and Mamet, B. L., 1977, Carboniferous microfacies, microfossils, and corals, Lisburne Group, Arctic Alaska: U.S. Geological Survey Professional Paper 849, 144 p.
- Armstrong, A. K., and Mamet, B. L., 1978, Microfacies of the Carboniferous Lisburne Group, Endicott Mountains, Arctic Alaska, in Stelk, C. R., and Chatterton, B. D. E., eds., Western and Arctic Biostratigraphy: Geological Association of Canada Special Paper 18, p. 333-394.
- Armstrong, A. K., Mamet, B. L., and Dutro, J. T., Jr., 1971, Lisburne Group, Cape Lewis-Niak Creek, Northwestern Alaska: U.S. Geological Survey Professional Paper 750-B, p. B23-B34.
- Armstrong, R. L., Harakal, J. E., Forbes, R. B., Evans, B. W., and Thurston, S. P., 1986, Rb-Sr and K-Ar study of metamorphic rocks of the Seward Peninsula and southwestern Brooks Range, Alaska, in Evans, B. W., and Brown, E. H., eds., Blueschists and Eclogites: Geological Society of America Memoir 164, p. 185-203.
- Bathurst, R. G. C., 1976, Carbonate sediments and their diagenesis: Developments in

Sedimentology, v. 12, Elsevier, New York, 658 p.

- Bathurst, R. C. G., 1987, Diagenetically enhanced bedding in argillaceous platform limestones: stratified cementation and selective compaction: *Sedimentology*, v. 34, p. 749-778.
- Bird, K. J., and Jordan, C. F., 1977, Lisburne Group (Mississippian and Pennsylvanian), potential major hydrocarbon objective of Arctic Slope, Alaska: *American Association of Petroleum Geologists Bulletin*, v. 61, p. 1493-1512.
- Bond, G. C., Kominz, M. A., Steckler, M. S., and Grotzinger, J. P., 1989, Role of subsidence, flexure, and eustasy in the evolution of early Paleozoic passive-margin carbonate platforms, in Crevello, P. D., Wilson, J. L., Sarg, J. F., and Read, J. F., eds., *Controls on carbonate platform and basin development*, SEPM Special Publication No. 44, p. 39-61.
- Bowsher, A. L., and Dutro, J. T., Jr., 1957, The Paleozoic section in the Shainin Lake area, central Brooks Range, Alaska: U.S. Geological Survey Professional Paper 303-A, p. 1-39.
- Brosgé, W. P., Nilsen, T. H., Moore, T. E., and Dutro, J. T., Jr., 1988, Geology of the Upper Devonian and Lower Mississippian(?) Kanayut Conglomerate in the central and eastern Brooks Range, in Gryc, G., ed., *Geology and Exploration of the National Petroleum Reserve in Alaska, 1974 to 1982*: U.S. Geological Survey Professional Paper 1399, p. 299-316.
- Brosgé, W. P., Reiser, H. M., and Tailleir, I. L., 1967, Copper analyses of selected samples, southwestern Brooks Range, Alaska: U.S. Geological Survey Open-File Report 274, 1 sheet, scale 1:1,000,000.
- Brosgé, W. P., and Tailleir, I. L., 1970, Depositional history of northern Alaska, in Adkison, W. L., and Brosgé, M. M., eds., *Proceedings of the geological seminar on the North Slope of Alaska*: American Association of Petroleum Geologists Pacific Section Meeting, Los Angeles, p. D1-D18.
- Campbell, R. H., 1967, Areal geology in the vicinity of the Chariot site, Lisburne Peninsula, northwest Alaska: U.S. Geological Survey Professional Paper 395, 71 p.
- Churkin, M., Jr., Nokleberg, W. J., and Huie, C., 1979, Collision-deformed Paleozoic continental margin, western Brooks Range, Alaska: *Geology*, v. 7, p. 379-383.
- Crane, R. C., 1987, Arctic reconstruction from an Alaskan viewpoint, in Tailleir, I. L., and Weimer, P., eds., *Alaskan North Slope Geology: Pacific Section*, Society of Economic Paleontologists and Mineralogists, Bakersfield, California, Book 50, p. 769-783.
- Curtis, S. M., Eilersieck, I., Mayfield, C. F., and Tailleir, I. L., 1983, Reconnaissance geologic map of the De Long Mountains A1, B1, and part of the C2 quadrangles, Alaska: U.S. Geological Survey Open-File Report 83-185, 53 p., 1 sheet, scale 1:63,360.
- Curtis, S. M., Eilersieck, I., Mayfield, C. F., and Tailleir, I. L., 1984, Reconnaissance geologic map of southwestern Misheguk Mountain quadrangle, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-1502, 2 sheets, scale 1:63,360.

- Davies, P. J., Symonds, P. A., Feary, D. A., and Pigram, C. J., 1988, Facies models in exploration--the carbonate platforms of north-east Australia: *APEA Journal*, p. 123-143.
- Davies, P. J., Symonds, P. A., Feary, D. A., and Pigram, C. J., 1989, The evolution of the carbonate platforms of northeast Australia, in Crevello, P. D., Wilson, J. L., Sarg, J. F., and Read, J. F., eds., *Controls on carbonate platform and basin development*, SEPM Special Publication No. 44, p. 233-258.
- Dickson, J. A. D., 1966, Carbonate identification and genesis as revealed by staining: *Journal of Sedimentary Petrology*, v. 36, p. 491-505.
- Dillon, J. T., Harris, A. G., and Dutro, J. T., Jr., 1987, Preliminary description and correlation of lower Paleozoic fossil-bearing strata in the Snowden Mountain area of the south-central Brooks Range, Alaska, in Tailleir, I. L., and Weimer, P., eds., *Alaskan North Slope Geology: Pacific Section*, Society of Economic Paleontologists and Mineralogists, Bakersfield, California, Book 50, p. 337-345.
- Doyle, L. J., and Roberts, H. H., 1988, Carbonate-Clastic Transitions: Developments in Sedimentology, v. 42, Elsevier, New York, 304 p.
- Dumoulin, J. A., and Harris, A. H., 1987, Lower Paleozoic carbonate rocks of the Baird Mountains quadrangle, western Brooks Range, Alaska, in Tailleir, I. L., and Weimer, P., eds., *Alaskan North Slope Geology: Pacific Section*, Society of Economic Paleontologists and Mineralogists, Bakersfield, California, Book 50, p. 311-336.
- Dunham, R. J., 1962, Classification of carbonate rocks according to depositional texture, in Ham, W. E., ed., *Classification of Carbonate Rocks*, American Association of Petroleum Geologists Memoir 1, p. 108-121.
- Dutro, J. T., Jr., 1987, Revised megafossil biostratigraphic zonation for the Carboniferous of northern Alaska, in Tailleir, I. L., and Weimer, P., eds., *Alaskan North Slope Geology: Pacific Section*, Society of Economic Paleontologists and Mineralogists, Bakersfield, California, Book 50, p. 359-364.
- Ellersieck, I., 1985, Tectonic shortening of the upper crust in the western Brooks Range, Alaska: structure of the northwestern Baird Mountains: M.S. thesis, University of California, Berkeley, 121 p.
- Ellersieck, I., Curtis, S. M., Mayfield, C. F., and Tailleir, I. L., 1983, Reconnaissance geologic map of the De Long Mountains A2, B2, and part of the C2 quadrangles, Alaska: U.S. Geological Survey Open-File Report 83-184, 53 p., 1 sheet, scale 1:63,360.
- Ellersieck, I., Curtis, S. M., Mayfield, C. F., and Tailleir, I. L., 1984, Reconnaissance geologic map of south-central Misheguk Mountain quadrangle, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-1504, 2 sheets, scale 1:63,360.
- Epstein, A. G., Epstein, J. B., and Harris, L. D., 1977, Conodont color alteration--an index to organic metamorphism: U.S. Geological Survey Professional Paper 995, 27 p.
- Fischbuch, N. R., 1968, Stratigraphy, Devonian Swan Hills reef complexes of central Alberta:

- Canadian Petroleum Geology Bulletin, v. 16, p. 444-587.
- Fischer, A. G., 1975, Tidal deposits, Dachstein Limestone of the North-Alpine Triassic, *in* Ginsburg, R. N., ed., *Tidal Deposits*: Springer-Verlag, New York, p. 235-242.
- Flood, P. G., and Orme, G. R., 1988, Mixed siliciclastic/carbonate sediments of the northern Great Barrier Reef, *in* Doyle, L. J., and Roberts, H. H., eds., *Carbonate-Clastic Transitions: Developments in Sedimentology*, v. 42, Elsevier, New York, p. 175-206.
- Gordon, M., Jr., 1957, Mississippian cephalopods of northern and eastern Alaska: U.S. Geological Survey Professional Paper 283, 61 p.
- Hallam, A., and Bradshaw, M. J., 1979, Bituminous shales and oolitic ironstones as indicators of transgressions and regressions: *Journal of the Geological Society of London*, v. 136, p. 157-164.
- Halley, R. B., Harris, P. M., and Hine, A. C., 1983, Bank margin, *in* Scholle, P. A., Bebout, D. G., and Moore, C. H., eds., *Carbonate Depositional Environments*: American Association of Petroleum Geologists Memoir 33, p. 463-506.
- Haq, B. U., and van Eysinga, F. W. B., 1987, *Geological Time Table*: Elsevier, New York, 4th ed., 1 sheet.
- Holdsworth, B. K., and Murchey, B. L., 1988, Paleozoic radiolarian biostratigraphy of the northern Brooks Range, Alaska, *in* Gryc, G., ed., *Geology and Exploration of the National Petroleum Reserve in Alaska, 1974 to 1982*: U.S. Geological Survey Professional Paper 1399, p. 777-797.
- Horowitz, A. S., and Potter, P. E., 1971, *Introductory Petrography of Fossils*: Springer-Verlag, New York, 302 p.
- James, N. P., 1984, Shallowing-upward sequences in carbonates, *in* Walker, R. G., ed., *Facies Models*: Geological Association of Canada, Toronto, p. 213-228.
- Johnson, J. G., Klapper, G., and Sandberg, C. A., 1985, Devonian eustatic fluctuations in Euramerica: *Geological Society of America Bulletin*, v. 96, p. 567-587.
- Karl, S. M., Dumoulin, J. A., Ellersieck, I., Harris, A. G., and Schmidt, J. M., 1989, Preliminary geologic map of the Baird Mountains and part of the Selawik quadrangles, Alaska: U.S. Geological Survey Open-File Report 89-551, 65 p., 1 pl., scale 1:250,000.
- Kelling, G., and Mullin, P. R., 1975, Graded limestone and limestone-quartzite couplets: possible storm-deposits from the Moroccan Carboniferous: *Sedimentary Geology*, v. 13, p. 161-190.
- Kendall, C. G. St. C., and Schlager, W., 1981, Carbonates and relative changes in sea level: *Marine Geology*, v. 44, p. 181-212.
- Lerand, M., 1973, Beaufort Sea, *in* McCrossam, R. G., ed., *The Future Petroleum Provinces of Canada--Their Geology and Potential*: Canadian Society of Petroleum Geology, Memoir 1,

p. 315-386.

- Lucia, F. J., 1962, Diagenesis of a crinoidal sediment: *Journal of Sedimentary Petrology*, v. 32, p. 848-865.
- Magoon, L. B., and Bird, K. J., 1988, Evaluation of petroleum source rocks in the National Petroleum Reserve in Alaska, using organic-carbon content, hydrocarbon content, visual kerogen, and vitrinite reflectance, in Gryc, G., ed., *Geology and Exploration of the National Petroleum Reserve in Alaska, 1974 to 1982*: U.S. Geological Survey Professional Paper 1399, p. 381-450.
- Mamet, B. L., and Skipp, B., 1970, Lower Carboniferous Foraminifera: preliminary zonation and stratigraphic implication for the Mississippian of North America: *Sixth International Congress Carboniferous Stratigraphy*, v. 3, p. 1129-1146.
- Mayfield, C. F., Curtis, S. M., Ellersieck, I., and TAILLEUR, I. L., 1983, Reconnaissance geologic map of the De Long Mountains A3, B3, and parts of the A4 and B4 quadrangles, Alaska: U.S. Geological Survey Open-File Report 83-183, 59 p., 1 sheet, scale 1:63,360.
- Mayfield, C. F., Curtis, S. M., Ellersieck, I., and TAILLEUR, I. L., 1984, Reconnaissance geologic map of southeastern Misheguk Mountain quadrangle, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-1503, 2 sheets, scale 1:63,360.
- Mayfield, C. F., Ellersieck, I., and TAILLEUR, I. L., 1987, Reconnaissance geologic map of the Noatak C5, D5, D6, and D7 quadrangles, Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-1814, 1 sheet, scale 1:63,360.
- Mayfield, C. F., TAILLEUR, I. L., and Ellersieck, I., 1988, Stratigraphy, structure, and palinspastic synthesis of the western Brooks Range, northwestern Alaska, in Gryc, G., ed., *Geology and Exploration of the National Petroleum Reserve in Alaska, 1974-1982*: U.S. Geological Survey Professional Paper 1399, p. 143-186.
- Mitchum, R. M., Jr., 1977, Glossary of terms used in seismic stratigraphy, in Payton, C. E., ed., *Seismic stratigraphy--applications to hydrocarbon exploration*, American Association of Petroleum Geologists Memoir 26, p. 205-212.
- Moore, T. E., and Nilsen, T. H., 1984, Regional sedimentological variations in the Upper Devonian and Lower Mississippian (?) Kanayut Conglomerate, Brooks Range, Alaska: *Sedimentary Geology*, v. 38, p. 464-398.
- Moore, T. E., Wallace, W. K., Bird, K. J., Karl, S. M., Mull, C. G., and Dillon, J. T., in press, *Geology of northern Alaska*, in Plafker, G., ed., *Geology of Alaska: Geological Society of America Decade of North American Geology*.
- Mount, J. F., 1984, Mixing of siliciclastic and carbonate sediments in shallow shelf environments: *Geology*, v. 12, p. 432-435.
- Mull, C. G., Roeder, D. H., TAILLEUR, I. L., Pessel, G. H., Grantz, A., and May, S. D., 1987, Geologic sections and maps across Brooks Range and Arctic slope to Beaufort Sea, Alaska: Geological Society of America Map and Chart Series MC-28S.

- Murchey, B. L., Jones, D. L., Holdsworth, B. K. and Wardlaw, B. R., 1988, Distribution patterns of facies, radiolarians, and conodonts in the Mississippian to Jurassic siliceous rocks of the northern Brooks Range, Alaska, *in* Gryc, G., ed., *Geology and Exploration of the National Petroleum Reserve in Alaska, 1974 to 1982*: U.S. Geological Survey Professional Paper 1399, p. 697-724.
- Nilsen, T. H., 1981, Upper Devonian and Lower Mississippian redbeds, Brooks Range, Alaska, *in* Miall, A. D., ed., *Sedimentation and tectonics in alluvial basins*: Geological Association of Canada Special Paper 23, p. 187-219.
- Oliver, W. A., Jr., Merriam, C. W., and Churkin, M., Jr., 1975, Ordovician, Silurian, and Devonian corals of Alaska: U.S. Geological Survey Professional Paper 823-B, p. 13-44.
- Pilkey, O. H., Bush, D. M., and Rodriguez, R. W., Carbonate-terrigenous sedimentation on the north Puerto Rico shelf, *in* Doyle, L. J., and Roberts, H. H., eds., *Carbonate-Clastic Transitions: Developments in Sedimentology*, v. 42, Elsevier, New York, p. 231-250.
- Read, J. F., 1985, Carbonate platform facies models: *American Association of Petroleum Geologists Bulletin*, v. 69, p. 1-21.
- Read, J. F., 1989, Controls on evolution of Cambrian-Ordovician passive margin, U. S. Appalachians, *in* Crevello, P. D., Wilson, J. L., Sarg, J. F., and Read, J. F., eds., *Controls on carbonate platform and basin development*, SEPM Special Publication No. 44, p. 147-165.
- Ross, C. A., and Ross, J. R. P., 1988, Late Paleozoic transgressive-regressive deposition, *in* Wilgus, C. K., Hastings, B. S., Posamentier, H., Van Wagoner, J., Ross, C. A., and Kendall, C. G. St. C., eds., *Sea-level changes: an integrated approach*: SEPM Special Publication No. 42, p. 227-247.
- Rowley, D. B., and Lottes, A. L., 1988, Plate-kinematic reconstructions of the North Atlantic and Arctic: Late Jurassic to Present: *Tectonophysics*, v. 155, p. 73-120.
- Rupp, A. W., 1966, Origin, structure, and environmental significance of recent and fossil calcispheres (abs.): *Geological Society of America Abstracts with Programs*, p. 186.
- Sable, E. G., and Dutro, J. T., Jr., 1961, New Devonian and Mississippian formations in the De Long Mountains, northern Alaska: *American Association of Petroleum Geologists Bulletin*, v. 45, p. 585-593.
- Sandberg, C. A., 1976, Conodont biofacies of Late Devonian *Polygnathus striatus* Zone in western United States, *in* Barnes, C. R., ed., *Conodont Paleogeology*: Geological Association of Canada Special Paper 15, p. 172-186.
- Sandberg, C. A., and Gutschick, R. C., 1984, Distribution, microfauna, and source-rock potential of Mississippian Delle Phosphatic Member of Woodman Formation and equivalents, Utah and adjacent states, *in* Woodward, J. and others, eds., *Hydrocarbon Source Rocks of the Greater Rocky Mountain Region*: Rocky Mountain Association of Geologists, Denver, Co., p. 135-178.
- Sandberg, C. A., and Ziegler, W., 1979, Taxonomy and biofacies of important conodonts of Late

- Devonian *striacus*-Zone, United States and Germany: *Geologica et Palaeontologica*, v. 13, p. 173-212.
- Sarg, J. F., 1988, Carbonate sequence stratigraphy, in Wilgus, C. K., Hastings, B. S., Posamentier, H., Van Wagoner, J., Ross, C. A., and Kendall, C. G. St. C., eds., *Sea-level changes: an integrated approach*: SEPM Special Publication No. 42, p. 155-181.
- Schmidt, J. M., 1987, Paleozoic extension of the western Brooks Range (WBR) continental margin—evidence from mineral deposits, igneous rocks and sedimentary facies (abs.): *Geological Society of America Abstracts with Programs*, p. 447.
- Shaw, A. B., 1964, *Time in Stratigraphy*: McGraw-Hill Publishing, New York, 353 p.
- Shinn, E. A., 1983a, Birdseyes, fenestrae, shrinkage pores, and loferites: a reevaluation: *Journal of Sedimentary Geology*, v. 53, p. 619-628.
- Shinn, E. A., 1983b, Tidal flat, in Scholle, P. A., Bebout, D. G., and Moore, C. H., eds., *Carbonate Depositional Environments*: American Association of Petroleum Geologists Memoir 33, p. 171-210.
- Smith, D. G., 1987, Late Paleozoic to Cenozoic reconstructions of the Arctic, in Tallleur, I. L., and Weimer, P., eds., *Alaskan North Slope Geology: Pacific Section*, Society of Economic Paleontologists and Mineralogists, Bakersfield, California, Book 50, p. 785-795.
- Tallleur, I. L., Brosgé, W. P., and Reiser, H. N., 1967, Palimpsestic analysis of Devonian rocks in northwestern Alaska, in Oswald, D. H., ed., *International Symposium on the Devonian System*, v. 2: Alberta Society of Petroleum Geologists, Calgary, Canada, p. 1345-1361.
- Till, A. B., Schmidt, J. M., and Nelson, S. W., 1988, Thrust involvement of metamorphic rocks, southwestern Brooks Range, Alaska: *Geology*, v. 16, p. 930-933.
- Vail, P. R., Mitchum, R. M., Jr., and Thompson, S., III, 1977, Global cycles of relative changes of sea level, in Payton, C.E., ed., *Seismic Stratigraphy—Applications to Hydrocarbon Exploration*, American Association of Petroleum Geologists Memoir 26, p. 83-98.
- Wilson, J. L., 1975, *Carbonate Facies In Geologic History*: Springer-Verlag, New York, 470 p.
- Wilson, J. L., and Jordan, C., 1983, Middle shelf, in Scholle, P. A., Bebout, D. G., and Moore, C. H., eds., *Carbonate Depositional Environments*: American Association of Petroleum Geologists Memoir 33, p. 297-344.
- Zuffa, G. G., 1980, Hybrid arenites: their composition and classification: *Journal of Sedimentary Petrology*, v. 50, p. 21-29.