

**STOLEN FROM
RICHARD D. REGER**

THE ALASKA EARTHQUAKE, MARCH 27, 1964:
EFFECTS ON THE HYDROLOGIC REGIMEN

Seismic Seiches
From the March 1964
Alaska Earthquake

By ARTHUR McGARR and ROBERT C. VORHIS

*An interpretation of the continental distribution
of seiches from the earthquake*

GEOLOGICAL SURVEY PROFESSIONAL PAPER 544-E

UNITED STATES DEPARTMENT OF THE INTERIOR

STEWART L. UDALL, *Secretary*

GEOLOGICAL SURVEY

William T. Pecora, *Director*



UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON : 1968

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C. 20402

THE
ALASKA EARTHQUAKE
SERIES

The U.S. Geological Survey is publishing the results of investigations of the Alaska earthquake of March 27, 1964, in a series of six Professional Papers. Professional Paper 544 describes the effects of the earthquake on the hydrologic regimen. Other chapters in this volume describe the effects of the earthquake on the hydrology of south-central Alaska, the Anchorage area, areas outside Alaska, and the effects on glaciers.

Other Professional Papers in the series describe the history of the field investigations and reconstruction; the effects of the earthquake on communities; the regional effects of the earthquake; and the effects on transportation, utilities, and communications

CONTENTS

| | Page | | Page | | Page |
|--|------|--|------|---|------|
| Abstract..... | E1 | Location and nature of the seiches—Continued | | Interpretation of seiche distribution—Continued | |
| Introduction..... | 1 | Geographic distribution..... | E11 | Relation to seismic surface waves..... | E15 |
| Purposes of the study..... | 2 | Hydrodynamic factors..... | 12 | Radiation pattern..... | 15 |
| Definition of terms..... | 2 | Interpretation of seiche distribution..... | 13 | Distance from epicenter.... | 15 |
| Previous studies of seismic seiches..... | 3 | Relation to geologic features.. | 13 | Lateral refraction..... | 16 |
| Sources of data..... | 3 | Sediment thickness..... | 13 | Local crustal structure..... | 18 |
| Acknowledgments..... | 4 | Thrust faults..... | 13 | Irregular structures..... | 19 |
| General theoretical background.. | 5 | Basins, arches, and domes... | 14 | Seiches and seismic intensity.. | 20 |
| Location and nature of the seiches..... | 7 | Edge of overlaps..... | 14 | Conclusions and recommendations..... | 22 |
| Seiche data..... | 7 | Rocky Mountain system.... | 14 | References..... | 23 |
| Gaging stations, instruments, and their records..... | 10 | Miscellaneous areas..... | 15 | | |

ILLUSTRATIONS

PLATE

1. Map of the conterminous United States and southern Canada showing occurrence of seiches from the Alaska earthquake and their relation to major tectonic features..... In pocket

FIGURES

| | Page |
|---|------|
| 1. The coordinate system applied to a theoretical water body and seiches of the first, third, and fifth modes..... | E6 |
| 2-5. Records: | |
| 2. The largest seiche recorded on a stream in each of eight States..... | 9 |
| 3. Some large seismic seiches on reservoirs..... | 9 |
| 4. Three types of bubble-gage records of Alaska earthquake seiches..... | 10 |
| 5. Seiche effects of Alaska earthquake on stage and flow, Miami area, Florida.. | 10 |
| 6-8. Maps: | |
| 6. Map of conterminous United States showing seiche density by State and by river basin..... | 11 |
| 7. Maximum horizontal acceleration at stations of the World-wide Standard Seismograph Network in the United States calculated for two aftershocks of the Alaska earthquake..... | 16 |
| 8. Phase-velocity distribution of 20-second Rayleigh waves in North America.. | 17 |
| 9. Amplification of Rayleigh-wave displacements in low-rigidity sediment overlying high-rigidity rock, (A) for 15- and (B) for 8-second period waves..... | 18 |
| 10. Variation in amplitude of surface horizontal acceleration as a function of "layer" shear wave velocity, for 6- and 10-second period Rayleigh waves..... | 20 |
| 11. Alaska earthquake seiches plotted on the intensity map of the Missouri earthquake of October 21, 1965..... | 21 |

TABLES

| | Page |
|--|------|
| 1. Summary of 859 seismic effects from the Alaska earthquake on surface-water bodies throughout the world..... | E7 |
| 2. First-, third-, and fifth-order modes, in seconds, for seiches on water bodies with selected widths and depths..... | 12 |
| 3. Seismic effects of the Alaska earthquake at surface-water gages..... | 25 |

THE ALASKA EARTHQUAKE, MARCH 27, 1964: EFFECTS ON THE HYDROLOGIC REGIMEN

SEISMIC SEICHES FROM THE MARCH 1964 ALASKA EARTHQUAKE¹

By Arthur McGarr, Lamont Geological Observatory of Columbia University, Palisades, N.Y., and Robert C. Vorhis, U.S. Geological Survey

ABSTRACT

Seismic seiches caused by the Alaska earthquake of March 27, 1964, were recorded at more than 850 surface-water gaging stations in North America and at 4 in Australia. In the United States, including Alaska and Hawaii, 763 of 6,435 gages registered seiches. Nearly all the seismic seiches were recorded at teleseismic distance. This is the first time such far-distant effects have been reported from surface-water bodies in North America. The densest occurrence of seiches was in States bordering the Gulf of Mexico.

The seiches were recorded on bodies of water having a wide range in depth, width, and rate of flow. In a region containing many bodies of water, seiche distribution is more dependent on geologic and seismic factors than on hydrodynamic ones. The concept that seiches are caused by the horizontal acceleration of water by seismic surface waves has been extended in this paper to show

that the distribution of seiches is related to the amplitude distribution of short-period seismic surface waves. These waves have their greatest horizontal acceleration when their periods range from 5 to 15 seconds. Similarly, the water bodies on which seiches were recorded have low-order modes whose periods of oscillation also range from 5 to 15 seconds.

Several factors seem to control the distribution of seiches. The most important is variations of thickness of low-rigidity sediments. This factor caused the abundance of seiches in the Gulf Coast area and along the edge of sedimentary overlaps. Major tectonic features such as thrust faults, basins, arches, and domes seem to control seismic waves and thus affect the distribution of seiches. Lateral refraction of seismic surface waves due to variations in local phase-velocity values was responsible for increase in seiche density in certain areas.

For example, the Rocky Mountains provided a wave guide along which seiches were more numerous than in areas to either side. In North America, neither direction nor distance from the epicenter had any apparent effect on the distribution of seiches.

Where seismic surface waves propagated into an area with thicker sediment, the horizontal acceleration increased about in proportion to the increasing thickness of the sediment. In the Mississippi Embayment however, where the waves emerged from high rigidity crust into the sediment, the horizontal acceleration increased near the edge of the embayment but decreased in the central part and formed a shadow zone.

Because both seiches and seismic intensity depend on the horizontal acceleration from surface waves, the distribution of seiches may be used to map the seismic intensity that can be expected from future local earthquakes.

INTRODUCTION

Seismic waves from the Alaska earthquake of March 28, 1964,² were so powerful that they caused

water bodies to oscillate at many places throughout North America. Those oscillations, or seismic seiches, were recorded at hundreds of surface-water gaging stations although they had rarely been reported following previous earthquakes and, when reported, had received little study. Local reports of numerous seiches resulting from the Alaska earthquake prompted one of the authors, Vorhis, to request records of Alaska earth-

quake seiches from his colleagues in the U.S. Geological Survey and from other hydrologic organizations both in North America and throughout the world. The replies identified most locations where seiches were recorded. In the United States, of all gages which could have recorded a seiche at the time of the Alaska earthquake, slightly more than 10 percent did. Factors other than the nature of the recording installation and the

¹ Lamont Geological Observatory Contribution 1070.

² The date and time of an earthquake can be given either as local or Greenwich time. In and near the epicentral region, it is customary to give the local time, such as 5:36 p.m. A.s.t. on March 27, 1964, for the Alaska earthquake. In studies of a worldwide nature, the date and time of an earthquake are usually given in Greenwich time. Thus, the Alaska earthquake occurred at 03:36 on March 28, 1964, G.c.t.

geometry of the water body seem to have controlled the pattern of seiche occurrence.

PURPOSES OF THE STUDY

The purposes of the study were (1) to assemble and present the data on all known seismic seiches resulting from the Alaska earthquake, (2) to analyze their distribution in relation to possible controls, (3) to apply existing theory to analysis of seiches recorded in bodies of known dimensions, and (4) to determine what hydrologic and seismologic implications can be drawn from seiche data.

In attempting to interpret seiche distribution, there are at least two approaches. One is to assume that the seismic waves causing the seiches were uniform throughout North America. Regional variations in seiche distribution would then result from variations in the capacity of water bodies to couple into the seismic waves. After preliminary studies, the authors decided that an alternative approach was needed.

There were 6,435 analog-type surface-water gages operating in the United States at the time of the earthquake. This number is assumed to be large enough to average out the varying response characteristics of individual stations within discrete regions of the country. The preferential concentration of seiches in certain regions implies varying amplitude distribution of seismic waves and serves to demonstrate again that geologic features materially influence seismic waves.

It should be noted that surface-water recorders are just one of at least three types of instruments maintained for nonseismic studies that can detect the passage of seismic waves. The other two are microbarographs and recorders on ground-water observation wells.

In a sense, the three types of instruments provide complementary seismic data: the surface-water gages record the effect of horizontal acceleration of seismic waves, microbarographs record the air-pressure fluctuations caused by vertical velocity of the ground, and the instruments on wells record the influences of transient and permanent strain induced by seismic waves on aquifers. Barometric disturbances due to the Alaska shock have been discussed by Donn and Posmentier (1964) and ground-water fluctuations have been treated by Vorhis (1967).

This auxiliary instrumentation was more important than usual at the time of the Alaska earthquake because nearly all operating seismographs in North America were temporarily put out of action by the extremely large amplitudes of the seismic waves.

DEFINITION OF TERMS

Because this paper is concerned with both hydrology and seismology, some of the terms which may be unfamiliar to the hydrologist or the nonseismologist are defined as they are used in this paper.

Amplitude. One half the wave height.

Double amplitude. The height of a wave from crest to trough.

Lateral refraction. A horizontal deflection of a seismic surface wave due to change in its phase velocity in passing from one rock medium to another.

Love wave. A seismic surface wave whose motion is horizontally polarized in a direction transverse to the direction of wave propagation.

Mode. One of the stationary patterns of vibration of which an oscillatory system is capable. In this paper, "mode" may refer both to seismic surface waves and to water waves. The

application to water waves is shown in figure 1. First-order mode is also commonly referred to as the fundamental mode.

Phase velocity. The velocity of a particular spectral component of a wave form.

Radiation pattern. The relative directional intensity of seismic surface waves.

Rayleigh wave. A seismic surface wave whose ground motion is elliptical in the plane defined by the vertical and the direction of propagation.

Seiche. A term first used by Forel (1895) to apply to standing waves set up on the surface of Lake Geneva by wind and by changes in barometric pressure. The term has been extended to all standing waves on any body of water whose period is determined by the resonant characteristics of the containing basin as controlled by its physical dimensions.

Seismic intensity. A measure of earthquake severity based on the damage produced by seismic waves in a given region.

Seismic seiche. A term first used by Kvale (1955) in discussing oscillation of lake levels in Norway and England caused by the Assam earthquake of August 15, 1950. His usage has been extended in this paper to apply to standing waves set up on rivers, reservoirs, ponds, and lakes at the time of passage of seismic waves from an earthquake.

Seismicity. The relative frequency of earthquake occurrence in a given region.

Shadow zone. An area or region where seiche activity is small or absent because of some sort of barrier to the transmission of seismic surface waves.

Standing wave. A single-frequency mode of vibration in which the nodes and antinodes have fixed

positions. In this paper, standing waves have the form shown in equation (1) on page E5.

Surface wave. A wave of Love or Rayleigh type that travels around rather than through the earth.

Teleseismic distance. A distance of 1,000 kilometers (600 miles) or more from the earthquake epicenter.

Wave guide. A part of the earth's crust and upper mantle that tends to channel seismic energy.

PREVIOUS STUDIES OF SEISMIC SEICHES

The first published mention of seismic seiches known to the authors is with respect to the great earthquake of November 1, 1755, at Lisbon, Portugal. In a review of hydrologic effects of that earthquake, Wilson (1953) referred to an article in *Scot's Magazine* in 1755 that described remarkable seismic seiches in Loch Lomond, Loch Long, Loch Katrine and Loch Ness. Richter (1958, p. 110) mentioned other descriptions of seismic seiches caused by the Lisbon earthquake. These were observed in English harbors and ponds and were described originally in the *Proceedings of the Royal Society* in 1755.

Earthquake effects recorded by surface-water gages were first noted by Piper (1933, p. 475, fig. 2). He reported that two of six gages on the Mokelumne River in California showed a slight fluctuation caused by the December 20, 1932, earthquake at Lodi, Calif. Two other gages on a nearby diversion canal showed double amplitudes of 0.08 and 0.04 feet (24 and 12 mm) from the same earthquake. These phenomena were definitely seismic seiches although they were not so designated by Piper.

The U.S. Coast and Geodetic Survey (1945, p. 26) listed effects recorded on 18 stream gages in New York State that were caused by the September 5, 1944, earthquake in the St. Lawrence Valley.

The earthquake of January 25, 1946, in Switzerland in the Canton of Valais was recorded on two gages maintained by the Swiss Federal Water Survey on Lake Geneva, or Lac Léman (Mercanton, 1946). According to Mercanton, not a single seismic seiche was recorded during the 17 years in which Forel studied the seiches of Lake Geneva. This absence is especially surprising because during those years 69 earthquakes with 123 shocks were felt in the area. Thus, seiche records, even though numerous for the Alaska earthquake, may be relatively rare for other earthquakes or generally restricted to small bodies of water.

Kvale (1955) discussed previous seismic seiches, mainly those from the Lisbon earthquake; he also described 29 seiches recorded in fiords and lakes in Norway and 4 seiches on reservoirs in England, all caused by the Assam earthquake of August 15, 1950. He did not mention any seiches recorded on river gages. Surprisingly, no surface-water body in Norway or England is known to have responded to the Alaska earthquake. Most of the seiches that Kvale described from Norway were recorded in the western part of the country where the surface geology consists of sedimentary units. This distribution suggests that these seiches, if compared with local geological features in Norway, would give interpretations similar to those obtained from study of the distribution of seiches from the Alaska earthquake.

Stermitz (1964, p. 144, table 10) listed 54 stream gages that

recorded seiches caused by the Hebgen Lake earthquake of August 17, 1959. They were in Montana, Wyoming, Idaho, and Alberta, Canada, the most distant one being 340 miles from the epicenter. Three of these gages later recorded seismic seiches caused by the Alaska earthquake.

SOURCES OF DATA

Some data on seismic seiches from the Alaska earthquake have been obtained from published sources. Miller and Reddell (1964, p. 661) mention a reservoir at Lubbock, Tex., that registered a seiche of about 0.5 foot. Wigen and White (1964, p. 6, figs. 1-4) listed seiches at 10 locations on the west coast and one on the north coast (Cambridge Bay) of Canada. The periods of the seismic seiches were smaller than the seiche-wave periods that are frequently recorded on tide records. P. W. Strilaeff (1964, written commun.) listed nine seiches that were recorded in the Winnipeg District of Canada. He pointed out that on Lakes Winnipeg and Manitoba, seiches were recorded only at the narrows of the lakes. Similarly, at Lake of the Woods, only the recorder at Clearwater Bay indicated a seiche.

Seiche data for Texas were compiled by W. B. Mills (written commun., 1964) and for Tennessee by Milburn Hassler (written commun., 1965). Donn (1964) mentioned reports of waves on the Gulf Coast as high as 6 feet (1.8 m) that were caused by the Alaska earthquake and suggested that these and a seiche recorded by a tide gage at Freeport, Tex., were generated in resonance with seismic waves.

Using the same record from Freeport, Tex., McGarr (1965) developed a theory to explain the interaction between seismic surface waves and a channel filled

with water. The analysis included a few factors influencing the size of the seismic surface waves and several possible damping mechanisms. This theory is discussed in the section on "General Theoretical Background" (p. E5).

In a paper on hydrologic effects of the Alaska earthquake outside Alaska, Vorhis (1967) summarized seiche records for the conterminous United States and Hawaii. Those records and others that were obtained subsequently are described and interpreted in the present paper. Most of the data were received from the Water Resources Division of the U.S. Geological Survey, others were furnished by the Tennessee Valley Authority, the Walla Walla District of the U.S. Corps of Engineers, and the Illinois State Water Survey.

Data on seiches in Canada were compiled by the Water Resources Branch of the Canadian Department of Natural Resources and were supplied by the Canadian National Committee for the International Hydrologic Decade. Some additional unpublished seiche data for Manitoba, Saskatchewan, and Ontario were compiled by P. W. Strilaeff (written commun., 1964).

Records of four seiches were received from Australia. One on the Victoria River in northern Australia was furnished by the Northern Territory Administration of the Commonwealth of Australia, one on the Tantangara Reservoir in New South Wales was furnished by the Snowy Mountains Hydro-Electric Authority, one on a reservoir at Canberra was furnished by Robert Underwood of the Australian National University, and one on the Melicke Munjie River in eastern Victoria was furnished by the State Electricity Commission of Victoria. These seiches were the most dis-

tant and were the only ones known from outside North America and Hawaii.

ACKNOWLEDGMENTS

A world-wide solicitation for seismic-seiche data from a major earthquake had never been undertaken prior to the Alaska earthquake. To ascertain the geographic distribution of seiches resulting from the earthquake, all organizations in the world that might be expected to operate a hydrologic network were requested to submit copies of all charts that seemed to show earthquake effects. Professor Gerard Tison of the International Association of Scientific Hydrology and Dr. R. Ambroggi, Food and Agriculture Organization of the United Nations, assisted in the solicitation of data.

The agencies that furnished seiche data have been mentioned above, and their help is acknowledged with gratitude. Many other agencies went to considerable expense and trouble to examine a large number of charts for seismic seiches. Even though they found none, the negative reports were useful. The efforts of the following countries and their hydrologic organizations are acknowledged with appreciation:

Austria: Hydrographical Central Office

Australia:

Victoria State Rivers and Water Supply Commission

South Australia Engineering and Water Supply Department

New South Wales—Sydney Metropolitan Water Sewerage and Drainage Board

Snowy Mountains Hydro-Electric Authority

Queensland Irrigation and Water Supply Commission

British Guiana: Ministry of Works and Hydraulics

Ceylon: Department of Meteorology

China: Geological Survey of Taiwan

Ethiopia: Ministry of Public Works and Communications, Water Resources Department

Ghana: National Construction Corporation

Hungary: Research Institute for Water Resources

Indonesia: Hydrological Survey

Nepal: Ministry of Irrigation, Hydrological Survey Department

New Zealand: Ministry of Works

Norway: Water Resources and Electricity Board

Papua and New Guinea Administration

Portugal: Geological Survey
Republic of the Philippines: Department of Public Works and Communications

Bureau of Public Works
Southern Rhodesia: Geological Survey Office

Switzerland: Federal Office of Water Resources

Tasmania:

Rivers and Water Supply Commission

Hydroelectric Commission

Turkey: State Hydraulics Works

Uganda: Water Development Department

Zambia: Ministry of Lands and Natural Resources, Department of Water Affairs

Mr. F. A. Ekker of the Dow Chemical Co. furnished the orig-

inal records of seiches in tanks at Plaquemines, La., to Dr. D. H. Kupfer of Louisiana State University, who in turn made the charts available to the authors. Mr. Claud R. Erickson, engineer with the Lansing Water Department, furnished data on seiches in reservoirs at Lansing, Mich.

Dr. Jack Oliver of Columbia

University made many helpful suggestions and reviewed the manuscript. Other reviewers include J. P. Eaton, J. H. Feth, R. M. Waller, and C. L. O'Donnell, all of the U.S. Geological Survey; Dr. William Stauder, S. J., of Saint Louis University; and Dr. L. E. Alsop and Dr. J. E. Nafe of Columbia University.

The work by McGarr, reported in this paper, was partly sponsored by the U.S. Air Force Cambridge Research Laboratories, Office of Aerospace Research, under contract AF 19(628)-4082 (Columbia University) as part of the Vela-Uniform Program of the Advanced Research Projects Agency.

GENERAL THEORETICAL BACKGROUND

The seiches caused by the Alaska earthquake can be considered for purposes of analysis to have occurred in two distinct regions. One region, comprising most parts of Alaska, is an area of great seismic intensity where seiches can be caused by mechanisms such as landslides, submarine slides, tilting, tsunamis, and seismic surface waves. This variety of mechanisms makes the determination of the cause of a given seiche difficult. Seiches in this epicentral region of the Alaska earthquake are therefore not discussed.

The other region is in effect the rest of the world outside Alaska. In this region, most of which is at teleseismic distances from the epicenter, inelastic effects are unimportant and seismic seiches are generated solely by seismic surface waves. Although tsunamis also may occur in coastal areas, they travel so much more slowly than surface waves and have such long periods that the two cannot be confused.

The data considered in this paper are chiefly from charts of water-level recorders operating on continental bodies of water, primarily rivers, reservoirs, small lakes, and ponds. The primary problem, then, is to determine how seismic surface waves interact

with bodies of water of various sizes and shapes. A theory of interaction has been developed only for the long channel with rectangular cross section (McGarr, 1965). Although this model is idealized, it contains most of the interesting features of realistic and complicated situations. Further, the natural periods of response for water

bodies can be approximated fairly well by using the long-channel results.

According to McGarr (1965) the free surface level of an infinitely long channel will behave under the influence of a uniform time-dependent horizontal force per unit mass, $F(t)$, according to

$$\eta(x, t) = + \frac{4H}{\pi c} \sum_{n=0}^{\infty} \frac{\cos [(2n+1)\pi x L^{-1}]}{2n+1} \cdot \int_0^t F(\tau) e^{-k(t-\tau)/2} \cdot \sin \left[\frac{(2n+1)\pi c(t-\tau)}{L} \right] d\tau \quad (1)$$

where

$\eta(x, t)$ = height of the free surface above the undisturbed level, H = depth, L = width, $c = \sqrt{gH}$, the velocity of long water waves, g = gravity field strength, k = a damping constant, τ = an integration variable, t = time in seconds, n = an integer variable of summation.


Figure 1 (next page) shows the cross section of a theoretical channel and the coordinate system applied to it. The force per unit

mass due to the horizontal acceleration is in the x direction. A water level recorder at the edge of the channel will record

$$\eta(0, t) = + \frac{4H}{\pi c} \sum_{n=0}^{\infty} \frac{1}{2n+1} \int_0^t F(\tau) e^{-k(t-\tau)/2} \cdot \sin \left[\frac{(2n+1)\pi c(t-\tau)}{L} \right] d\tau \quad (2)$$

where

$\eta(0, t)$ = the height of the free surface above the undisturbed level at the edge of the channel.



1.—The coordinate system applied to a theoretical water body and seiches of the first, third, and fifth modes. Because of the nature of the seismic forcing function, only the odd-order modes are excited.

This expression shows that the height of a seiche is directly proportional to the horizontal acceleration provided by the seismic surface waves and \sqrt{H} , because $c = \sqrt{gH}$. Thus for a given surface-wave acceleration, a deeper channel will produce a higher seiche.

The damping constant k is included in equation (2) under the assumption that the attenuation of the seiche will be proportional to the velocity of water-particle motion. This assumption is not exactly true for all the factors contributing to the damping. However, the most important factors in dissipation, such as a sloping

beach, will yield damping curves that look similar to $e^{-kt/2}$; the assumption of a linear damping term is therefore probably acceptable.

The most important term in computing $\eta(0, t)$ is $F(t)$, the driving force. The fact that both Love and Rayleigh waves have a horizontal component of motion means that, no matter what the orientation of the channel, there will always be a component of horizontal acceleration parallel to the width. The primary problem is to determine the Love- and Rayleigh-wave amplitudes as a function of period for various distances and directions from the source. Because the horizontal acceleration produces the seiches, the short-period components of the seismic surface waves are very important. The tilt caused by the Rayleigh waves has been shown to be unimportant in causing seiches, especially for periods less than 600 seconds (McGarr, 1965, p. 851). The predominant surface-accelerations probably lie in the period range of 5 to 15 seconds. If everything else is equal, bodies of water with fundamental modes of oscillation in this period range should have the most numerous seiches.

In the Alaska earthquake of 1964, almost all of the known recorded seiches occurred in North America. Furthermore, most of the recorded seiches in North America were in the United States, most occurring in the Gulf Coast region. Our main attempt has been to explain the distribution of seiches in the United States because there we have the best data

control and the greatest density of records.

Throughout the United States the network of water-level recorders is reasonably well distributed. Our main assumption has therefore, been that, in a given geographical area containing a large number of them, a certain percentage of the water-level recorders are on bodies of water that are favorable for generating seiches. Because information about the size and shape of the various bodies of water is not readily available, such an assumption is the only realistic way to treat the data in a preliminary study such as this. Therefore, the problem of explaining the seiche distribution becomes one of identifying places where the horizontal components of the shorter period seismic surface waves were large enough in amplitude to provide a generating force. Other forces, such as seismic body waves, might induce seismic seiches, but preliminary studies imply that they are unimportant.

The fundamental hypothesis of this paper is that seiche distribution is a direct function of the amplitude distribution of Love and Rayleigh waves in a period range from 5 to 15 seconds. The occurrence of seiches is explained in terms of those waves, although surface-wave theory does not explain many features of the seiche distribution. The actual explanation may involve factors other than seismic surface waves or aspects of the behavior of surface waves that are not yet known. Perhaps this presentation of seiche data will promote further development of surface-wave theory.

LOCATION AND NATURE OF THE SEICHES

SEICHE DATA

The authors considered two types of data to ascertain seiche distribution: negative and positive. They did not examine the negative data, that is, the water-level records which showed no trace of a seismic seiche. A few recordings of seismic seiches may have been missed, but this source of error is not considered significant. All the recorded seismic seiches were examined by both

authors. The locations and double amplitudes of the seismic seiches in the conterminous United States and southern Canada are shown on plate 1.

The seiche data are summarized in table 1 by State or Province; data from gages on rivers and streams are grouped separately from those from gages on lakes, reservoirs, and ponds. The seiches recorded on rivers and streams generally were of short duration, lasting no more than 5 to 10


minutes. Seiches recorded in reservoirs, especially in the west, lasted for 2 hours or longer. The fluctuations decreased so gradually that the point of cessation of fluctuation and resumption of normal water level could not be distinguished on the records. These seiches lasted longer than stream seiches because reservoirs usually have much greater resonance qualities than other types of water bodies, as is discussed under "Hydrodynamic Factors" (p. E12).

TABLE 1.—Summary of 859 seismic effects from the Alaska earthquake on surface-water bodies throughout the world

| State or Province | On rivers and streams | | | | On lakes, reservoirs, and ponds | | | | Gages at time of earthquake | |
|---------------------|-----------------------|------------------------------------|---------------------------------------|---------|---------------------------------|------------------------------------|---------------------|---------|-----------------------------|----------------------------------|
| | Number recorded | Amplitude of maximum seiche (feet) | Discharge with seiche (cu ft per sec) | | Number recorded | Amplitude of maximum seiche (feet) | Storage (acre-feet) | | Number | Percent that recorded earthquake |
| | | | Maximum | Minimum | | | Maximum | Minimum | | |
| United States | | | | | | | | | | |
| Alabama..... | 24 | 0.22 | 109,000 | 11 | 5 | 0.18 | 1,100,000 | 120,000 | 103 | 28.1 |
| Alaska..... | 32 | ----- | 400 | 4 | 0 | ----- | ----- | ----- | 42 | 76.2 |
| Arizona..... | 6 | .02 | 260 | 3.1 | 2 | .35 | 14,952,000 | 77 | 119 | 6.7 |
| Arkansas..... | 36 | .48 | 58,000 | 1 | 5 | 1.45 | 1,970,000 | ----- | 89 | 46.0 |
| California..... | 8 | .05 | 1,580 | 15 | 19 | .42 | 3,257,100 | 4,000 | 661 | 4.1 |
| Colorado..... | 14 | .30 | 260 | .1 | 0 | ----- | ----- | ----- | 212 | 6.6 |
| Connecticut..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 70 | .0 |
| Delaware..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 6 | .0 |
| Florida..... | 97 | .66 | 26,800 | 2 | 3 | .04 | ? | ----- | 288 | 34.7 |
| Georgia..... | 28 | .22 | 43,000 | 100 | 0 | ----- | ----- | ----- | 75 | 37.4 |
| Hawaii..... | 5 | .17 | 302 | 7.4 | 0 | ----- | ----- | ----- | 146 | 3.4 |
| Idaho..... | 3 | .03 | 1,110 | 18 | 2 | .56 | 146,000 | ? | 191 | 2.6 |
| Illinois..... | 6 | .10 | 8,700 | 1,200 | 2 | .05 | ? | ? | 144 | 5.6 |
| Indiana..... | 13 | .39 | 15,000 | 35 | 3 | .07 | ? | ? | 131 | 12.2 |
| Iowa..... | 1 | ----- | 225 | ----- | 1 | .02 | ? | ----- | 129 | 1.6 |
| Kansas..... | 12 | .17 | 400 | .2 | 2 | .05 | 15,000 | 13,000 | 82 | 17.1 |
| Kentucky..... | 0 | ----- | ----- | ----- | 4 | .57 | 200,000 | 88 | 84 | 4.8 |
| Louisiana..... | 69 | .68 | 31,000 | .2 | 0 | ----- | ----- | ----- | 103 | 67.0 |
| Maine..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 52 | .0 |
| Maryland..... | 3 | .04 | ? | ? | 0 | ----- | ----- | ----- | 46 | 6.5 |
| Massachusetts..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 7 | .0 |
| Michigan..... | 13 | .10 | 860 | .8 | 3 | 1.83 | 30 | 21 | 140 | 11.4 |
| Minnesota..... | 1 | .03 | 5.0 | ----- | 0 | ----- | ----- | ----- | 91 | 1.1 |
| Mississippi..... | 22 | .37 | 22,500 | 24 | 0 | ----- | ----- | ----- | 61 | 36.1 |
| Missouri..... | 18 | .87 | 1,600 | 5 | 0 | ----- | ----- | ----- | 108 | 16.6 |
| Montana..... | 16 | .10 | 2,150 | 6 | 0 | ----- | ----- | ----- | 168 | 9.5 |
| Nebraska..... | 13 | .18 | 1,300 | 23 | 1 | .08 | 267,100 | ----- | 152 | 9.2 |
| Nevada..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 76 | .0 |
| New Hampshire..... | 1 | Tr. | 2,200 | ----- | 0 | ----- | ----- | ----- | 11 | 9.1 |
| New Jersey..... | 0 | ----- | ----- | ----- | 1 | .08 | 20,000 | ----- | 82 | 1.2 |
| New Mexico..... | 27 | .26 | 470 | 1 | 0 | ----- | ----- | ----- | 156 | 17.3 |
| New York..... | 4 | Tr. | 130 | 80 | 0 | ----- | ----- | ----- | 176 | 2.3 |
| North Carolina..... | 0 | ----- | ----- | ----- | 1 | .05 | 1,000,000 | ----- | 63 | 1.6 |
| North Dakota..... | 2 | .06 | 57 | 47 | 1 | ----- | 21,000 | ----- | 89 | 3.4 |
| Ohio..... | 16 | .14 | 1,650 | 11 | 9 | .25 | 60,600 | 1,500 | 188 | 13.3 |
| Oklahoma..... | 28 | .13 | 1,870 | .1 | 9 | .44 | 1,117,000 | 7,100 | 129 | 28.7 |
| Oregon..... | 10 | .14 | 21,000 | 2.8 | 7 | .11 | 272,000 | 18,000 | 239 | 7.1 |
| Pennsylvania..... | 2 | .05 | 1,400 | 7.7 | 0 | ----- | ----- | ----- | 108 | 1.8 |

TABLE 1.—Summary of 859 seismic effects from the Alaska earthquake on surface-water bodies throughout the world—Continued

| State or Province | On rivers and streams | | | | On lakes, reservoirs, and ponds | | | | Gages at time of earthquake | |
|----------------------------------|-----------------------|------------------------------------|---------------------------------------|---------|---------------------------------|------------------------------------|---------------------|---------|-----------------------------|----------------------------------|
| | Number recorded | Amplitude of maximum seiche (feet) | Discharge with seiche (cu ft per sec) | | Number recorded | Amplitude of maximum seiche (feet) | Storage (acre-feet) | | Number | Percent that recorded earthquake |
| | | | Maximum | Minimum | | | Maximum | Minimum | | |
| United States—Continued | | | | | | | | | | |
| Rhode Island..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 3 | 0.0 |
| South Carolina..... | 8 | .12 | 34,500 | 500 | 0 | ----- | ----- | ----- | 40 | 20 |
| South Dakota..... | 6 | .14 | 24,500 | 2 | 0 | ----- | ----- | ----- | 90 | 6.7 |
| Tennessee..... | 24 | .42 | 170,000 | 35 | 8 | .14 | 3,400,000 | 150,000 | 130 | 24.6 |
| Texas..... | 57 | .67 | 6,920 | .0 | 13 | .14 | 1,777,200 | 50 | 346 | 20.2 |
| Utah..... | 8 | .06 | 90 | 2 | 0 | ----- | ----- | ----- | 126 | 6.4 |
| Vermont..... | 0 | ----- | ----- | ----- | 2 | .23 | 29,000 | 8,500 | 8 | 25.0 |
| Virginia..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 155 | .0 |
| Washington..... | 6 | .45 | <10,000 | 6 | 15 | 1.04 | 6,900,000 | ? | 356 | 5.9 |
| West Virginia..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 91 | .0 |
| Wisconsin..... | 6 | .02 | 1,300 | 50 | 0 | ----- | ----- | ----- | 74 | 8.1 |
| Wyoming..... | 12 | .08 | 660 | 1 | 0 | ----- | ----- | ----- | 199 | 6.0 |
| Total..... | 658 | ----- | ----- | ----- | 118 | ----- | ----- | ----- | 6,435 | 12.0 |
| Puerto Rico..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 16 | 0.0 |
| Virgin Islands..... | 0 | ----- | ----- | ----- | 0 | ----- | ----- | ----- | 9 | .0 |
| Australia | | | | | | | | | | |
| Australia Capital Territory..... | 0 | ----- | ----- | ----- | 1 | Tr. | 21 | ----- | ----- | ----- |
| New South Wales..... | 0 | ----- | ----- | ----- | 1 | 0.02 | 23,680 | ----- | ----- | ----- |
| Northern Territory..... | 1 | 0.02 | ----- | ----- | 0 | ----- | ----- | ----- | ----- | ----- |
| Victoria..... | 1 | .02 | ----- | ----- | 0 | ----- | ----- | ----- | ----- | ----- |
| Total..... | 2 | ----- | ----- | ----- | 2 | ----- | ----- | ----- | ----- | ----- |
| Canada | | | | | | | | | | |
| Alberta..... | 28 | 0.31 | ----- | ----- | 0 | ----- | ----- | ----- | ----- | ----- |
| British Columbia..... | 4 | .29 | ----- | ----- | 23 | 3± | ----- | ----- | ----- | ----- |
| Northwest Territory..... | 5 | .15 | ----- | ----- | 2 | .30 | ----- | ----- | ----- | ----- |
| Ontario..... | 6 | .14 | ----- | ----- | 2 | .13 | ----- | ----- | ----- | ----- |
| Saskatchewan..... | 7 | .30 | ----- | ----- | 2 | .08 | ----- | ----- | ----- | ----- |
| Total..... | 50 | ----- | ----- | ----- | 29 | ----- | ----- | ----- | ----- | ----- |
| Grand total..... | 710 | ----- | ----- | ----- | 149 | ----- | ----- | ----- | ----- | ----- |




2.—The largest seiche recorded on a stream in each of eight States.

The seiches from the Alaska earthquake at surface-water gages that have been reported from throughout the world are separately listed and described in table 3 (p. E25); the station number, name, and location are those in current use.

Ideally, the table should give average depth and width of the body of water on which the seiche was observed. In their place a more easily obtained measurement is given, either the discharge in cubic feet per second ($\times 28.317 =$ liters per second) for flowing streams or acre-feet of water in storage ($\times 1,233.49 =$ cubic meters) for lakes, reservoirs, and ponds. The recorded seismically caused water-level motion is given under "seiche double amplitude." This amplitude may be less than the true amplitude because of the response of the gage. Furthermore, the fluctuations at the bubble-gages and at some of the float-gages were not symmetrical above and below the stage immediately prior to the seiche. For the asymmetrical double amplitudes, motion upward from prior stage is shown above a slash line and motion downward is shown below.

The largest seiche recorded on a stream in each of eight States is shown in figure 2. The largest one in California was only 0.05 feet (15 mm) in double amplitude. This seiche contrasts markedly both in size and duration with the seiches recorded in California reservoirs. The thinness of some of the pen lines on recorder charts suggests that there may have been only one or a very few oscillations associated with the seiche and that the oscillations were damped out almost immediately after passage of the seismic wave.

Some of the largest seiches recorded in reservoirs are shown in figure 3. Most of the seiches



3.—Some large seismic seiches on reservoirs.

shown continued for 2 hours or more, but the one for Wheeler Reservoir on the Tennessee River at Triana, Ala., lasted only about 40 minutes.


GAGING STATIONS, INSTRUMENTS, AND THEIR RECORDS

At the time of the Alaska earthquake, the Water Resources Division of the U.S. Geological Survey had about 8,150 recorders in operation, of which 6,435 were equipped to give a continuous record on which an event such as a seismic seiche could be recorded. Seiches were recorded on 763 charts. About half (356) were recorded in the States on or near the Gulf Coast and most distant from the epicenter, namely, Alabama, Arkansas, Florida, Georgia, Mississippi, Louisiana, and Texas (pl. 1).

The remaining 1,700 stations were equipped with a digital-type instrument that records a water-level measurement at 15-minute intervals and consequently cannot record any sudden changes such as seismic seiches. Because the trend currently is to install such instruments in place of the continuous-record type, the Alaska earthquake may be the last major earthquake for which seismic seiches can be widely recorded.

Seismic seiches were recognized on charts from three types of recorders, the continuous-analog, the bubble-gage, and the deflection-meter. The last records direction and velocity of flow and is used on streams and canals in Florida where stage-discharge relations that prevail elsewhere cannot be used, because gradients are so low and directions of flow vary with changing stages of the ocean tides.


Each type of gage and recorder has its special characteristics that




4.—Three types of bubble-gage records of Alaska earthquake seiches.

in part govern the kinds of seiche records that were obtained. Those characteristics and their effects were discussed in some detail by Vorhis (1967, p. C5, C6, C9). In brief, the continuous-analog records of stage generally are the most revealing. The movement tends to be symmetrical above and below the level prevailing before the onset of the seiches. Because of damping effects in the stilling wells in which the recorder floats operate, the fluctuations in stage recorded during seiches are smaller than the actual amplitudes of the seiche waves. There is no consistent degree of damping, for each installation has its individual character. Consequently, it is impossible currently to derive a factor by which

to convert recorded amplitude to true amplitude. The seiches illustrated in figures 2 and 3 are from continuous-analog recorders. The bubble gages have a built-in delay that may cause a seiche to be recorded as a brief or prolonged drop in stage or rise in stage or as an asymmetrical fluctuation (fig. 4). Simultaneous traces of stage and flow, recorded on continuous-analog charts in Florida, and the effects of the seiches are shown in figure 5.



5.—Seiche effects of Alaska earthquake on stage and flow, Miami area, Florida. A, Fluctuation in flow, no change in stage; B, fluctuation in stage, no change in flow; C, fluctuation in both stage and flow, "permanent" decrease in flow.



6.—Map of conterminous United States showing seiche density, in percent, by State and by river basin.

GEOGRAPHIC DISTRIBUTION

With the exception of four in Australia, three on the Island of Kauai, and two on the Island of Hawaii, all known seismic seiches caused by the Alaska earthquake were recorded at gaging stations in Canada and the continental United States. All data from other parts of the world were negative.

Seiche distribution was studied by areas, in terms of the percentage of the total number of gages that showed seiches. It was necessary to assume that all the charts had been examined and that the reported instrumentation of gaging stations was accurate. Neither assumption is entirely valid. Therefore, the method is

not highly precise, but it does permit a reasonably accurate comparison of seiche density by area.

The areas chosen are the major river basins within each State, that is, about 100 areas in the United States, for which percentage of seiche density could be computed. The map (fig. 6) presents the data. The percent values have been contoured to display the gross features of the distribution.

The southeastern part of the United States, notably, Louisiana, Arkansas, Florida, eastern Oklahoma, and eastern Mississippi, had by far the highest density of seiches. Other high-density areas include north-central New Mexico, eastern Kansas, and the area ad-

acent to the southern tip of Lake Michigan. The areas west of the Rocky Mountains, the area immediately to the east of the Rockies, and the Middle Atlantic States and New England experienced few or no seiches. Anomalous low-density areas occur in a strip along northwestern Mississippi, western Tennessee, and western Kentucky and in an area of southern Alabama. The distribution does not have any obvious dependence on distance or azimuth from the epicenter. On the other hand, the distribution seems to form definite regional patterns. It is highly improbable that these regional patterns have anything to do with the abilities of the individual bodies of water to couple

into the seismic waves. Possible controls over the distribution pattern are considered after the following discussion of hydrodynamic factors.

HYDRODYNAMIC FACTORS

Alaska earthquake seiches occurred in many different kinds of water bodies, including lakes, rivers, streams, ponds, and reservoirs, and in tanks that contained chemicals. Several factors influence the amplitude and duration of seiches in different types of fluid bodies affected by a given seismic surface wave. These factors include the regularity of the geometry, the depth, and the size of the fluid body as well as the physical characteristics of the fluid. The following discussion deals only with water. In principle, the exact response, including the effects of damping, can be calculated for a body of water of any shape and size. In this study, however, the necessary information was not available so calculations of various responses are only approximate.

Seismic surface waves excite maximum response in deep, regular bodies of water that have low-order odd modes (fig. 1) and periods of 5–15 seconds. These waves excite only odd-order seiches. Rivers and creeks are considered to be similar to the idealized channel for which the exact response is known. Assume a river with width L and average depth H . The approximate periods of the normal modes of the river are then given by

$$T_{2n+1} = \frac{1}{2n+1} \frac{2L}{\sqrt{gH}}; n=0,1,\dots$$

These periods are approximate to the extent that the river departs from the shape of the idealized channel. The theory for a long canal may also be applied in a rough fashion to a narrow lake or a lake with a narrow inlet. In fact, in this paper the cross section of any body of water is considered to be the cross

TABLE 2.—First-, third-, and fifth-order modes, in seconds, for seiches on water bodies with selected widths and depths

| Depth (meters) | Mode | Width (meters) | | | | | | |
|----------------|------|----------------|-------|-------|-------|-------|-------|-------|
| | | 5 | 10 | 20 | 40 | 60 | 100 | 200 |
| 1 | 1 | 3.2 | 6.3 | 12.7 | 25.3 | 38.0 | 63.3 | 126.6 |
| | 3 | ----- | ----- | 4.2 | 8.4 | 12.7 | 21.1 | 42.2 |
| | 5 | ----- | ----- | ----- | 5.1 | 7.6 | 12.7 | 25.3 |
| 2 | 1 | 2.2 | 4.5 | 9.0 | 17.9 | 26.9 | 44.8 | 89.7 |
| | 3 | ----- | ----- | 3.0 | 6.0 | 9.0 | 14.9 | 30.0 |
| | 5 | ----- | ----- | ----- | 3.6 | 5.4 | 9.0 | 17.9 |
| 4 | 1 | ----- | 3.2 | 6.3 | 12.7 | 19.0 | 31.6 | 63.3 |
| | 3 | ----- | ----- | ----- | 4.2 | 6.3 | 10.5 | 21.1 |
| | 5 | ----- | ----- | ----- | 2.5 | 3.8 | 6.3 | 12.7 |
| 6 | 1 | ----- | ----- | 5.2 | 10.3 | 15.5 | 25.8 | 51.6 |
| | 3 | ----- | ----- | ----- | 3.4 | 5.2 | 8.6 | 17.2 |
| | 5 | ----- | ----- | ----- | ----- | 3.1 | 5.2 | 10.3 |
| 10 | 1 | ----- | ----- | 4.0 | 8.0 | 12.0 | 20.0 | 40.0 |
| | 3 | ----- | ----- | ----- | 2.7 | 4.0 | 6.7 | 13.3 |
| | 5 | ----- | ----- | ----- | ----- | ----- | 4.0 | 8.0 |
| 20 | 1 | ----- | ----- | ----- | 5.7 | 8.5 | 14.1 | 28.4 |
| | 3 | ----- | ----- | ----- | 1.9 | 2.8 | 4.7 | 9.4 |
| | 5 | ----- | ----- | ----- | ----- | ----- | 2.8 | 5.7 |
| 30 | 1 | ----- | ----- | ----- | 4.6 | 6.9 | 11.6 | 23.1 |
| | 3 | ----- | ----- | ----- | ----- | ----- | 3.8 | 7.7 |
| | 5 | ----- | ----- | ----- | ----- | ----- | ----- | 4.6 |

section of an infinitely long channel. For instance, the normal modes of a cylindrical tank are given approximately by

$$T_{2n+1} = \frac{2D}{(2n+1)\sqrt{gH}}$$

where D is the tank diameter. Table 2 lists the periods for modes 1, 3, and 5 for various combinations of width and depth where depth represents the average depth of the cross section. Table 2 shows that there are many possible cross sections that will have at least one of the periods of the first three nonzero modes in the 5- to 15-second period range. The periods of table 2 were computed on the basis of assumed long wavelength; these assumptions are not entirely valid for places where the length is not much greater than the depth. For those places, the period of the table is an underestimate of the true period. Table 2 shows which dimensions are in the optimal range for producing seiches.

In general, the seiches having the highest amplitudes and longest durations occurred in reservoirs. The lowest amplitudes and shortest durations were on creeks and

small rivers, owing probably to the combination of shallowness and irregularity of cross section.

The dimensions of a few of the bodies of water for which seiches were recorded are known. In California, a seiche in the Isabella Reservoir lasted more than 3 hours. The recorder on this reservoir which is formed behind a dam, is near one end of the dam. The most likely cross section to consider seems to be that parallel to the dam; its length is about 300 meters and its average depth is roughly 15 meters. The approximate periods of the first three modes are $T=49$, 16, and 10 seconds. These periods are in the approximate range required for coupling into the seismic surface waves.

Two partly buried water-storage reservoirs at Lansing, Mich., recorded fluctuations of 22 inches and 15 inches shortly after the Alaska earthquake. The reservoir which recorded the 22-inch seiche is cylindrical; its depth is about 8 meters and its diameter is about 50 meters. The periods of the first two seiche modes for that

reservoir would be 11 and 4 seconds. The reservoir that had the 15-inch seiche is a rectangular prism whose length, width, and depth are about 130, 41, and 8 meters, respectively. If the seiche had water movement parallel to the length, then the first three modes had periods of 29, 10, and 6 seconds. If the seiche was parallel to the width, then the periods of the first two seiche modes were 9.2 and 3.1 seconds.

Two seiches, that lasted somewhat more than an hour each, were recorded in two drums of liquid ethylene (density=0.529 gm per cm⁻³) at the Louisiana Division of the Dow Chemical Co. in Plaquemine, La. The tanks are about 18 meters long and the average depth of the liquid was about 1.0 meter. The fundamental seiche mode would have had a period of about 10 seconds and the third mode a period of 3½ seconds.

Thus, in all examples where the size and shape of the body of liquid is known, and for which a seiche was recorded, at least

one of the first three seiche modes lies in the period range of 5 to 15 seconds. Modes which are of higher order cannot be expected to be important because of the factor $\frac{1}{2n+1}$ which occurs in equation (2).

For the purposes of this study, it would have been ideal if all the bodies of water had been of the same shape, size, and orientation. Then measurements of the seiche amplitudes would indicate only the distribution of seismic surface-wave acceleration. This ideal situation is not even approached, so some assumptions were necessary. As stated on page E6, one major assumption was that in an area having a large number of surface-water recorders, most of the recorders were able to record a marginally detectable seiche. If the seismic waves were amplified, a larger percentage of recorders would show a seiche. Conversely, if the seismic waves were attenuated, no seiches would have been generated or recorded. The data support these assumptions. To

make the data more homogeneous, little emphasis was placed on those from reservoirs and canals, which are such good resonators that any in any part of North America probably would have experienced a seiche at the time of the Alaska shock. The data considered most valid for deducing the seismic surface-wave horizontal-acceleration distribution are from creeks and small rivers, which are generally poor resonators. As table 2 shows, nearly all the bodies of water in this study (mostly small rivers and streams) have low-order modes whose periods are in the 5- to 15-second range.

The observed geographic distribution of seiches from the Alaska earthquake was apparently controlled both by geologic features and by certain characteristics of seismic surface waves. The two kinds of control will be discussed separately, but their effects are not wholly separable because the surface waves may be strongly modified by the geologic materials and structural features they traverse.

INTERPRETATION OF SEICHE DISTRIBUTION

RELATION TO GEOLOGIC FEATURES

The influence of major geologic features on the distribution of seiches became apparent when seiche locations were plotted on the tectonic map of the United States (U.S. Geol. Survey and Am. Assoc. Petroleum Geologists, 1962). A simplified version of this map is shown as plate 1.

SEDIMENT THICKNESS

In all but three areas of North America—the northeast end of the Mississippi Embayment, the

area near Miami, Fla., and the Great Valley of California—the density of seiches seems to be roughly proportional to the thickness of low-rigidity sediments. Extreme examples of this density distribution are shown by the concentration of seiches in the Mississippi Delta region along the Gulf Coast of Louisiana, where sediment thickness is maximum, and by near absence of seiches on the Canadian Shield, where sediments are almost nonexistent. Along the Gulf Coast eastward and westward from Louisiana the regular decrease in number of

seiches as the deposits become thinner is particularly striking. The anomalously high density of seiches near Miami and the anomalously low densities at the head of the Mississippi Embayment and in the Central Valley of California are discussed on pages E19 and E20.

THRUST FAULTS

Thrust faults apparently provide a favorable environment for the generation of seiches. The relationship is especially clear in Georgia, where seiches were recorded at gages on the Brevard Rome, Towaliga, and Whitestone

thrust faults; a cluster of 11 seiches in west-central Alabama may be related to extensions of these faults. The Ouachita Mountains and the Ridge and Valley Province of Tennessee and Alabama—regions where thrust faults are numerous—show high concentrations of seiches; the Ouachita area, in fact, has a density comparable to that of central Florida. In several other places seiches were recorded over possible extensions of known thrust faults: in Utah west of the Wasatch Mountains, in Montana below Hebgen Lake on the Madison River (Irving J. Witkind, oral commun., October 1966), in Wyoming at Moran on the Snake River, and at Valley on the South Fork of the Shoshone River.

BASINS, ARCHES, AND DOMES

The locations of many seiches seemingly were controlled by structural basins and uplifts.

In the Williston basin (pl. 1) a few large seiches occurred on the side toward the epicenter but most occurred on the southeast or "lee" side. The presence of Lake Michigan makes observation of seiches on the northwest side of the Michigan basin impossible, but small seiches were recorded on its lee side. Three small seiches in the northern part of the basin overlie and may have been related to a pronounced positive Bouguer anomaly as shown on the gravity map of Woollard and Joesting (1964).

The greatly elongated Appalachian basin (pl. 1) lies with its long axis about perpendicular to the great-circle path for surface waves that propagated from Alaska. In that basin, seiches were recorded only on the northwest side in a belt trending northeastward through Ohio. Perhaps the elongated shape focused waves less than did the nearly circular shape

of the Williston and Michigan basins, for only one seiche was recorded on the lee side of the Appalachian basin.

These major basins may have damped the surface-wave energy near the land surface, because the waves as they traveled beyond a basin were able to generate relatively few seiches until well beyond its limit. For example, southeast of the Appalachian basin, in Virginia, New Jersey, southeastern Pennsylvania, and most of North Carolina, no seiches were recorded, and only three seiches were recorded in Maryland, two of which were at the lower limit of perceptibility.

A large seiche occurred on the Wichita Mountain uplift in southwestern Oklahoma and another good-sized one on its lee side, but from there to the Gulf Coast none was recorded in the 375-mile-long drainage basin of the Trinity River although many recorders were in operation and although some of the largest seiches were recorded in rivers on the flanks of the Trinity basin. Thus it seems that the Wichita Mountain uplift and possibly the Muenster arch shielded the Trinity River from surface waves and left it in a shadow zone of little or no seismic intensity. The Adirondack uplift also seems to have acted either as a shield or a deflector, for the data indicate a shadow zone to the southeast of it.

The elongated Arkoma basin (pl. 1) had abundant seiche activity throughout, at about the same positions with respect to the base of the Pennsylvanian rocks as in the Appalachian basin. Because the Arkoma basin trends in roughly the same direction as the Appalachian basin with respect to surface-wave propagation paths from Alaska, the same factors may account for the similar seiche distribution in both basins. In the

Delaware basin, seiches were concentrated along the northeast side, and in the San Juan basin along the northern and eastern edges. The Black Warrior basin had many seiches along its northwest and northern edges.

In the Nashville dome area, a fairly large number of seiches were recorded. Because all but one of the seiches in that area were on large rivers, however, there may be little or no geological significance to this seiche concentration. Many basins, domes, and arches did not seem to control seiche distribution, perhaps because they are much smaller than those named above.

EDGE OF OVERLAPS

The feather edges of sediments deposited by marine invasions seem to have been areas favorable for the generation of seiches. Seven seiches occurred along the edge of the Cretaceous overlap in Oklahoma and Arkansas although they may have been related to thrust faults, synclines, and compressed anticlines that extend below the overlap. In Tennessee and Alabama, six seiches occurred along the edge of the Cretaceous overlap, and three more were recorded along its edge in Georgia and South Carolina, only one of which may also be associated with a thrust fault.

ROCKY MOUNTAIN SYSTEM

In the western United States most of the seiche activity seems to be related to the Rocky Mountain tectonic belt (pl. 1). Apparently the surface waves traveled along the Rockies and produced seiches wherever they met an irregularity in the wave guides, such as the Sangre de Cristo uplift and the White River uplift. Other areas in the Rockies where many seiches were noted include much-faulted areas in north-

central Utah, southwestern Montana, and east-central Arizona. By acting as a wave guide, the Rocky Mountains seemingly channeled so much energy along the mountains that a shadow zone, shown on plate 1, was created along the foot of the Rocky Mountains from Canada to the Gulf of Mexico.

MISCELLANEOUS AREAS

By far the greatest density of seiches in North America was recorded in the Miami area of Florida. Most of the seiches occurred on the canals that lace the region. The sedimentary deposits there are relatively thin compared to those on many parts of the Gulf Coast that had much lower seiche densities. The high density around Miami may have been due to the fact that most canals are of optimum size and shape for coupling into seismic surface waves. Because their geometrical shapes are better defined than those of most rivers, canals are presumably much better resonators.

Many seiches were recorded on the western edge of the Sierra Nevada batholith, mostly in reservoirs and lakes. The Sierra Nevada and the Cascades may form a continuous wave guide for surface waves, similar to the one along the Rocky Mountains.

RELATION TO SEISMIC SURFACE WAVES

A basic thesis of this paper is that the distribution of seiches corresponds directly to horizontal acceleration by seismic surface waves whose periods range from 5 to 15 seconds. The only waves that can provide sufficient horizontal acceleration are the fundamental-mode Love and Rayleigh waves. Such waves with periods of less than 5 seconds do not propagate efficiently at teleseismic dis-

tances, and waves with periods longer than 15 seconds produce little acceleration. Factors that determine the relative horizontal acceleration at a given point for the surface waves with periods that range from 5 to 15 seconds may include (1) nature of the radiation pattern, (2) distance from the epicenter, (3) focusing and defocusing of the surface waves by lateral refraction, (4) local crustal structure, especially the thickness of surficial sediments of low rigidity, and (5) structural irregularity of the crust. The relative importance of these factors must be considered in the light of the seiche data that have been studied.

RADIATION PATTERN

The radiation pattern of surface waves from the Alaska earthquake cannot be ascertained from seismograms because nearly all long-period seismographs were driven off scale. However, a study of the aftershocks, which according to Stauder and Bollinger (1966) had fault-plane solutions similar to those for the main shock, indicates that whatever surface-wave radiation pattern existed did not noticeably affect the horizontal acceleration of surface waves throughout the United States.

Data from two aftershocks (nos. 17 and 21 in table 1 of Stauder and Bollinger, 1966), as recorded at each of the World-wide Standard Seismograph Network stations (WWSSN) in the United States, were used to determine the maximum horizontal displacement in the period range of 5 to 15 seconds on the two horizontal long-period seismograph components. These displacements were added vectorially and divided by the square of their period to derive a value that is proportional to acceleration. The values were then adjusted to account for the different


gain settings at each station. The resulting values, (\bar{u} in fig. 7) indicate the relative distribution of horizontal acceleration from the main shock of the earthquake, based on the assumption that the selected aftershocks and the main shock had similar patterns of surface-wave radiation.

The distribution of \bar{u} values does not seem to correlate with the distribution of seiches, partly perhaps because there are too few WWSSN stations, but partly because an ideal site for a seismograph station is a poor location for the generation of a seiche. At most seismograph sites low-rigidity sediments are thin or absent. The only major exception is the station at Spring Hill, Ala., which is in a region where no ideal seismograph site was available. The Spring Hill station record yielded the largest value of \bar{u} calculated in this study. This high value corresponds to the high seiche density along the Gulf Coast. The relation of seiche density to sediment thickness is discussed further on page E18.

The fact that both Love and Rayleigh waves produce horizontal acceleration also tends to diminish the importance of the radiation pattern because the radiation patterns of Love and Rayleigh waves are generally different. The aftershock records indicate that in the United States short-period Rayleigh waves had slightly larger amplitudes than did the Love waves. Thus, within North America, the radiation pattern was probably not an important factor in determining seiche distribution.

DISTANCE FROM EPICENTER

If the crustal wave guide were perfectly homogeneous and elastic between the epicenter and a given point, then any frequency component of the surface waves would



7.—Maximum horizontal acceleration (\ddot{u}) at stations of the World-wide Standard Seismograph Network in the United States calculated for two aftershocks of the Alaska earthquake.

decrease in amplitude according to $1/\sqrt{\sin \Delta}$, because of geometrical spreading on a sphere. The effect of this decrease is probably unimportant within North America in comparison with other factors. In theory, this effect would cause the surface-wave amplitude 10° from the epicenter to be about twice as large as the amplitude at the tip of Florida. The seiche data definitely do not suggest such a relation. Seismograms of Alaskan aftershocks indicate similarly that these smaller earthquakes in the epicentral region of the main shock sent out surface waves that did not diminish materially with distance within North America (fig. 7).

The effect of dispersion of seismic surface waves on seiche amplitudes is not well understood. In theory, surface-wave trains decrease in amplitude proportionally to either $1/\sqrt{\Delta}$ or $1/\sqrt[3]{\Delta}$ because of dispersion. This effect was seemingly unimportant in determining the amplitude distribution of either the seiches or the aftershocks.

LATERAL REFRACTION

The seiche data suggest that lateral refraction of seismic surface waves occurred in some areas. Exact theoretical calculation of this effect is impossible because detailed knowledge is lacking on phase velocity of surface waves in

North America. An example of lateral refraction was the apparent concentration of seismic energy along the Rocky Mountains (pl. 1, fig. 6). This effect could have been predicted qualitatively on the basis of work by John T. Kuo on distribution of phase velocity (fig. 8). Although the map shows contours of phase velocity for waves with a period of 20 seconds, it is probably also a valid guide to the relative distribution of velocity of the 5- to 15-second period waves considered in the present paper. According to geometrical ray theory, energy would have been concentrated in the low-velocity channel down the axis of the Rockies that is nearly parallel to

gated very inefficiently across the Appalachian basin as demonstrated by the few seiches recorded east of the mountains. In contrast, the long-period waves were not similarly affected, for in New Jersey alone 40 ground-water observation wells recorded hydro-seisms from the earthquake.


Large circular basins seem to be capable of focusing surface-wave energy. In the Michigan and the Williston basins the seismic surface waves traveled from northwest to southeast. The fact that local concentrations of seiches occurred on the southeast sides of the basins suggests that seismic energy was focused by the lenticular shape of the sedimentary basin fill. Because the sediments are deepest in the center of a basin, the local phase velocity of the surface waves would be smallest at the center and would increase with distance from the center of the basin. Geometrical ray theory indicates that wave crests, which were parallel while the waves were still northwest of the basin, would cross each other to the southeast of the basin and would produce amplification there. The analogous situation for water waves passing over a circular shoal was shown by Stoker (1957, p. 135).

In summary, lateral variations in phase velocity appeared to channel seismic energy along geosynclinal belts and focus energy on the lee sides of basins.

LOCAL CRUSTAL STRUCTURE

The thickness of sediments of low rigidity seems to be an important cause of amplification of horizontal motion resulting from surface waves. The following examples indicate the type of amplification this mechanism may produce.

Application of an approximate theory of Rayleigh-wave trans-



9.—Amplification of Rayleigh-wave displacements $\frac{u'}{u}$ and $\frac{w'}{w}$ (also accelerations $\frac{\ddot{u}'}{u}$ and $\frac{\ddot{w}'}{w}$) in low-rigidity sediment overlying high-rigidity rock, for (A) 15- and (B) 8-second period waves.

mission and reflection developed by McGarr and Alsop (1967) shows (fig. 9) the amplifications of horizontal and vertical components of motion of 15- and 8-second period Rayleigh waves that have crossed a structural boundary. In those examples,

waves traveling in a Canadian Shield model (Brune and Dorman, 1963) are incident on a model in which the upper part has been replaced by a layer of elastic surficial sediments. The layer has a compressional velocity, α , of 3 km sec⁻¹, a shear velocity, β , of

1.55 km sec⁻¹ and a density, ρ , of 2.17 gm cm⁻³. The thickness of the layer ranges from $H=0$ to $H=6.0$ km. As shown in figure 9, an amplification of as much as 2.5 can be provided by a thick layer of sediments. This mechanism for amplification of surface horizontal displacement and acceleration predicts that the density of occurrence of seiches will be approximately proportional to the thickness of the elastic sedimentary layer. This theory seems to agree well with the density of seiches along the Gulf Coast.

In the northeast part of the Mississippi Embayment, however, the theory is less well substantiated, for the seiche density was much lower in the embayment where sediments are thick than in the surrounding areas (pl. 1, fig. 6). We have considered the possibility that the theory for normal-mode surface waves may explain the apparent attenuation of horizontal acceleration in the areas of extremely low rigidity sediments such as may be found in that part of the Mississippi Embayment.

Figure 10 (next page) shows the variation in amplitude of surface horizontal acceleration (which is proportional to the amplitude of surface horizontal displacement) as a function of "layer" shear velocity for 6- and 10-second period Rayleigh waves propagating in a crustal model. This crustal model has the same structure as the Canadian Shield except that the upper 1 km has been replaced by a layer with a compressional-wave velocity of 3.0 km sec⁻¹, a density of 2.3 gm cm⁻³, and a shear velocity that ranges from 1.0 to 0.1 km sec⁻¹. The horizontal displacement has been normalized, so all the waves of a given period transport the same amount of energy. For reference, the horizontal acceleration

produced by 6- and 10-second waves in an unmodified Canadian Shield model are -0.94 and -0.93 (expressed in the same relative units used in fig. 10). If only the waves of 10-second period are considered, then low horizontal acceleration would result if the shear velocity were in a narrow region near 0.475 km sec⁻¹. However, the 6-second waves have a horizontal displacement of more than 2 for $\beta=0.475$. Similarly, the value for the 6-second waves is zero where the 10-second waves provide a horizontal acceleration of more than 1.5. We are considering a band of periods between 5 and 15 seconds and low accelerations for the entire band, or even for a large fraction of the band, obviously will not occur where shear velocities are greater than 0.1 km sec⁻¹. Thus, ordinary surface-wave theory does not seem to explain the low seiche density observed in the northeastern part of the Mississippi Embayment.

The data suggest that the boundary between hard and soft material and possibly the finite extent of the sediments must be considered in any theory that seeks to explain phenomena such as those observed in the upper Mississippi Embayment.

In summary, sediments of low rigidity seem to be capable of amplifying or, in isolated cases, attenuating the horizontal acceleration of surface waves. Surface-wave theory can predict the amplification of horizontal acceleration for crustal models having a surficial layer of elastic sediments, but it cannot predict attenuation.

IRREGULAR STRUCTURES

Short-period surface waves are generally observed to travel more efficiently parallel to tectonic features than perpendicular to them (Sutton and others, 1967). Waves traveling in a direction perpen-


dicular to a tectonic trend are attenuated rather rapidly, although the mechanism of attenuation is not understood at present (Richter, 1958, p. 143). The distribution of seiches indicates that, in addition, the horizontal displacement of short-period surface waves is amplified in regions of rapidly changing crustal structure, especially where surface waves travel across structural features in a direction normal to their trends.

In the Appalachian basin, nearly all of the seiche activity occurred on the northwest side of the basin; there was a pronounced shadow zone to the southeast. Seiche activity was strongest in the region where the beds begin to dip under the Appalachian basin. In Ohio, there is a belt of activity parallel to the contacts of Pennsylvanian beds that dip under the basin.

In the Valley and Ridge province of southern Tennessee, the areas of high seiche density coincide with surface contacts of southeast-dipping beds and with traces of thrust faults. There is no pronounced shadow zone on the lee side of the tectonic belt; rather, the seiche activity seems to continue at a somewhat diminished, but constant, level across Georgia and South Carolina to the coast. The Arkoma basin did not produce a shadow zone, perhaps because it is narrower and not nearly as deep as the Appalachian basin.

In summary, beds that thicken in the direction of wave propagation seem locally to amplify the horizontal acceleration of seismic surface waves; extremely deep sedimentary basins may attenuate short-period surface waves and thus cause shadow zones.

The continental margin also appears to attenuate short-period waves. Great-circle paths from the



10.—Variation in amplitude of surface horizontal acceleration, as a function of “layer” shear-wave velocity, for 6- and 10-second period Rayleigh waves propagating in the modified Canadian Shield model discussed in the text.

epicenter of the Alaska earthquake to all of California and parts of Oregon, Washington, and Nevada cross part of the Pacific Ocean. The data suggest that seiches in that part of the United States occurred for the most part only on bodies of water, such as reservoirs, that were capable of coupling into rather long-period seismic surface waves. Otherwise, the Central Valley of California might have had a very high seiche density because of its thick filling of low-rigidity sediments.

SEICHES AND SEISMIC INTENSITY

According to Richter (1958, p. 140), a passable relation between ground acceleration and the modified Mercalli intensity scale is given by the expression $\log a = \frac{I}{3} - \frac{1}{2}$ where I is the intensity and a is the acceleration in centimeters per second per second. Because both seiches and seismic intensity are related to horizontal ground acceleration, the authors investigated the possibility of


using seiches in seismic-intensity studies. Richter (1958, p. 138) included seiche occurrence among the long-period intensity effects. Distribution of analog water-level recorders in the United States is now sufficiently dense that their records might be a more reliable indication of intensity than eyewitness reports, at least in some situations.

The seiche distribution from a major shock, such as the Alaska earthquake, might also be used to predict the potential distri-

bution of intensity in areas before a local earthquake occurred. To find out how effectively seiche distribution from the Alaska earthquake might be so used, the seiche distribution was plotted on an intensity map (prepared by Kisslinger and Nuttli, 1965) of the south-central Missouri earthquake of October 21, 1965. All seiches resulting from the Alaska shock, which occurred within the perceptibility ellipse of the Missouri shock, were plotted to see whether or not seiche distribution was correlated with ground response to horizontal acceleration caused by local shocks (fig. 11). Several features of the intensity map could have been predicted from the seiche distribution. Both the seiche distribution and the local-intensity were anomalously low in the Mississippi Embayment. A local high in seiche density occurred near the axis of the perceptibility ellipse, about 125 km northwest of the epicenter. There was a local high in both seiche density and local-shock intensity at the southeast end of the ellipse, which is also on the southeast side of the embayment.

Some features of the intensity map, of course, would not have been predicted from study of the seiche distribution, possibly because:

1. Seiches from the Alaska shock were caused by seismic surface waves having periods greater than 5 seconds, whereas most intensity effects are caused by seismic waves having periods of less than 1 second.
2. The direction of wave propagation seems to have a strong effect. High correlations occurred northwest and southeast from the epicenter, that is, parallel or antiparallel to the waves from the Alaska



EXPLANATION

| | | |
|--------------------------------------|-------------------------|---|
| ----- | ----- | ○ |
| Boundary of region of perceptibility | "Not felt" zone | Seismic seiche of the Alaska earthquake, March 28, 1964 |
| ----- | △ | II |
| Perceptibility ellipse | Reported damage | Intensity (Modified Mercalli scale) |
| ----- | × | |
| Main region of reported damage | Not felt near epicenter | |

11.—Alaska earthquake seiches plotted on the intensity map of the Missouri earthquake of October 21, 1965.

shock. Perhaps if the seiche distribution which resulted from waves traveling from the northwest could be combined with the distribution of seiches resulting from waves propagated either from

the southwest or from the northeast, we would be able to predict potential seismicity more precisely for any area desired.

Apparent attenuation of seismic intensity, such as occurred in the

Mississippi Embayment, seems to occur in other areas as well. Richter (1958, p. 143) stated that where seismic waves emerge from hard rock into alluvium or unconsolidated sediments there is con-

siderable absorption, accompanied by increase of local intensity. This statement was based largely on observations of seismic intensity in California. It agrees with the seiche distribution in the Missis-

sippi Embayment for an unusually high number of seiches occurred at the northwest edge of the embayment along the Tertiary overlap, but there were almost none across the rest of the embayment.

CONCLUSIONS AND RECOMMENDATIONS

The factors of greatest influence on the distribution of short-period seismic surface-wave amplitudes seem to be (1) local crustal structure, especially the thickness of surficial material of low rigidity, (2) tectonic trends, (3) homogeneity of the path of surface-wave travel from the epicenter to a given locale, and (4) focusing of surface-wave energy by lateral phase velocity variations. Epicentral distance and radiation pattern seem to be of little importance.

There may be other controls on the seismic amplitude distribution. In areas of soft sediments, such as the Gulf Coast, there may have been horizontal displacements of as much as 10 cm due to the surface waves. If the period of the waves was as short as 6 seconds, then the horizontal displacement at land surface was about 0.01 of gravity. Locally, this displacement may have been sufficient to cause inelastic effects, some of which may correspond to the square symbols on plate 1.

There seems to be a correlation between the distribution of seiches and the potential intensity of a local earthquake in a given region. If seiches are indeed valid indicators of potential intensity, then an earthquake of a given magnitude in Louisiana might be of greater intensity than one of comparable magnitude at any other location in North America.

The distribution of seiches may contain implications that will lead

to further developments in seismic surface-wave theory. For instance, the seiche distribution resulting from the Alaska earthquake suggests that:

1. Unusually large horizontal amplitudes of short-period seismic surface waves occur in areas where absorption of the waves is most rapid. Waves that travel transverse to tectonic trends produce large horizontal amplitudes in the vicinity of the trend.
2. Lateral variations of local-phase velocity can focus and channel surface waves.

If the assumptions made in this study are valid, then analog water-level recorders are a valuable tool both for the theoretical and for the disaster-prevention aspects of seismology because the recorders are equivalent in many respects to a relatively dense network of horizontal accelerometers. For further study of seismic seiches, the authors recommend that:

1. A network of analog water-level recorders be maintained throughout the United States, or preferably throughout the world.
2. Analog recorders with an expanded time scale be maintained on selected bodies of water in areas of high seismicity.
3. Seismographs be installed on appropriate tectonic features to permit study of the local amplification of surface waves

such as is suggested by the seiche data.

4. Seiche recordings for smaller magnitude shocks be collected to investigate the possibility of a relation between seiche distribution and earthquake magnitude.
5. Seiches or their absence in epicentral areas be studied as a potentially reliable method for measuring earthquake intensity.

Because this study of seiches resulting from a major earthquake is the first of its type, the interpretations must be regarded as preliminary. Furthermore, the seiche data have not been used fully, for little attention was paid to amplitudes, periods, or durations. Most of the interpretation is based on the number of seiches that were recorded in a given region compared to the number of recorders in operation. Because of the great variation in response at the various recording sites and because more than 750 seiches were recorded in the United States, it seemed prudent to keep the data analysis relatively simple. In the future, it may be possible to analyze the records of seiche amplitudes from sites where the response to seismic surface waves can be calculated. Bodies of water with well-known regular shapes, such as canals and reservoirs, would be the best sites for such studies.

REFERENCES

- Brune, James, and Dorman, James, 1963, Seismic waves and earth structure in the Canadian Shield: *Seismol. Soc. America Bull.*, v. 53, no. 1, p. 167-209.
- Callahan, J. T., 1964, The yield of sedimentary aquifers of the Coastal Plain, Southeast River Basins: U.S. Geol. Survey Water-Supply Paper 1669-W, p. W1-W56.
- Donn, W. L., 1964, Alaskan earthquake of 27 March 1964—remote seiche stimulation: *Science*, v. 145, no. 3629, p. 261-262.
- Donn, W. L., and Posmentier, E. S., 1964, Ground-coupled air waves from the great Alaskan earthquake: *Jour. Geophys. Research*, v. 69, no. 24, p. 5357-5361.
- Forel, F. A., 1895, *Le Léman—Monographie limnologique*: Lausanne, F. Rouge, v. 2, 651 p.
- Kisslinger, Carl, and Nuttli, O. W., 1965, The earthquake of October 21, 1965, and Precambrian structure in Missouri: *Earthquake Notes*, v. 36, nos. 3-4, p. 32-36.
- Kvale, Anders, 1955, Seismic seiches in Norway and England during the Assam earthquake of August 15, 1950: *Seismol. Soc. America Bull.*, v. 45, no. 2, p. 93-113.
- McGarr, Arthur, 1965, Excitation of seiches in channels by seismic waves: *Jour. Geophys. Research*, v. 70, no. 4, p. 847-854.
- McGarr, Arthur, and Alsop, L. E., 1967, Transmission and reflection of Rayleigh waves at vertical boundaries: *Jour. Geophys. Research*, v. 72, no. 8, p. 2169-2180.
- Mercanton, Paul-Louis, 1946, *Le sisme du 25 janvier 1946—son effet sur les lacs suisses*: *Soc. Vaudoise Sci. Nat. Bull.*, v. 63, no. 267, p. 321-323.
- Miller, W. D., and Reddell, D. L., 1964, Alaskan earthquake damages Texas High Plains water wells: *Am. Geophys. Union Trans.*, v. 45, no. 4, p. 659-663.
- Oliver, Jack, and Ewing, W. M., 1958, The effect of surficial sedimentary layers on continental surface waves: *Seismol. Soc. America Bull.*, v. 48, no. 4, p. 339-354.
- Piper, A. M., 1933, Fluctuation of water surface in observation wells and at stream-gaging stations in the Mokelumne area, California, during the earthquake of December 20, 1932: *Am. Geophys. Union Trans.*, 14th Ann. Mtg., p. 471-475.
- Richter, C. F., 1958, *Elementary seismology*: San Francisco, Calif., W. H. Freeman and Co., 768 p.
- Sever, C. W., 1966, Miocene structural movements in Thomas County, Georgia, in *Geological Survey research, 1966*: U.S. Geol. Survey Prof. Paper 550-C, p. C12-C16.
- Stauder, William, and Bollinger, G. A., 1966, The focal mechanism of the Alaska earthquake of March 28, 1964, and of its aftershock sequence: *Jour. Geophys. Research*, v. 71, no. 22, p. 5283-5296.
- Stermitz, Frank, 1964, Effects of the Hebgen Lake earthquake on surface water, in *Effects of the Hebgen Lake, Montana, earthquake of August 17, 1959*: U.S. Geol. Survey Prof. Paper 435, p. 139-150.
- Stoker, J. J., 1957, *Water waves; the mathematical theory with applications*: New York, Interscience Publishers, Inc., 567 p.
- Sutton, G. H., Mitronovas, Walter, and Pomeroy, P. W., 1967, Short-period seismic energy radiation patterns from underground nuclear explosions and small-magnitude earthquakes: *Seismol. Soc. America Bull.*, v. 57, no. 2, p. 249-267.
- U.S. Coast and Geodetic Survey, 1945, *United States earthquakes, 1943*: Washington, U.S. Govt. Printing Office, 47 p.
- U.S. Geological Survey and American Association of Petroleum Geologists, 1962, *Tectonic map of the United States, exclusive of Alaska and Hawaii*, scale 1:2,500,000.
- Vorhis, R. C., 1967, Hydrologic effects of the earthquake of March 27, 1964, outside Alaska, *with sections on Hydroseismograms from the Nunn-Bush Shoe Co. well, Wisconsin*, by Elmer E. Rexin and Robert C. Vorhis, and Alaska earthquake effects on ground water in Iowa by R. W. Coble: U.S. Geol. Survey Prof. Paper 544-C, p. C1-C54.
- Wigen, S. O., and White, W. R. H., 1964, Tsunami of March 27-29, 1964, west coast of Canada: Canada Dept. of Mines and Tech. Surveys, 12 p.
- Wilson, B. W., 1953, Coastal seiches, pt. 1 of *Oscillations of the sea and the phenomenon of range*: The Dock and Harbour Authority [London], June, p. 41-45.
- Woollard, G. P., chm., and Joesting, H. R., coordinator, 1964, *Bouguer gravity anomaly map of the United States (exclusive of Alaska and Hawaii)*: *Am. Geophys. Union and U.S. Geol. Survey*, scale 1:2,500,000.

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages

[North latitude, west longitude, unless otherwise indicated. Time: March 28, 1964, Greenwich civil time. Discharge (in cubic feet per second) in roman type, storage (in acre feet) in *italic*; for asymmetrical double amplitudes, motion upward is shown above a slash line and motion downward is shown below. Latitude and longitude in degrees, minutes, and seconds where the location has been accurately determined; in degrees and minutes or in degrees only where location is less certain. Datum is altitude of an arbitrary point at each gaging station below the lowest level to which streamflow is likely to fall and from which all stage levels at a station are measured; altitude of the water surface above sea level is the sum of the stage plus altitude of the datum. Time is given mainly to indicate that the reported fluctuation occurred at about the time the seismic waves arrived. Many of the times as given might be subject to some correction if the entire chart could be examined for systematic clock error]

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|----------------------|--|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| UNITED STATES | | | | | | | | | |
| Alabama | | | | | | | | | |
| 2-3440 | Chattahoochee River at Alaga | 31°07' | 85°03' | 62.72 | 19.50 | 04:00 | 40,000 | 0.18 | Seiche lasted about 30 min. |
| 2-3785 | Fish River near Silver Hill | 30°32'45" | 87°47'55" | 20 | 1.93 | 03:50 | 75 | .03 | |
| 2-3995 | Cosa River at Weiss Dam at Leesburg | 34°11' | 85°45' | 517.77 | 68.42 | 04:00 | | .15/.00 | On Rome fault. Bubble gage. |
| 2-4001 | Terrapin Creek at Ellenville | 34°04' | 85°37' | 539.07 | 9.45 | 04:10 | 1,750 | .13 | In Coosa syncline and on a possible extension of a thrust fault. |
| 2-4015 | Big Cane at Gadsden | 33°54'11" | 86°06'37" | 490.56 | 12.65 | 04:00 | 3,900 | .10 | On a thrust fault. |
| 2-4120 | Tallapoosa River near Heflin | 33°37' | 85°31' | 830 | 17.20 | 03:45 | 6,400 | .12 | On Whitestone thrust fault. |
| 2-4285 | Flat Creek at Fountain | 31°37' | 87°25' | 45.43 | 2.68 | 04:10 | 240 | .12 | |
| 2-4295 | Alabama River at Claiborne | 31°32' | 87°31' | .4 | 40.7 | 04:15 | 109,000 | .18 | On possible extension of fault zone. |
| 2-4380 | Buttahatchee River below Hamilton | 34°06' | 87°58' | 360.80 | 5.30 | 04:00 | 1,350 | .22 | Fault(?) buried under Cretaceous overlap. |
| 2-4420 | Luxapilla Creek near Fayette | 33°43' | 87°52' | 322.33 | 1.60 | 02:40 | 280 | .03 | On possible extension of a buried fault. |
| 2-4450 | Lubbub Creek near Carrollton | 33°15' | 88°05' | 174.24 | 6.40 | 04:00 | 345 | .05 | On crest of compressed anticline. |
| 2-4451.55 | Tombigbee River at Epes | 32°41'45" | 88°06'55" | | 36.90 | 04:00 | | .12 | On west edge of buried Appalachian front. |
| 2-4565 | Locust Fork at Sayre | 33°42'35" | 86°59'00" | 258.64 | 21.00 | 04:00 | 13,500 | .20 | On an echelon fault. |
| 2-4645 | North River near Tuscaloosa | 33°21'10" | 87°33'25" | 155.24 | 2.93 | 04:10 | 840 | .08 | |
| 2-4670 | Tombigbee River at Demopolis Lock and Dam near Coatopa | 32°31'15" | 87°52'05" | 56.00 | 37.40 | 04:00 | 78,000 | .06/.10 | On possible extension of Appalachian faults. |
| 2-4680 | Alamuchee Creek near Cuba | 32°26' | 88°20' | 161.50 | 2.53 | 04:00 | 92 | .04 | On west edge of buried Appalachians. |
| 2-4695 | Tuckabum Creek near Butler | 32°11' | 88°10' | | 1.94 | 03:45 | 170 | .10 | On possible extension of Appalachian faults. |
| 2-4695.5 | Horse Creek near Sweetwater | 32°03' | 87°52' | 130 | 2.55 | 04:05 | 62 | .07 | On possible extension of a buried fault. |
| 2-4696 | Bashi Creek near Campbell | 31°56' | 87°59' | | 4.92 | 04:10 | 205 | .11 | Do. |
| 2-4700 | Tombigbee River near Leroy | 31°34' | 88°02' | 7.28 | 35.4 | 04:30 | 180,000 | .18 | On Hatchetigbee anticline. Bubble gage. |
| 2-4701 | East Bassett Creek near Walker Springs | 31°32' | 87°47' | 60.02 | 3.40 | 04:30 | 300 | .10 | On fault zone. |
| 2-4710.65 | Montilmar Creek at U.S. Hwy 90 at Mobile | 30°39'03" | 88°07'28" | | 2.38 | 04:00 | 11 | .05 | |
| 2-4795 | Escatawpa River near Wilmer | 30°52' | 88°25' | 60 | 5.23 | 04:15 | 720 | .08 | On Wiggins uplift. |
| 3-5853 | Sugar Creek near Goodsprings | 34°56'40" | 87°09'20" | 575 | 4.25 | 04:10 | 460 | .05 | |
| 3-5905 | Tuscumbia Spring at Tuscumbia | 34°43'45" | 87°42'15" | 409.65 | 9.03 | 04:15 | 121 | .06 | A residual 0.02-ft. rise in stage. |
| 3-5923 | Little Bear Creek at Halltown | 34°29'19" | 88°02'07" | 499.30 | 4.10 | 03:20 | 380 | .06 | |
| | Tennessee River at Waterloo | 34° | 88° | | | 04:15 | 900,000 | .03 | A residual 0.01-ft. drop in stage. |
| | Tennessee River at Triana | 34° | 86° | MSL | 559.78 | 04:35 | 1,100,000 | .18 | |
| | Tennessee River near Smithsonia | 34° | 87° | | 12.60 | 04:00 | 900,000 | .07 | Seiche lasted about 50 min. |
| Alaska | | | | | | | | | |
| 30-0115 | Red River near Metlakatla | 55°08'29" | 130°31'50" | 5 | 2.72 | 03:45 | 140 | 0.15 | Tsunami crests were recorded at 03:30, 10:00, 11:50, 21:20, and 22:20. |
| 30-0120 | Winstanley Creek near Ketchikan | 55°25'00" | 130°52'05" | 290 | 1.51 | 03:30 | 50 | .12 | |
| 30-0201 | Tyee Creek near Wrangell | 56°12'54" | 131°30'25" | 4.62 | 1.05 | 03:55 | 22 | .12 | Tsunami waves superimposed on high tide. |
| 30-0220 | Harding River near Wrangell | 56°13' | 131°38' | 20 | 4.65 | 04:00 | 100 | No seiche | Water rose 0.02 ft. in 20 min, then dropped and rose once during 80-min period. |
| 30-0280 | Cascade Creek near Petersburg | 57°01' | 132°47' | 120 | 1.86 | 04:00 | 30 | .02/.00 | |
| 30-0340 | Long River near Juneau | 58°10'00" | 133°41'50" | 183 | 1.44 | 03:20 | 45 | No seiche | Water level rose 0.07 ft. in 30 min, declined 0.65 ft. in next 340 min, then gradually rose to preearthquake level during 24 hr. |
| 30-0360 | Speel River near Juneau | 58°12'10" | 133°36'40" | 140 | .34 | 03:30 | 400 | .46 | Bubble gage; seiche lasted about 60 min. |
| 30-0400 | Dorothy Creek near Juneau | 58°13'40" | 134°02'25" | 350 | 1.79 | 03:40 | 19 | | At 04:30, water level began decline of 0.08 ft. during 70 min. |
| 30-0480 | Sheep Creek near Juneau | 58°16'30" | 134°18'50" | 629.8 | 1.55 | 03:50 | 4 | .04 | |
| 30-0600 | Perseverance Creek near Wacker | 55°24'40" | 131°40'05" | 600 | 1.65 | 03:30 | 10 | | A residual 0.02-ft drop in stage. |
| 30-0720 | Fish Creek near Ketchikan | 55°23'30" | 131°11'40" | 20 | .98 | 03:25 | 120 | .52/.16 | |
| 30-0760 | Manzanita Creek near Ketchikan | 55°36' | 130°59' | 140 | 2.10 | 04:00 | 200 | .35 | |
| 30-0780 | Grace Creek near Ketchikan | 55°39'28" | 130°58'14" | 15 | 2.01 | 03:40 | 100 | .07 | |
| 30-0865 | Neck Creek near Point Baker | 56°05'55" | 133°08'20" | 4 | 1.10 | 04:00 | 80 | .06/.03 | Tsunami crest at 09:20. |
| 30-0940 | Deer Lake Outlet near Port Alexander | 56°31'10" | 134°40'10" | 1 | 2.01 | 03:25 | 56 | .07 | Tsunami crest at 10:40. Stage dropped 0.05 ft after seiche was recorded, then recovered in 2½ hr; Tsunami crests superimposed on high tide at 09:25, 10:05, 10:55, and 22:35. |
| 30-0980 | Baranof River at Baranof | 57°05'15" | 134°50'30" | 140 | 3.05 | 04:00 | 170 | .025/.075 | Bubble gage. |
| 30-1000 | Takatz River near Baranof | 57°08'35" | 134°51'50" | 4 | 1.63 | 03:45 | 50 | .02 | Waves from lake or tsunami crests at 09:55 and 10:45. |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|-------------------------|---|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| Alaska—Continued | | | | | | | | | |
| 30-1020 | Hasselborg Creek near Angoon..... | 57°39'40" | 134°14'55" | 295 | 1.45 | ? | 80 | 0.15 | Float was frozen solidly in ice. Stage dropped 0.07 ft, rose gradually 1.88 ft in 70 min, then declined 0.48 ft in 3 hr. Earthquake dislodged batteries of manometer control unit and caused loss of record. |
| 30-1080 | Pavlov River near Tenakee..... | 57°50'30" | 135°02'10" | 15 | 4.18 | 03:50 | 30 | .72 | |
| 30-2115 | Tobay River near Chitina..... | 61°13'55" | 144°11'50" | 1,796.23 | ----- | 03:50 | ice | .03+ | |
| 30-2160 | Power Creek near Cordova..... | 60°35'15" | 145°37'05" | 33.5 | .70 | ----- | 50 | .27 | |
| 30-2370 | Nellie Juan River near Hunter..... | 60°28'20" | 148°43'30" | 90 | 4.97 | ----- | 28 | .02+ | Chart indicates only one up-and-down seiche motion. Water level then receded 0.40 ft in 6 hr, and gradually rose. Many aftershocks were recorded. |
| 30-2390 | Bradley River near Homer..... | 59°45'25" | 150°51'00" | 1,050 | .97 | 04:00 | 30 | .25/.33 | |
| 30-2435 | Snow River near Divide..... | 60°18'05" | 149°14'10" | 1,050 | 2.88 | 03:30 | 16 | No seiche | Water rose 1.02 ft in 20 min, then returned to normal over 24 hr. Three aftershocks were recorded. |
| 30-2480 | Trail River near Lawing..... | 60°26'00" | 149°22'20" | 460 | 2.8 | ----- | 63 | 1.02 | Float was frozen in before and after quake. Earthquake dammed creek upstream and thus shut off flow till March 29th. Float released from ice by quake. Irregular change of stage during 18 hr after quake. |
| 30-2610 | Cooper Creek at mouth near Cooper Landing..... | 60°28'30" | 149°52'30" | 450 | ----- | 03:20 | 6 | Tr. | |
| 30-2760 | Ship Creek near Anchorage..... | 61°13'25" | 149°38'00" | 530 | .23 | 03:00 | 11 | .95/.58 | |
| 30-2900 | Little Susitna River near Palmer..... | 61°42'40" | 149°13'40" | 920.6 | ----- | 03:30 | 19 | .17/.13 | Tsunami crests 330, 460, 500, 530, and 610 min after seiche was recorded. |
| 30-2957 | Terror River at mouth near Kodiak.. | 57°41'50" | 153°10'10" | 10 | 1.90 | 03:20 | 13 | .27 | Tsunami crests 330, 450, and 520 min after seiche was recorded. |
| 30-2960 | Uganik River near Kodiak..... | 57°41'05" | 153°25'10" | 20 | 4.17 | 03:25 | 75 | .00/.03 | 0.2 ft surge began shortly after quake was recorded; it continued through Mar h 28 and diminished through 29th. |
| 30-2963 | Spiridon Lake outlet near Larsen Bay. | 57°40'40" | 153°39'00" | 440 | .52 | 03:35 | 30 | 1.18/.02 | Tsunami crests 60, 120, and 170 min after seiche was recorded. |
| 30-2972 | Myrtle Creek near Kodiak..... | 57°36'15" | 152°24'10" | 50 | 1.15 | 04:10 | ----- | .25 | |
| Arizona | | | | | | | | | |
| 9-3834 | Little Colorado River at Greer..... | 34°01' | 109°27' | 8,500 | 1.97 | 03:30 | 1.6 | No seiche | Temporary 0.002 ft drop in stage. |
| 9-3880 | Little Colorado River near Hunt..... | 34°39' | 109°42' | 5,371.59 | 6.32 | 04:00 | .0 | No seiche | A residual 0.005-ft drop in stage. |
| 9-3935 | Silver Creek near Snowflake..... | 34°40'00" | 110°02'30" | 5,204.1 | 1.70 | 04:15 | 3.1 | .02 | Seiche lasted about 60 min near a fault. |
| 9-3975 | Chevelon Fork below Wildcat Canyon, near Winslow. | 34°38' | 110°43' | 5,905.16 | 2.66 | 03:30 | 3.3 | .1 | |
| 9-4210 | Lake Mead at Hoover Dam..... | 36°00'58" | 114°44'13" | MSL | 1,123.75 | 03:45 | 14,952,000 | .11 | Seiche lasted about 90 min near both a fault and a graben. |
| 9-4690 | San Carlos Reservoir at Coolidge Dam. | 33°10'30" | 110°31'45" | MSL | 2,412.22 | 03:50 | 53,460 | .35 | On extension of a fault. |
| 9-4897 | Big Bonita Creek near Fort Apache.. | 33°40'10" | 109°50'45" | 5,910 | 2.77 | 03:40 | 25 | .02 | A residual 0.005-ft drop in stage. |
| 9-4975 | Salt River near Chrysotile..... | 33°48' | 110°30' | 3,354.57 | 1.81 | 04:00 | 200 | Tr. | |
| 9-4985 | Salt River near Roosevelt..... | 33°37'10" | 110°55'15" | 2,177.14 | 7.80 | 03:40 | 260 | .02 | On a fault. |
| Arkansas | | | | | | | | | |
| 7-0475 | St. Francis River at Marked Tree.... | 35°31'58" | 90°25'25" | 196.44 | 6.60 | 03:50 | 2,080 | 0.26 | Near edge of Tertiary overlap. |
| 7-0480 | Auxiliary..... | 35°31' | 90°25' | ----- | 8.18 | 04:05 | 2,080 | .06 | |
| 7-0490 | West Fork White River at Greenland. | 35°59' | 94°10' | 1,233.00 | 1.14 | 03:50 | 34 | .08 | |
| 7-0500 | War Eagle Creek near Hindsville..... | 36°12'02" | 93°51'16" | 1,170.06 | ----- | ----- | ----- | .05 | |
| 7-0640 | Buffalo River near St. Joe..... | 35°59' | 92°45' | 560.35 | 5.55 | 03:40 | 1,250 | .12 | On edge of Tertiary overlap. |
| 7-0690 | Black River near Corning..... | 36°24'05" | 90°32'03" | 272.90 | 10.70 | 03:30 | 4,100 | .04 | |
| 7-0690 | Black River at Pocahtontas..... | 36°15' | 90°58' | 242.43 | 14.40 | 04:00 | 11,200 | .11 | Seiche may have lasted about 30 minutes near edge of Tertiary overlap. |
| 7-0695 | Spring River at Imboden..... | 36°12' | 91°10' | 254.07 | 5.08 | 04:00 | 1,500 | .04 | |
| 7-0745 | White River at Newport..... | 35°36'20" | 91°17'20" | 194.09 | 16.93 | 03:50 | 36,000 | .30 | Seiche lasted about 110 min. |
| 7-0759 | Greers Ferry Reservoir near Heber Springs. | 35°31'15" | 91°52'42" | ----- | 441.12 | 04:10 | 1,345 | .44 | On edge of Tertiary overlap. |
| 7-0768.5 | Cypress Bayou near Beebe..... | 35°01'30" | 91°52'23" | ----- | 10.90 | 04:10 | ----- | .04 | On Choctaw thrust fault. |
| 7-0770 | White River at De Valls Bluff..... | 34°47' | 91°27' | 152.93 | 22.40 | 03:50 | 88,000 | .16 | |
| 7-1950 | Osage River near Elm Springs..... | 36°13'15" | 94°17'20" | 1,052 | 1.58 | 04:10 | 36 | .02 | On extension of anormal fault. |
| 7-2470 | Poteau River at Cauthron..... | 34°55'08" | 94°17'53" | 569.53 | 5.00 | 03:30 | 40 | .02 | |
| 7-2494 | James Fork near Hackett..... | 35°09'45" | 94°24'25" | 459.71 | 3.02 | 03:50 | 64 | .16 | |
| 7-2495 | Cove Creek near Lee Creek..... | 35°43'20" | 94°24'30" | 852 | 1.57 | 03:50 | 8 | .07 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|--------------------------------|--|-----------|------------|--------------------|------------|--------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| Arkansas—Continued | | | | | | | | | |
| 7-2515 | Frog Bayou at Rudy..... | 35°31'25" | 94°16'30" | 475.08 | 2.96 | 04:00 | 106 | 0.15 | |
| 7-2540 | Six Mile Creek Subwatershed 5 near Chismville. | 35°13'45" | 93°54'50" | 475.83 | 13.44 | ----- | 1 | .03 | On possible extension of axis of anticline. |
| 7-2551 | Six Mile Creek subwatershed 23 near Branch. | 35°21'15" | 93°59'00" | 400.00 | 22.58 | ----- | 3.3/77 | .01 | On extension of a normal fault. Bubble gage. |
| 7-2555 | Hurricane Creek near Branch..... | 35°21' | 93°56' | 379.87 | 2.60 | 04:00 | 31 | .03 | On extension of a normal fault. |
| 7-2570 | Piney Creek near Dover..... | 35°33'00" | 93°09'25" | 487.66 | 3.56 | 04:10 | 580 | .48 | On axis of syncline. Seiche from long way round world at 05:05? |
| 7-2575 | Illinois Bayou near Scottsville..... | 35°27'58" | 92°02'28" | 447.54 | 6.40 | 03:50 | 485 | .06 | |
| 7-2615 | Fourche La Fave River near Gravelly. | 34°52' | 93°39' | 410.50 | 2.48 | 03:50 | 188 | .26 | Seiche lasted about 30 min. On possible extension of thrust fault. |
| 7-2640 | Bayou Meto near Lonoke..... | 34°44'10" | 91°54'58" | 199.11 | 10.80 | 04:00 | 370 | .03 | On possible extension of thrust fault. |
| 7-3370 | Red River at Index..... | 33°33'05" | 94°02'25" | 246.87 | 6.62 | 04:05 | 36,600 | .14 | On edge of Tertiary overlap. |
| 7-3395 | Rolling Fork near DeQueen..... | 34°03' | 94°25' | 318.24 | 4.50 | 04:00 | 340 | .04 | Near edge of Cretaceous overlap. |
| 7-3400 | Little River near Horatio..... | 33°55'10" | 94°23'15" | 272.89 | 9.37 | 03:50 | 3,000 | .00/.08 | Bubble gage. |
| 7-3405 | Cossatot River near DeQueen..... | 34°03' | 94°13' | 335.48 | 6.20 | 04:10 | 982 | .08 | Near edge of Cretaceous overlap. |
| 7-3410 | Saline River near Dierks..... | 34°06' | 94°05' | 353.09 | 5.90 | 04:05 | 95 | .07 | Do. |
| 7-3494.3 | Bodcau Creek at Stamps..... | 33°22'00" | 93°31'20" | ----- | 4.82 | 03:55 | 429 | .01 | On South Arkansas fault zone. |
| 7-3565 | Ouachita River South Fork at Mt. Ida. | 34°34' | 93°38' | 612.05 | 2.30 | 03:50 | 96 | .11 | A residual 0.02-ft. drop in stage. |
| 7-3575 | Lake Ouachita near Hot Springs..... | 34°34'20" | 93°11'50" | ----- | 573.10 | 03:20 | 1,970,000 | 1.45 | Seiche lasted about 140 min. Near both an anticline and a fault. |
| 7-3605 | Lake Greason near Murfreesboro..... | 34°08'55" | 93°42'55" | ----- | 537.10 | 04:00± | 208,000 | .45 | Seiche lasted about 60 min. On fault and near intrusive body. |
| 7-3615 | Antoine River at Antoine..... | 34°02'20" | 93°25'05" | 229.33 | 4.10 | 03:45 | 165 | .00/.02 | Near edge of Cretaceous overlap. Bubble gage? |
| 7-3621 | Smackover Creek near Smackover... | 33°20'40" | 92°46'45" | ----- | 5.80 | 03:50 | 235 | .18 | Near Arkansas fault zone. |
| 7-3625 | Moro Creek near Fordyce..... | 33°47' | 92°20' | 160.63 | 6.35 | 04:00 | 286 | .06 | |
| 7-3633 | Hurricane Creek near Sheridan..... | 34°19'10" | 92°20'40" | ----- | 9.70 | 03:40 | 300 | .04 | |
| 7-3635 | Saline River near Rye..... | 33°42' | 92°02' | 95 | 11.03 | 04:00 | 2,090 | .10 | |
| 7-3658 | Cornie Bayou near Three Forks..... | 33°02' | 92°56' | ----- | 5.08 | 03:55 | 72 | .02 | |
| 7-3659 | Three Creek near Three Creeks..... | 33°04' | 92°53' | ----- | 1.91 | 04:00 | 13 | .05 | |
| California | | | | | | | | | |
| 10-2904 | Lower Twin Lake near Bridgeport... | 38°09'20" | 119°20'20" | MSL | 7,208.58 | 03:50 | 4,000 | 0.06 | Seiche lasted about 240 min. On a normal fault. |
| 10-3385 | Donner Creek at Donner Lake near Truckee. | 39°19'25" | 120°14'00" | 5,930 | 1.70 | 03:10 | 23 | No seiche | Slight drop in stage. |
| 11-1445 | Salinas Reservoir near Pezo..... | 35°20'15" | 120°30'05" | MSL | 1,293.41 | 04:00 | 20,600 | .42 | Seiche lasted about 300 min. On a fault. |
| 11-1812 | Chabot Reservoir near San Leandro... | 37°43'17" | 122°07'15" | MSL | 227.30 | 03:50 | ----- | .30 | Seiche lasted 190 min. |
| 11-1814.9 | San Pablo Reservoir near Residence. | 37°56'31" | 122°15'40" | MSL | 305.88 | 03:45 | ----- | .06 | Seiche lasted about 140 min but was poorly recorded. On Hayward fault. |
| 11-1829.2 | Lafayette Reservoir near Briones Valley. | 37°53'05" | 122°15'40" | MSL | 445.64 | 04:00 | ----- | .00/.02 | Bubble gage? Seiche lasted about 240 min. On Hayward fault. |
| 11-1905 | Isabella Reservoir near Isabella..... | 36°38'50" | 118°28'50" | MSL | 2,557.45 | 03:20 | 187,700 | ----- | May be effect of wind. Duration about 230 min. Near Kern Canyon fault. |
| 11-2047 | Lake Success near Success..... | 36°03'40" | 118°55'18" | MSL | 598.42 | 04:00 | 13,400 | No seiche | Water level rose 0.02 ft in 10 min. Near edge of Sierra Nevada batholith. |
| 11-2109 | Lake Kaweah near Lemoncove..... | 36°24'53" | 119°00'07" | MSL | 571.06 | 04:00 | 8,450 | .06 | Seiche lasted about 50 min. On edge of Sierra Nevada batholith. |
| 11-2150 | North Fork Kings River near Cliff Camp. | 36°59'38" | 118°58'50" | 6,143.95 | 3.03 | 03:50 | 15 | .05 | Do. |
| 11-2210 | Pine Flat Reservoir near Piedra..... | 36°49'55" | 119°19'25" | MSL | 861.01 | 03:50 | 543,000 | .14 | Seiche seemingly lasted about 560 min. |
| 11-2501 | Millerton Lake at Friant..... | 37°00'00" | 119°42'10" | MSL | 518.07 | 03:50 | 274,300 | .03 | Seiche lasted about 100 min. Near edge of Sierra Nevada batholith. |
| 11-2713.5 | Merced River at Cressey..... | 37°25'28" | 120°39'47" | ----- | 10.34 | 03:45 | ----- | .01 | In Central Valley. |
| 11-2745.5 | San Joaquin River at Crows Landing Bridge. | 37°26'52" | 121°00'44" | ----- | 38.75 | 04:00 | ----- | .04 | Do. |
| 11-2875 | Don Pedro Reservoir near La Grange. | 37°42'48" | 120°24'14" | MSL | 575.40 | 03:30 | 200,700 | .02 | Record rather indistinct. |
| 11-2884 | Tuolumne River at La Grange Bridge. | 37°39'59" | 120°27'40" | ----- | 167.34 | 04:00 | ----- | .01 | |
| 11-2905 | San Joaquin River at Maze Road Bridge. | 37°38'28" | 121°13'37" | ----- | 14.56 | 03:50 | ----- | .02 | In Central Valley. |
| 11-2999.95 | Tulloch Reservoir near Knights Ferry. | 37°52'30" | 120°36'15" | MSL | 496.10 | 04:20 | 51,400 | .07 | Seiche may have lasted about 270 min. |
| 11-3087 | New Hogan Reservoir near Valley Springs. | 38°09'00" | 120°48'45" | MSL | 598.45 | 03:50 | 25,800 | .12 | Seiche lasted about 60 min. |
| 11-3166 | North Fork Mokelumne River above Tiger Creek. | 38°26'45" | 120°29'15" | ----- | 2.73 | 03:45 | ----- | .02 | Slight residual drop in stage. |
| 11-3200 | Pardee Reservoir near Spring Valley. | 38°15'30" | 120°51'00" | MSL | 551.83 | 04:00 | 176,400 | .38 | Seiche lasted about 180 min. |
| 11-3700 | Shasta Lake near Redding..... | 40°43'10" | 122°25'10" | MSL | 1,018.75 | 04:00 | 5,267,100 | .25 | Seiche lasted about 120 min. |
| 11-3879.95 | Black Butte Reservoir near Orland.. | 39°48'50" | 122°20'10" | MSL | 429.40 | 03:45 | 27,900 | .02 | Seiche lasted about 60 min. On a fault. |

Table 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|---|--|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| California—Continued | | | | | | | | | |
| 11-4180 | Yuba River at Englebright Dam..... | 39°14'22" | 121°16'00" | MSL | 627.76 | 03:30 | 1,580 | 0.05 | Storm or seiche recorded about 240 min. On edge of batholith. |
| 11-4270 | North Fork American River at North Fork Dam. | 38°56'15" | 121°01'25" | MSL | 715 | 03:30 | 599 | .02 | Seiche lasted about 60 min. |
| 11-4539 | Lake Berryessa near Winters..... | 38°30'50" | 122°06'15" | MSL | 437.76 | 03:50 | 1,559,300 | .18 | Seiche lasted about 190 min. |
| 11-4560 | Napa River near St. Helena..... | 38°29'40" | 122°25'50" | 200 | 1.05 | 03:45 | 20 | .01 | Temperature record unaffected by earthquake. |
| Colorado | | | | | | | | | |
| [About 40 gaging stations were out of operation owing to ice conditions during period of earthquake. All those that did record were in western half of State] | | | | | | | | | |
| 9-0664 | Red Sandstone Creek near Minturn. | 39°40'55" | 106°24'05" | 9,150 | 2.42 | 03:55 | 0.9 | 0.02 | Close to several faults. |
| 9-0802 | Fryingpan River at Ruedi..... | 39°21'40" | 106°49'10" | 7,500 | 2.15 | 04:00 | 30 | Tr. | On a fault. |
| 9-0850 | Roaring Fork at Glenwood Springs... | 39°32'50" | 107°19'50" | 5,720.73 | .92 | 03:45 | 260 | .02 | West of a thrust fault. |
| 9-0890 | West Divide Creek below Willow Creek, near Raven. | 39°16'32" | 107°31'10" | 7,820 | 1.90 | 04:10 | 2.4 | .04 | At southeast end of Piceance basin. |
| 9-1122 | East River below Cement Creek, near Crested Butte. | 38°47'25" | 106°52'20" | 8,450 | 3.76 | 04:20 | 42 | .03 | On a fault. |
| 9-1465 | East Fork Dailas Creek near Ridgeway. | 38°05'40" | 107°48'40" | 7,980 | 1.95 | 04:00 | 5.0 | .01 | On west edge of San Juan volcanic area. |
| 9-1712 | San Miguel River near Telluride..... | 37°56'55" | 107°52'35" | 8,622.81 | ----- | 04:00 | 16 | .01 | On a fault. |
| 9-2410 | Elk River at Clark..... | 40°43'03" | 106°54'55" | 7,267.75 | .75 | 04:00 | 32 | Tr. | On west edge of Sierra Madre uplift. A 0.001-ft. rise in stage. |
| 9-3028 | White River near Buford..... | 40°02' | 107°31' | ----- | 2.75 | 03:50 | 121 | .03 | On White River uplift. |
| 9-3042 | White River above Coal Creek, near Meeker. | 40°00'20" | 107°49'30" | 6,400 | 1.56 | 03:45 | 260 | .30 | Do. |
| 9-3443 | Navaho River near Chromo..... | 37°01'55" | 106°43'56" | 7,700 | 3.41 | 04:00 | 26 | .01 | Near dikes and faults. |
| 9-3610 | Hermosa Creek near Hermosa..... | 37°25'30" | 107°50'20" | 6,705.88 | .51 | 04:00 | 14 | Tr. | |
| 9-3612 | Falls Creek near Durango..... | 37°22'00" | 107°52'00" | 7,120 | 3.05 | 04:00 | 1 | .02 | |
| 9-3614 | Junction Creek near Durango..... | 37°20'05" | 107°54'30" | 7,045.65 | 2.44 | 03:50 | 3.0 | Tr. | |
| Connecticut | | | | | | | | | |
| No seismic seiche was recorded at any gaging station. | | | | | | | | | |
| Delaware | | | | | | | | | |
| No report received. | | | | | | | | | |
| Florida | | | | | | | | | |
| 2-2310 | St. Marys River near Macclenny.... | 30°21'35" | 82°04'55" | 40.00 | 5.20 | 04:15 | 490 | 0.66 | Seiche lasted about 40 min. |
| 2-2313.5 | St. Johns headwaters near Vero Beach. | 27°38'35" | 80°40'28" | 18.56 | 6.05 | 04:50 | ----- | .02 | |
| 2-2321 | Lake Washington near Eau Gallie... | 28°08'50" | 80°44'10" | 10.39 | 4.17 | 03:55 | 4,298 | .04 | |
| 2-2324 | St. Johns River near Cocoa..... | 28°22'10" | 80°52'22" | MSL | 12.50 | 04:20 | 870 | .10 | |
| 2-2332 | Little Econlockhatchee River near Union Park. | 28°31'29" | 81°14'39" | 56.19 | 6.60 | 04:20 | 20 | .02 | |
| 2-2360 | St. Johns River at St. Francis Landing, near Deland. | 29°02'14" | 81°25'05" | -1.11 | 1.66 | 04:10 | 3,700 | .02 | |
| 2-2369 | Palatka Creek at Cherry Lake outlet, near Groveland. | 28°36' | 81°49' | MSL | 95.64 | 04:35 | 20 | .01 | |
| 2-2445 | Auxiliary..... | 28°36' | 81°49' | MSL | 94.56 | 04:10 | 20 | .05 | |
| 2-2445 | Little Haw Creek near Seville..... | 29°19' | 81°23' | 5.74 | 3.83 | 04:10 | 80 | No seiche | Stage declined 0.34 ft in 20 min, then began to rise. |
| 2-2465 | St. Johns River at Jacksonville..... | 30°19'13" | 81°39'32" | -10.00 | ? | 04:05 | ----- | .06 | |
| 2-2465 | St. Johns River at Naval Air Station, near Jacksonville. | 30°13'39" | 81°39'58" | -10.00 | 10.78 | 04:30 | ----- | .03 | |
| 2-2469 | Moultrie Creek near St. Augustine (State Hwy. 207). | 29°50'50" | 81°21'39" | 14.24 | 4.11 | 04:00 | 19 | No seiche | A 0.01-ft drop in stage. |
| 2-2500 | Turkey Creek near Palm Bay..... | 28°00'46" | 80°36'28" | -1.03 | 2.36 | 03:45 | 34 | .05 | |
| 2-2520 | Fellsmere Canal near Fellsmere..... | 27°49'18" | 80°36'27" | 7.90 | 1.50 | 04:10 | 34 | .01 | |
| 2-2540 | North Fork St. Lucie River at White City. | 27°22'26" | 80°20'33" | MSL | ----- | 04:15 | ----- | .13 | Seiche lasted about 20 min. |
| 2-2560 | Fisheating Creek near Venus..... | 27°03'57" | 81°25'52" | 46.52 | 9.92 | 04:35 | 2 | .04 | |
| 2-2638 | Shingle Creek at airport, near Kissimmee. | 28°18'14" | 81°27'04" | 60.66 | 5.02 | 04:05 | 45 | .04 | Seiche lasted about 15 min. |
| 2-2674 | Lake Hatchineha near Lake Wales... | 28°00'00" | 81°22'50" | 47.23 | 4.90 | 04:40 | 6,636 | Tr. | |
| 2-2691 | Kissimmee River at Fort Kissimmee. | 27°35'27" | 81°09'20" | 37.98 | 7.03 | 04:40 | ----- | .04 | |
| 2-2715 | Josephine Creek near DeSoto City... | 27°22'26" | 81°23'37" | 52.99 | 3.75 | 04:20 | 21 | Tr. | |
| 2-2720 | Istokpoga Canal near Cornwall..... | 27°22'56" | 81°09'45" | 27.91 | 35.00 | 04:10 | 10 | Tr. | |
| 2-2720 | Auxiliary..... | 27°23'16" | 81°10'50" | ----- | 5.46 | 04:10 | 10 | Tr. | |
| 2-2784.5 | West Palm Beach Canal near Loxahatchee. | 26°41'05" | 80°22'15" | MSL | ----- | 04:35 | 135 | ----- | 0.14/0.06 units on deflection meter. |
| 2-2975 | Auxiliary..... | 26°41'05" | 80°22'00" | MSL | 12.75 | 03:50 | 135 | .22 | Seiche lasted about 60 min. |
| 2-2980 | Joshua Creek at Nocatee..... | 27°09'59" | 81°52'47" | 3.94 | 4.32 | 04:20 | 13 | .07 | Seiche lasted about 20 min. |
| 2-2980 | Horse Creek near Arcadia..... | 27°11'57" | 81°59'19" | 10.96 | 3.03 | 04:10 | 76 | .04 | |
| 2-2982 | Myakka River at Myakka City..... | 27°20'47" | 82°09'27" | 23.81 | 5.91 | 04:20 | 433 | Tr. | |
| 2-2990 | Myakka River near Sarasota..... | 27°14'25" | 82°18'50" | 7.92 | 4.31 | 04:15 | 74 | .02 | |
| 2-3014 | Turkey Creek near Durant..... | 27°56'15" | 82°11'39" | 43.00 | 2.52 | 03:50 | ----- | .03 | |
| 2-3034 | Cypress Creek near San Antonio..... | 28°19'25" | 82°23'03" | MSL | 73.18 | 04:25 | 29 | Tr. | |
| 2-3038 | Cypress Creek near Sulphur Springs. | 28°05'20" | 82°24'33" | MSL | 28.52 | 04:15 | ----- | .02 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|-------------------------|---|-----------|-----------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| Florida—Continued | | | | | | | | | |
| 2-3045 | Hillsborough River at 22d Street, near Tampa. | 28°01'15" | 82°26'08" | MSL | 0.50 | 04:15 | 472 | 0.15 | Seiche superimposed on tidal curve. |
| 2-3065 | Sweetwater Creek near Sulphur Springs. | 28°02'33" | 82°30'44" | 30.68 | .50 | 04:20 | 41 | Tr. | |
| 2-3103 | Pithlachascotee River nr. New Port Richey. | 28°18'19" | 82°39'37" | 7.06 | 4.28 | 04:40 | 33 | Tr. | |
| 2-3105.5 | Weekiwachee River near Bayport. | 28°31'58" | 82°37'38" | -10.00 | 10.34 | 03:45 | ----- | .01 | 0.38 units on deflection meter. |
| 2-3106.5 | Chassahowitzka River near Homosassa. | 28°42'54" | 82°34'38" | -10.00 | 11.64 | 04:00 | ----- | .03 | |
| 2-3107 | Homosassa River at Homosassa. | 28°47'08" | 82°37'08" | -10.00 | ? | 04:15 | ----- | Tr. | Possibly 0.2 units on deflection meter. |
| 2-3107.5 | Crystal River near Crystal River. | 28°54'17" | 82°38'13" | -10.00 | ----- | 03:50 | ----- | .06 | Seiche superimposed on tidal curve. |
| 2-3142 | Tenmile Creek at Lebanon Station. | 29°09'39" | 82°38'21" | 15.00 | 6.35 | 04:40 | 63 | .02 | |
| 2-3155 | Suwannee River at White Springs. | 30°19'32" | 82°44'18" | 48.54 | ----- | 03:50 | 4,450 | Tr. | |
| 2-3155.5 | Suwannee River at Suwannee Springs. | 30°23'34" | 82°56'00" | MSL | 55.25 | 03:40 | ----- | .13 | |
| 2-3195 | Suwannee River at Ellaville. | 30°23'04" | 83°10'19" | 27.22 | 18.50 | 03:45 | 17,700 | .06 | |
| 2-3235 | Suwannee River near Wilcox. | 29°36' | 82°56' | MSL | 12.10 | 03:45 | 26,800 | .24 | |
| 2-3290 | Chipola River near Altha. | 30°22'02" | 85°09'58" | 19.95 | 17.25 | 03:50 | 4,120 | .30 | |
| 2-3280 | Yellow River at Milligan. | 30°45'10" | 86°37'45" | 45.00 | 6.34 | 03:40 | 1,860 | .01 | |
| 2-3765 | Perdido River at Barrineau Park. | 30°41'25" | 87°26'25" | 25.77 | 3.44 | 03:10 | 855 | .20 | |
| 2-2785 | West Palm Beach Canal near Loxahatchee (S-5A). | 26°41'00" | 80°22'10" | MSL | 12.70 | 04:35 | 132 | .03 | |
| 2-2785.5 | Levee 8 Canal at West Palm Beach Canal, near Loxahatchee. | 26°41'05" | 80°21'35" | MSL | 7.30 | 04:35 | 112 | .32 | On head water; brief decline of 0.01 ft on tall water. No trace on deflection meter. |
| 2-2790 | West Palm Beach Canal at West Palm Beach. | 26°38'40" | 80°03'32" | MSL | 8.23 | 04:20 | 182 | .06 | 0.02 units on deflection meter. |
| 2-2805 | Hillsboro Canal below HGS-4, near South Bay. | 26°42'00" | 80°42'45" | MSL | -.52 | 04:20 | 238 | .30 | A 0.08-ft drop in stage. |
| 2-2813 | Hillsboro Canal near Deerfield Beach. | 26°21'20" | 80°17'58" | MSL | 15.83 | 04:00 | 46 | .01 | No trace on deflection meter. |
| 2-2815 | do. | 26°19'39" | 80°07'51" | MSL | 1.22 | 04:30 | 67 | .13 | Seiche superimposed on tidal curve; no trace on deflection meter. |
| 2-2817 | Pompano Canal at S-33, near Pompano Beach. | 26°13'45" | 80°17'50" | MSL | 6.50 | 04:00 | 3 | .20 | |
| 2-2820 | Pompano Canal at Pompano Beach. | 26°13'51" | 80°07'28" | MSL | 3.74 | 04:10 | ----- | .04 | No trace on deflection meter. |
| 2-2821 | Cypress Creek at S-37A, near Pompano Beach. | 26°12'20" | 80°07'57" | MSL | 3.82 | 04:00 | ----- | .03 | 0.44 units on deflection meter. |
| 2-2832 | Plantation Road Canal at S-33, near Fort Lauderdale. | 26°08'05" | 80°11'42" | MSL | 5.96 | 03:55 | ----- | .04 | |
| 2-2850 | North New River Canal near Fort Lauderdale (auxiliary). | 26°05'39" | 80°13'50" | MSL | ? | 04:40 | 39 | ? | Seiche superimposed on tidal curve. |
| 2-2854 | South New River Canal (east of S-9) near Davie. | 26°03'40" | 80°28'30" | MSL | ? | 04:10 | 0 | ----- | 0.02 ft on lower stage; 0.05 ft on upper stage. 0.04 units on deflection meter. |
| 2-2861 | South New River Canal at S-13 near Davie. | 26°03'57" | 80°12'32" | MSL | ? | 04:15 | ----- | ----- | No trace on upper stage; trace on lower stage. 0.09 units on deflection meter. |
| 2-2861.8 | Snake Creek Canal at S-30 near Hialeah. | 25°57'22" | 80°25'54" | MSL | 5.53 | 04:00 | ----- | .06 | 0.48 on deflection meter with a slight decrease in flow. |
| 2-2862 | Snake Creek Canal at NW 67th Ave., near Hialeah. | 25°37'50" | 80°18'40" | MSL | 2.52 | 04:00 | ----- | .00 | 0.18 deflection units on deflection meter. |
| 2-2863 | Snake Creek Canal at S-29 at North Miami Beach. | 25°33'41" | 80°09'22" | MSL | 2.52 | 03:55 | 26 | .11 | Seiche lasted about 60 min; 0.29/0.36 units on deflection meter followed by slight decrease in flow. |
| 2-2863.4 | Biscayne Canal at S-28 near Miami. | 25°52'24" | 80°10'58" | MSL | 2.00 | 04:15 | 36 | .01 | 0.41 units on deflection meter of which 0.19 was lasting decrease in flow. |
| 2-2863.5 | Little River Canal at Palm Avenue, in Hialeah. | 25°52'13" | 80°17'00" | MSL | 2.05 | 04:35 | ----- | .01 | |
| 2-2863.8 | Little River Canal at S-27, in Miami. | 25°51'11" | 80°11'38" | MSL | ----- | 04:40 | ----- | Tr. | Seiche lasted about 60 min; 0.40 units on deflection meter with small permanent decrease in flow. |
| 2-2864 | Miami Canal at HGS-3 and S-3, in Lake Harbor. | 26°41'55" | 80°48'25" | MSL | 13.45 | 04:15 | ----- | .15 | Quake affected the lakeside gage but not the landside gage; 0.56/0.12 units on deflection meter with apparent lasting increase of 0.02 units. |
| 2-2864 | Miami Canal south of S-3 at Lake Harbor. | 26°41'55" | 80°48'25" | MSL | ----- | 04:00 | ----- | ----- | 0.38/0.40 units on deflection meter with no lasting change in flow. |
| 2-2874 | Miami Canal at broken dam near Miami. | 25°56'00" | 80°25'50" | MSL | ----- | ----- | ----- | ----- | Trace of quake on both stage and deflection records. |
| 2-2875 | Miami Canal at Pennsuco near Miami. | 25°53'40" | 80°22'45" | MSL | 2.95 | ? | ----- | .05 | Seiche lasted about 150 min. |
| 2-2882 | Miami Canal at Palmetto By-pass, near Hialeah. | 25°51'11" | 80°19'22" | MSL | 2.55 | 04:05 | ----- | .07 | 0.02 units on deflection meter. |
| 2-2886 | Miami Canal at NW 36th St., Miami. | 25°48'29" | 80°15'44" | MSL | 2.42 | 04:15 | ----- | .04 | 0.33/0.29 units on deflection meter; seiche lasted about 40 min. |
| 2-2888 | Tamiami Canal outlets, Monroe to Carnestown (at bridge 84). | 25°53'10" | 81°15'30" | MSL | 1.33 | 03:30 | ----- | .05 | |
| | Tamiami Canal at bridge 77 near Carnestown (auxiliary). | 25°54' | 81°21' | 3.14 | 4.00 | 03:30 | ----- | .05 | Seiche superimposed on tidal (?) curve. |
| 2-2889 | Tamiami Canal at 40-mile bend, near Miami (auxiliary). | 25°45'50" | 80°49'50" | MSL | 7.28 | 03:45 | 10 | .06 | |
| 2-2890 | Tamiami Canal at bridge 45, near Miami. | 25°45'40" | 80°37'40" | ----- | 6.22 | 04:45 | ----- | .10 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|-------------------------|--|-----------|-----------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| Florida—Continued | | | | | | | | | |
| 2-2890.4 | Tamiami Canal below S-12-C, near Miami (auxiliary). | 25°45'40" | 80°43'34" | 0.04 | 6.88 | 05:00 | ----- | 0.03 | |
| | Tamiami Canal below S-12-B, near Miami (auxiliary). | 25°45'40" | 80°46'05" | .04 | 6.98 | 03:35 | ----- | .05 | |
| | Tamiami Canal above S-12-B, near Miami (auxiliary). | 25°45'42" | 80°46'05" | .05 | 7.15 | 04:15 | ----- | .04 | |
| | Tamiami Canal above S-12-C, near Miami (auxiliary). | 25°45'42" | 80°43'34" | ----- | 7.15 | 03:40 | ----- | .04 | |
| 2-2895 | Tamiami Canal near Coral Gables... | 25°45'43" | 80°19'42" | MSL | 2.50 | 05:00 | 50 | .04 | No trace on deflection meter. |
| 2-2905.1 | Miami Canal at NW 27th Ave., Miami. | 25°47'32" | 80°14'24" | MSL | 1.20 | 04:10 | ----- | .37 | Seiche superimposed on tidal curve. |
| 2-2905.2 | South Fork Miami River at NW 29th Ave., Miami. | 25°47'00" | 80°14'32" | MSL | ----- | 04:10 | ----- | .03 | |
| 2-2905.3 | Miami River at Brickell Ave., Miami. | 25°45'11" | 80°11'25" | MSL | 0.87 | 03:55 | ----- | .17 | Seiche superimposed on tidal curve. 1.09 units on deflection meter with no lasting change in flow. |
| 2-2905.6 | Coral Gables Canal at Red Road, in Coral Gables. | 25°44'17" | 80°17'13" | MSL | 2.53 | 04:30 | ----- | .02 | |
| 2-2905.8 | Coral Gables Canal near South Miami. | 25°42'20" | 80°15'40" | MSL | .25± | 04:25 | ----- | .15 | Seiche superimposed on tidal curve. 0.70 units on deflection meter. |
| 2-2906 | Snapper Creek Canal near Coral Gables. | 25°45'40" | 80°23'05" | MSL | 3.05 | 04:10 | ----- | .02 | Pen lines of stage and deflection were both slightly displaced downward; 0.1 units on deflection meter. |
| | Snapper Creek Canal at Miller Drive, near South Miami (auxiliary). | 25°42'56" | 80°22'59" | MSL | 3.00 | 04:10 | ----- | .09 | Seiche lasted about 40 min. |
| 2-2907 | Snapper Creek Canal at S-22, near South Miami. | 25°40'11" | 80°17'03" | MSL | 2.94 | 03:45 | ----- | .03 | 0.07 units on deflection meter; seiche lasted about 30 min. |
| 2-2907.15 | Goulds Canal near Goulds..... | 25°32'15" | 80°19'55" | MSL | ----- | 03:25 | ----- | .03 | |
| 2-2907.2 | Military Canal near Homestead..... | 25°29'20" | 80°20'55" | MSL | .79 | 04:05 | ----- | .05 | Seiche lasted about 20 min. |
| 2-2907.45 | Model Land Canal at control, near Florida City (auxiliary). | 25°21'59" | 80°25'53" | MSL | ----- | 04:20 | ----- | .05 | Seiche lasted about 20 min. |
| 2-2908.5 | Shark River near Homestead..... | 25°23'10" | 81°01'00" | ----- | ----- | 04:00 | ----- | .30 | Seiche superimposed on tidal curve; 0.75 units on deflection meter. |
| 2-2934.8 | Lake Otis at Winter Haven..... | 28°01'10" | 81°42'35" | 120.00 | 6.15 | 03:55 | 144 | Tr. | |
| 2-2949 | Saddle Creek at structure P-11, near Bartow. | 27°56'17" | 81°51'05" | 94.08 | 1.02 | 03:55 | 2 | .01 | |
| 2-2962 | Little Charlie Bowlegs Creek near Sebring (auxiliary). | 27°48'40" | 81°33'25" | 62.32 | 16.52 | 04:40 | 3 | .02 | |
| 2-2965 | Charlie Creek near Gardner..... | 27°22'29" | 81°47'48" | 21.66 | 3.21 | ? | 322 | Tr. | |
| Georgia | | | | | | | | | |
| 2-1872.5 | Hartwell Reservoir near Hartwell.... | 34°21'25" | 82°49'20" | ----- | 664.39 | 03:50 | ----- | 0.05 | |
| 2-1975.5 | Little Brier Creek near Thomson.... | 33°20'24" | 82°27'29" | 313.95 | 6.47 | 04:20 | 100 | .04 | On edge of Cretaceous overlap. |
| 2-1980 | Brier Creek at Millhaven..... | 32°56'00" | 81°39'05" | 95.88 | 6.94 | 04:20 | 1,380 | .05 | |
| 2-2030 | Canoochee River near Claxton..... | 32°11'05" | 81°53'25" | 80.5 | 8.00 | 03:55 | 1,190 | .09 | On Ochlockonee Fault of Sever (1966). |
| 2-2130.5 | Walnut Creek near Gray..... | 32°58'20" | 83°37'10" | 390 | 2.10 | 04:00 | 60 | .03 | |
| 2-2210 | Murder Creek near Monticello..... | 33°25' | 83°40' | 498.21 | 1.32 | 04:00 | 60 | .02 | On Towaliga fault. |
| 2-2255 | Ohoopce River near Reidsville..... | 32°04' | 82°11' | 73.8 | 10.75 | 04:30 | 2,750 | .09 | On possible extension of Ochlockonee fault of Sever (1966). |
| 2-2261 | Penholoway Creek near Jesup..... | 31°34'00" | 81°50'18" | ----- | 6.74 | 04:05 | 118 | .03 | On fault of Callahan (1964, fig. 5). |
| 2-2265 | Satilla River near Waycross..... | 31°14' | 82°19' | 66.43 | 11.78 | 04:40 | 2,000 | .06 | Do. |
| 2-3145 | Suwannee River at Fargo..... | 30°41' | 82°34' | 91.90 | 10.76 | ----- | 2,200 | .07 | |
| | Auxiliary | 30° | ----- | ----- | ----- | ----- | ----- | .03 | |
| 2-3160 | Alapaha River near Alapaha..... | 31°23' | 83°10' | 209.34 | 9.80 | 03:35 | 1,480 | .09 | On possible extension of Ochlockonee fault of Sever (1966). |
| 2-3175 | Alapaha River at Statenville..... | 30°42' | 83°01' | 76.77 | 12.19 | 04:40 | 2,650 | .22 | On fault of Callahan (1964, fig. 5). |
| 2-3275 | Ochlockonee River near Thomasville. | 30°52' | 84°03' | 133.6 | 14.10 | 04:30 | 3,300 | .05 | On Ochlockonee fault of Sever (1966). |
| 2-3316 | Chattahoochee River near Cornelia... | 34°33' | 83°37' | 1,128.53 | 3.24 | 04:10 | 3,000 | .03 | |
| 2-3350 | Chattahoochee River near Norcross... | 34°00' | 84°12' | 878.14 | ----- | ----- | ----- | .18 | On Brevard fault zone. |
| 2-3390 | Yellowjacket Creek near La Grange... | 33°05'25" | 85°03'45" | 601 | 4.35 | 03:40 | 245 | .12 | |
| 2-3432 | Pataula Creek near Lumpkin..... | 31°56' | 84°48' | 224.34 | 2.44 | 04:10 | 120 | .15 | On edge of Tertiary overlap. |
| 2-3465 | Potato Creek near Thomaston..... | 32°54'15" | 84°21'45" | 600 | 4.35 | 05:00 | 880 | .03 | On SE flank of Wacochee anticlinal belt. |
| 2-3490 | Whitewater Creek below Rambulette Creek, nr. Butler. | 32°28' | 84°16' | 365.85 | 1.86 | 04:10 | 180 | .015 | Near edge of Tertiary overlap. |
| 2-3499 | Turkey Creek at Byromville..... | 32°12' | 83°54' | ----- | 8.34 | 04:20 | 130 | .05 | Near Andersonville fault. |
| 2-3506 | Kinchafoonee Creek at Preston..... | 32°03' | 84°33' | 337.7 | 4.86 | 04:10 | 375 | .06 | |
| 2-3534 | Pachita Creek near Edison..... | 31°33' | 84°41' | 212.64 | 5.34 | 04:30 | 440 | .11 | |
| 2-3560 | Flint River at Bainbridge..... | 30°55' | 84°34' | 58.06 | 20.80 | 04:00 | 15,000 | .13 | |
| 2-3570 | Spring Creek near Iron City..... | 31°03' | 84°43' | 85.7 | 9.60 | 04:20 | 1,100 | .09 | |
| 2-3800 | Elijah River at Elijah..... | 34°42' | 84°29' | 1242.32 | 5.63 | 04:40 | 800 | .06 | On Murphy syncline. |
| 2-3870 | Conasauga River at Tilton..... | 34°40' | 84°56' | 622.28 | 21.30 | 04:00 | 12,000 | .10 | On Rome fault. |
| 2-3885 | Oostanaula River near Rome..... | 34°18' | 85°08' | 561.70 | 32.10 | 04:05 | 28,000 | .09 | Do. |
| 2-3970 | Coosa River near Rome..... | 34°12' | 85°16' | 563.05 | 31.10 | 04:00 | 43,000 | .12 | In Coosa syncline extended and near Rome fault. |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|---|--|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| Hawaii | | | | | | | | | |
| [No effects of the Alaska earthquake were found on records of stations on the islands of Oahu, Maui, and Molokai in the Hawaiian group nor of stations on Okinawa and on the islands of Guam and Tutuila, American Samoa] | | | | | | | | | |
| 40-0310 | Waimea River near Waimea, Kauai | 21°59'02" | 159°39'46" | 25 | 4.59 | 03:50 | 169 | Tr. | |
| 40-0610 | North Waimea ditch near Lihue, Kauai | 22°03'55" | 159°28'12" | 1,105.45 | 7.23 | 04:00 | 24 | 0.03 | |
| 40-1000 | Hanalei tunnel outlet near Lihue, Kauai | 22°04'57" | 159°27'52" | 1,201 | 1.00 | 03:45 | 45 | Tr. | |
| 40-7040 | Waikuku River above Hila School ditch, near Hilo, Hawaii | 19°42'55" | 155°09'10" | 1,060 | 4.58 | 03:45 | 302 | .17 | |
| 40-7580 | Waikoloa Stream at Marine Dam, near Kamuela, Hawaii | 20°02'48" | 155°39'58" | 3,450 | 1.60 | 03:45 | 7.4 | .01 | |
| Idaho | | | | | | | | | |
| 13-0320 | Bear Creek above reservoir near Irwin | 43°16'45" | 111°13'15" | 5,640 | ----- | ----- | 18 | 0.01 | |
| 13-0505 | Henrys Fork at St. Anthony | 43°58' | 111°40'20" | 4,950.7 | ----- | ----- | 1,110 | .02 | |
| 13-0522 | Teton River near Driggs | 43°47' | 111°13' | 5,952.9 | ----- | ----- | 236 | .03 | |
| ----- | Disposal Pond at National Reactor Testing Station | 43° | 112° | ----- | ----- | 03:40 | 4,919.10 | .56 | Seiche lasted about .140 min. |
| 3-2015 | Lucky Peak Reservoir near Boise | 43°32' | 116°04' | MSL | 2,991.30 | ----- | 146,100 | .24 | Seiche lasted more than an hour. On a normal fault. |
| Illinois | | | | | | | | | |
| 3-3815 | Little Wabash River at Carmi | 38°03'40" | 88°09'35" | 339.91 | 26.74 | 04:00 | 8,700 | Tr. | On a fault trending north-northeast. |
| ----- | Auxiliary | 38°05'30" | 88°09'20" | 339.91 | 26.23 | 04:00 | 8,700 | 0.10 | Do. |
| 3-3825 | Saline River near Junction | 37°41'52" | 88°16'00" | 320.40 | 37.07 | ----- | 1,200 | .02 | On extension of a fault trending north-northeast. |
| ----- | Auxiliary | 37°39'15" | 88°15'10" | 320.42 | 36.25 | 03:50 | 1,200 | .02 | Do. |
| 4-0925 | Wolf Lake at Chicago | 41°39'53" | 87°32'22" | 580.45 | 1.25 | 04:00 | ----- | .04 | |
| 4- | West Branch Du Page River | 41°43'20" | 88°07'45" | ----- | 2.00 | ----- | ----- | .04 | |
| 4- | East Branch Du Page River | 41°44'10" | 88°07'59" | ----- | 2.14 | ----- | ----- | .03 | |
| 5- | Money Creek at Lake Bloomington | 40°39'47" | 88°56'23" | 700.00 | 8.32 | 04:00 | ----- | .052 | |
| Indiana | | | | | | | | | |
| 3-3285 | Eel River near Logansport | 40°46'55" | 86°15'50" | 621.50 | 5.80 | 04:00 | 2,000 | Tr. | Bubble gage. |
| 3-3301.4 | Smalley Lake near Washington Center | 41°18'52" | 85°35'03" | ----- | 2.77 | 04:15 | 63 | 0.03 | A residual 0.01-ft rise in stage. On south side of Michigan basin. |
| 3-3355 | Wabash River at Lafayette | 40°25'19" | 86°53'49" | 504.14 | 9.40 | 04:20 | 13,000 | .07 | |
| 3-3405 | Wabash River at Montezuma | 39°47'33" | 87°22'26" | 457.75 | 10.70 | 04:00 | 15,000 | .24 | |
| 3-3485 | White River near Noblesville | 40°07' | 85°38' | 763.08 | 5.38 | 04:40 | 760 | .02 | |
| 3-3488 | White River at Clare | 40°06' | 85°38' | ----- | 15.40 | 03:50 | ----- | .08 | |
| 3-3510 | White River at Broad Ripple near Nora (auxiliary) | 39°52'18" | 86°08'30" | 710.94 | 3.55 | 03:50 | 1,300 | .39 | |
| 3-3530 | White River at Indianapolis | 39°45'05" | 86°10'30" | 662.26 | 4.72 | 04:00 | 1,720 | .04 | A residual 0.02-ft drop in stage. |
| 3-3532 | Eagle Creek at Zionsville | 39°56'56" | 86°15'22" | 816.85 | 3.34 | 03:50 | 146 | Tr. | A residual 0.01-ft drop in stage. |
| 3-3630 | Driftwood River near Edinburg | 39°20'21" | 85°59'11" | 636.99 | 4.35 | 03:25 | 1,200 | No seiche | A 0.05-ft drop in stage. |
| 3-3715 | East Fork White River near Bedford (auxiliary) | 38°49'33" | 86°30'48" | 473.59 | ----- | 03:50 | 5,100 | .06 | |
| 3-3752 | Beaver Creek Reservoir near Jasper | 38°24'10" | 86°50'30" | ----- | 27.82 | 04:00 | ----- | 0.07 | On east side of Illinois basin. |
| 4-0930 | Deep River at Lake George outlet at Hobart | 41°32'05" | 87°15'30" | 588.17 | 2.32(?) | 04:10 | 100 | .03 | |
| 4-0976.8 | Jimerson Lake at Nevada Mills | 41°43'31" | 85°04'55" | 964.44 | 4.45 | 04:15 | 283 | .05 | On south side of Michigan basin. |
| 4-0995 | Pigeon Creek at Hogback Lake outlet near Angola | 41°37'24" | 85°05'44" | 940.00 | 8.93 | 03:50 | 35 | .05 | Do. |
| 4-1004.5 | Syracuse Lake at Syracuse | 41°25'23" | 85°44'41" | 858.57 | 8.20 | 04:00 | 414 | .02 | Do. |
| Iowa | | | | | | | | | |
| 5-4870 | Lake Ahquabi near Indianola | 41°17'35" | 93°35'40" | ----- | 5.42 | 03:45 | ----- | 0.02 | |
| 5-4590 | Shell Rock River at Northwood | 43°24'50" | 93°13'10" | ----- | ----- | ----- | 225 | No seiche | A lasting 0.02-ft. drop in stage. On southwest flank of syncline. |
| Kansas | | | | | | | | | |
| 6-8535 | Republican River near Hardy | 40°00' | 97°56' | 1,501.46 | 3.80 | 04:05 | 157 | 0.00/0.07 | On northeast flank of Salina basin. Bubble gage. |
| 6-8665 | Smoky Hill River at Mentor | 38°47'54" | 97°34'28" | 1,211.40 | 6.20 | 03:50 | 66 | .00/0.04 | On Abilene arch. Bubble gage. |
| 6-8870 | Big Blue River near Manhattan (auxiliary) | 39°14'14" | 96°34'16" | 991.86 | 3.80 | 03:55 | 400 | .00/0.17 | On Nemaha uplift. Bubble gage. |
| 6-9110 | Marsias des Cygnes at Melvern | 38°31'50" | 95°46'40" | 939.11 | 5.60 | 04:00 | .2 | .00/0.07 | Bubble gage. A residual 0.02-ft. drop in stage. |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|--------------------------------|--|-----------|-----------|--------------------|------------|----------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| Kansas—Continued | | | | | | | | | |
| 7-1423 | Rattlesnake Creek near Macksville... | 37°52'20" | 98°52'30" | 1,963.46 | 3.85 | 04:00 | 26 | 0.00/.03 | South-southeast of Central Kansas uplift. Bubble gage. On trough on east side of Nemaha uplift. Bubble gage. |
| 7-1478 | Walnut River at Winfield..... | 37°14' | 97°00' | 1,082.86 | 2.67 | 03:55 | 40 | .00/.05 | |
| 7-1659 | Toronto Reservoir near Toronto..... | 37°44'30" | 95°56'00" | 897.46 | ----- | 04:10 | 13,000 | .05 | On crest of Precambrian rise. A residual 0.002-ft drop in stage. |
| 7-1675 | Otter Creek near Climax..... | 37°42'30" | 96°13'30" | 977.76 | 2.99 | 04:10 | 0 | .02 | |
| 7-1680 | Fall River Reservoir near Fall River.. | 37°39' | 96°04' | 943.11 | ----- | 04:10 | 15,000 | .04 | On east flank of Nemaha uplift. Bubble gage. On crest of Precambrian rise. Bubble gage. |
| 7-1685 | Fall River near Fall River..... | 37°38' | 96°03' | 898 | 3.81 | 04:10 | 14 | .14 | |
| 7-1800 | Cottonwood River near Marion..... | 38°21' | 97°04' | 1,289.85 | 1.84 | 03:55 | 13 | .03/.06 | |
| 7-1832 | Neosho River near Chanute..... | 37°43'49" | 95°26'28" | 887.94 | 7.75 | 04:05 | 71 | .00/.13 | |
| Kentucky | | | | | | | | | |
| 3-2808 | Buckhorn Reservoir at Buckhorn..... | 37°20'24" | 83°28'13" | MSL | 766.70 | 03:30 | 17,000 | 0.57 | Reservoir covers about 5,800 acres. At east end of Moor-man syncline. Reservoir covers about 5,000 acres. On a northeast-trending fault. |
| 3-2960 | Plum Creek subwatershed 4 near Simpsonville..... | 36°10'27" | 85°22'05" | 687.99 | 15.84 | 03:45 | 88 | .02 | |
| 3-3109 | Nolin River Reservoir near Kyrock.. | 37°16'40" | 86°14'51" | MSL | 514.38 | 03:40 | 200,000 | .40 | |
| 3-3180.06 | Rough River Reservoir near Falls of Rough. | 37°37'11" | 86°29'59" | MSL | 462.43 | 04:00 | 19,000 | .02 | |
| Louisiana | | | | | | | | | |
| 2-4895 | Pearl River near Bogalusa..... | 30°47'35" | 89°49'15" | 55.00 | 19.15 | 04:30 | 31,000 | 0.34 | Float gage. |
| 2-4900 | Bogue Lusa Creek near Franklinton.. | 30°52'05" | 90°00'10" | 210.56 | 1.87 | 03:55 | 12 | .02 | |
| 2-4901.05 | Bogue Lusa Creek at Hwy 439 at Bogalusa. | 30°46'58" | 89°52'24" | 76.60 | 4.10 | 04:00 | 120 | .00/.03 | |
| 2-4920 | Bogue Chitto near Bush..... | 30°37'45" | 89°53'50" | 44.25 | 6.20 | 04:00 | 2,000 | .62 | Between a dome and a basin. |
| 7-3444.5 | Paw Paw Bayou near Greenwood..... | 32°31'00" | 93°58'20" | 170.35 | 2.77 | 03:40 | 23 | .05 | |
| 7-3470 | Kelly Bayou near Hosston..... | 32°51'25" | 93°52'20" | 165.53 | 3.18 | 04:00 | 70 | .05 | |
| 7-3487 | Bayou Dorcheat near Springhill..... | 32°59'40" | 93°23'45" | 173.91 | 9.08 | ----- | 450 | .15 | |
| 7-3488 | Flat Lick Bayou near Leton..... | 32°46'10" | 93°16'00" | 182.79 | 3.82 | 04:00 | 40 | .03 | |
| 7-3490 | Bayou Dorcheat near Minden (auxiliary). | 32°38'40" | 93°20'15" | ----- | 6.90 | ----- | ----- | .14 | |
| 7-3498 | Cypress Bayou near Benton..... | 32°43'20" | 93°41'15" | 165.98 | 4.48 | 03:45 | 94 | .07 | |
| 7-3500 | Loggy Bayou near Ninock..... | 32°14'10" | 93°25'35" | ----- | 19.75 | 04:00 | ----- | .28 | |
| 7-3510 | Auxiliary..... | 32°11'40" | 93°28'30" | ----- | 18.90 | 04:00 | ----- | .68 | |
| 7-3517 | Boggy Bayou near Keithville..... | 32°22'35" | 93°49'20" | 145.13 | 9.87 | 06:00(?) | 14 | .05 | |
| 7-3519 | Bayou Na Bonchasse near Mansfield.. | 32°06'05" | 93°41'45" | 165.78 | 2.34 | 04:00 | 4 | No seiche | |
| 7-3519 | Bayou Dupont near Marthaville..... | 31°42'00" | 93°22'45" | ----- | 1.90 | ----- | ----- | .02 | |
| 7-3520 | Bayou Dupont near Robeline..... | 31°42'15" | 93°19'38" | 123.51 | 1.83 | 04:00 | 6 | .07 | |
| 7-3528 | Saline Bayou near Lucky..... | 32°15'00" | 92°58'35" | 162.65 | 3.68 | 04:10 | 55 | .05 | |
| 7-3528 | Grand Bayou near Coushatta..... | 32°02'55" | 93°18'10" | 136.26 | 2.25 | 03:45 | 25 | .02 | |
| 7-3530 | Saline Bayou near Clarence..... | 31°49'05" | 92°56'55" | 72.75 | 10.0 | 04:00 | 900 | .12 | |
| 7-3545 | Auxiliary..... | 31°49' | 92°56' | 72.97 | 7.85 | 04:10 | ----- | .18 | |
| 7-3545 | Horsepen Creek near Provencal..... | 31°36'05" | 93°12'05" | 149.06 | 2.31 | ----- | ----- | No seiche | |
| 7-3641 | Ouachita River near Arkansas-Louisiana State Line..... | 33°01'55" | 92°05'10" | 44.09 | 20.10 | 04:20 | ----- | .00/.10 | |
| 7-3642 | Bayou Bartholomew near Jones..... | 32°59'25" | 91°39'20" | 79.21 | 15.00 | 03:50 | 2,390 | .17 | |
| 7-3643 | Chemin-a-Haut Bayou near Beekman. | 32°58'55" | 91°48'20" | 85.58 | 2.66 | 04:30 | 31 | .06 | |
| 7-3645 | Bayou Bartholomew near Beekman..... | 32°52'20" | 91°52'04" | 70.60 | 11.5 | 04:00 | ----- | .26 | |
| 7-3647 | Bayou de Loutre near Laran..... | 32°57'20" | 92°30'00" | 112.34 | 3.06 | 04:10 | 118 | .04 | |
| 7-3650 | Bayou D'Arbonne near Dubach..... | 32°40'50" | 92°39'10" | 83.25 | 6.7 | 04:10 | 200 | .08 | |
| 7-3662 | Little Corney Bayou near Lillie..... | 32°55'40" | 92°37'55" | 91.48 | 3.88 | 04:10 | 100 | .14 | |
| 7-3677 | Boeuf River near Arkansas-Louisiana State line. | 32°58'35" | 91°26'20" | 74.11 | 3.07 | 04:50 | 580 | .57 | |
| 7-3695 | Auxiliary..... | 32°57'35" | 91°27'35" | 74.35 | 2.60 | 04:50 | ----- | .00/.09 | |
| 7-3695 | Tensas River at Tendal..... | 32°25'55" | 91°22'00" | 50.07 | 6.65 | ----- | 70 | .05? | |
| 7-3695 | Auxiliary..... | 32°23'35" | 91°19'55" | 50.07 | 5.78 | 04:00 | 70 | .00/.20 | |
| 7-3697 | Bayou Macon near Kilbourne..... | 32°59'35" | 91°15'45" | 77.41 | 2.07 | 03:50 | 250 | .08 | |
| 7-3700 | Bayou Macon near Delhi..... | 32°27'25" | 91°28'30" | 50.05 | 7.04 | 04:00 | 450 | .28 | |
| 7-3705 | Castor Creek near Grayson..... | 32°04'55" | 92°12'25" | 89.89 | 6.15 | 04:10 | 200 | .06 | |
| 7-3722 | Little River near Rochelle..... | 31°45'15" | 92°20'40" | 24.79 | 16.68 | 04:00 | 1,400 | .41 | |
| 7-3725 | Auxiliary..... | 31°47'25" | 92°21'40" | 24.79 | 17.72 | 04:10 | 1,400 | .28 | |
| 7-3730 | Bayou Funny Louis near Trout..... | 31°43'00" | 92°13'20" | 81.51 | 2.92 | 03:50 | 42 | .08 | |
| 7-3730 | Big Creek at Pollock..... | 31°32'10" | 92°24'30" | 76.79 | 2.24 | 03:40 | 40 | .08 | |
| 7-3750 | Tchefuncta River near Folsom..... | 30°36'55" | 90°14'55" | 62.11 | 7.15 | 03:50 | 175 | .23 | |
| 7-3758 | Ticklaw River at Liverpool..... | 30°55'47" | 90°40'41" | 206 | 2.37 | 04:10 | 68 | .19 | |
| 7-3780 | Comite River trib. at Sharp Station Pond near Baton Rouge. | 30°28'45" | 91°03'23" | ----- | 1.95 | 04:00 | ----- | .12 | |
| 7-3780 | Comite River near Comite..... | 30°30'45" | 91°04'25" | 25.85 | -.29 | 03:50 | 240 | .52(+?) | |
| 7-3813 | Bayou Lafourche at Golden Meadow. | 29°23'25" | 90°15'55" | MSL | .20 | 04:00 | ----- | .52/.00 | |
| 7-3820 | Bayou Cocodrie near Clearwater..... | 31°00'00" | 92°22'46" | 40.00 | 13.67 | 03:40 | 815 | .00/.02 | |
| 7-3825 | Cocodrie Lake near Clearwater..... | 31°00'00" | 92°22'57" | ----- | 13.85 | 04:20 | ----- | .35 | |
| 7-3825 | Bayou Courtaubeau at Washington.. | 30°37'05" | 92°03'20" | MSL | 19.22 | 04:00 | 1,200 | .11/.19 | |
| 7-3835 | Bayou des Blaises diversion channel at Moreauville. | 31°01'59" | 91°58'57" | 28.30 | 8.40 | 04:30 | 480 | .10 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|---|--|-----------|-----------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| Louisiana—Continued | | | | | | | | | |
| 7-3840 | Twelve mile Bayou near Dixie | 32°38'45" | 93°52'40" | 140.00 | 4.63 | 04:10 | 1,400 | 0.14 | On south side of dome. Float gage. Sharp change in water-level trend after seiche. Bubble gage. |
| 7-3855 | Bayou Teche at Arnaudville | 30°23'50" | 91°53'50" | MSL | 13.60 | 03:55 | 1,140 | .11/.17 | |
| 7-3866 | Bayou Bourbeau at Shuteston | 30°25'40" | 92°05'30" | 27.14 | 2.00 | 04:00 | .2 | .02 | |
| 7-3867 | Ruth Canal near Ruth | 30°14'35" | 91°53'05" | MSL | 10.45 | 03:40 | 186 | .09/.15 | Chart time not corrected. Two possible earthquake effects. Float gage. |
| 8-0120 | Bayou Nezpique near Basile | 30°28'50" | 92°37'55" | 3.39 | 8.95 | 03:55 | ----- | .22 | |
| 8-0130 | Calcasieu River near Glenmora | 30°59'45" | 92°40'25" | 110.77 | 9.25 | 04:00 | ----- | Tr. | |
| 8-0135 | Calcasieu River near Oberlin | 30°38'25" | 92°48'50" | 39.43 | 8.96 | 03:50 | 920 | .20 | |
| 8-0140 | Six mile Creek near Sugartown | 30°48'52" | 92°55'34" | 82.16 | 3.70 | 04:05 | 165 | .08 | |
| 8-0142 | Ten mile Creek near Elizabeth | 30°50'11" | 92°52'28" | 94.38 | 3.76 | ----- | 82 | ----- | |
| 8-0145 | Whiskey Chitto Creek near Oberlin | 30°41'55" | 92°53'35" | 46.24 | 5.07 | 04:05 | 450 | 10/.13 | |
| 8-0148 | Bundick Creek near De Ridder | 30°49'09" | 93°13'51" | 113.75 | 3.81 | 04:00 | 92 | .14 | |
| 8-0150 | Bundick Creek near Dry Creek | 30°40'55" | 93°02'15" | 56.92 | 3.90 | 04:00 | 170 | .12 | |
| 8-0155 | Calcasieu River near Kinder | 30°30'10" | 92°54'55" | 11.95 | 6.80 | 04:00 | 1,800 | .04 | |
| 8-0160 | English Bayou near Lake Charles | 30°16'17" | 93°10'37" | MSL | 1.99 | 04:00 | ----- | .24 | |
| 8-0164 | Beckwith Creek near De Quincy | 30°28'15" | 93°21'35" | 25.29 | 3.40 | 04:00 | 54 | .05 | |
| 8-0168 | Bear Head Creek near Starks | 30°13'50" | 93°37'44" | 16.34 | 9.14 | 04:00 | 56 | .12 | |
| 8-0230 | Bayou Castor near Logansport | 31°58'25" | 93°58'10" | 171.20 | 2.65 | 04:00 | 12 | .03 | |
| 8-0235 | Bayou San Patricio near Noble | 31°43'15" | 93°42'25" | 169.73 | 5.16 | 04:15 | 64 | .04 | |
| 8-0240, 6 | Blackwell Creek at Many | 31°34'50" | 93°27'45" | 224.12 | 2.35 | 04:00 | 3 | .04 | |
| 8-0255 | Bayou Toro near Toro | 31°18'25" | 93°30'58" | 138.00 | 4.30 | 03:50 | 80 | .15 | |
| 8-0275 | Bayou Anacoco near Leesville | 31°09'35" | 93°21'05" | 190.58 | 6.82 | 03:55 | 212 | .07 | |
| 8-0280 | Bayou Anacoco near Rosepine | 30°57'10" | 93°21'10" | 118.09 | 6.23 | 03:55 | 380 | .16 | |
| Maine | | | | | | | | | |
| No seiche was recorded at any gaging station. | | | | | | | | | |
| Maryland | | | | | | | | | |
| 1-4900 | Chicamacomico River near Salem | 38°30'45" | 75°52'50" | 10 | 1.85 | 03:50 | 30 | 0.04 | |
| 1-5892 | Gwynns Falls near Owings Mills | 39°26'16" | 76°46'57" | 520 | 1.24 | 03:50 | 4.0 | .006 | |
| 1-5948 | St. Leonard Creek near St. Leonard | 38°26'57" | 76°29'43" | 5 | 2.94 | 04:10 | 7.6 | .01 | |
| Massachusetts | | | | | | | | | |
| No seismic seiche was recorded at any gaging station. | | | | | | | | | |
| Michigan | | | | | | | | | |
| 4-0964 | St. Joseph River near Burlington | 42°06'10" | 85°02'25" | 930 | 2.74 | 04:00 | 140 | 0.01 | On edge of Michigan basin. |
| 4-0966 | Coldwater River near Hodunk | 42°01'45" | 85°06'25" | 900 | 2.99 | 04:00 | 120 | .01 | |
| 4-1115 | Deer Creek near Dansville | 42°36'30" | 84°19'15" | 889.08 | 2.98 | 04:00 | 5 | .01 | On edge of Michigan basin; a residual 0.01-ft rise in stage. On south side of Michigan basin; a residual 0.01-ft drop in stage. |
| 4-1120 | Sloan Creek near Williamston | 42°40'30" | 84°21'50" | 862.12 | 1.89 | 03:50 | 2.1 | .01 | Do. |
| 4-1125 | Cedar River at East Lansing | 42°43'40" | 84°28'40" | 824.39 | 3.65 | 03:40 | 115 | No seiche | Do. |
| 4-1300 | Cheboygan River near Cheboygan | 45°34'40" | 84°29'15" | 591.21 | 1.40 | 04:10 | 860 | .00/.03 | East of 10-mgal high. |
| 4-1355 | Au Sable River at Grayling | 44°38'35" | 84°42'45" | 1,123.49 | 1.28 | 03:40 | 60 | .03 | East of 0-mgal high. |
| 4-1356 | East Branch Au Sable River at Grayling | 44°40'10" | 84°42'20" | 1,110 | 3.42 | 04:10 | 34 | .05/.00 | Do. |
| 4-1460 | Farmers Creek near Leaper | 43°02' | 83°20' | 805.79 | 15.50 | 03:40 | 19 | .02 | On southeast side of Michigan basin. |
| 4-1505 | Cass River at Cass City | 43°35'10" | 83°10'35" | ----- | ----- | ? | ----- | No seiche | A residual 0.01-ft rise in stage. |
| 4-1606 | Belle River at Memphis | 42°54'03" | 82°46'09" | 720 | 1.78 | 04:00 | 27 | .02 | On southeast side of Michigan basin. |
| 4-1635 | Plum Brook near Utica | 42°35'01" | 83°01'49" | 610 | 1.58 | 03:40 | 12 | .015 | Do. |
| 4-1640, 1 | North Branch Clinton River at Almont | 42°54'59" | 83°02'42" | 830 | 2.95 | 04:00 | 2 | .01 | Do. |
| 4-1644 | Deer Creek near Meade | 42°42'39" | 82°51'32" | 610 | .70 | 04:00 | .8 | .02 | Do. |
| 4- | Kent Lake near New Hudson | 42°30'45" | 83°40'35" | 868.00 | 13.55 | 04:00 | ----- | .07 | On Howell anticline. On south east side of Michigan basin and 10-mgal high. |
| ----- | Reservoirs of City of Lansing | 42° | 84° | ----- | ----- | 03:55 | 21 | 1.83 | 7-million gallon reservoir. |
| ----- | | 42° | 84° | ----- | ----- | 03:55 | 30 | 1.25 | |
| Minnesota | | | | | | | | | |
| 5-1075 | Roseau River at Ross | 48°54'37" | 95°55'18" | 1,018.44 | 1.55 | 03:50 | 5.0 | 0.03 | Near edge of Cretaceous overlap. |
| Mississippi | | | | | | | | | |
| 2-4330 | Bull Mountain Creek near Smithville | 34°05' | 88°24' | 234.81 | 10.16 | 04:10 | 2,700 | 0.06 | Peseiche effect(?). |
| 2-4340 | Old Town Creek near Tupelo | 34°17'40" | 88°42'35" | 244.24 | 5.92 | 05:00 | 190 | .08 | |
| 2-4345 | Euclatubba Creek at Saffillo | 34°22'20" | 88°42'00" | 280 | 4.10 | 04:10 | 24 | .03 | |
| 2-4365 | West Fork Tombigbee River near Nettleton | 34°03'32" | 88°37'40" | 194.01 | 10.48 | 04:20 | 940 | .17 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|--------------------------------|---|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| Mississippi—Continued | | | | | | | | | |
| 2-4370 | Tombigbee River near Amory | 33°59'10" | 88°33'05" | 178.34 | 18.35 | 03:40 | 9,800 | 0.27 | |
| 2-4400 | Chookatonchee Creek near Egypt | 33°50'30" | 88°46'30" | 226.07 | 1.80 | 03:50 | 215 | .04 | |
| 2-4750 | Leaf River near McLain | 31°06'10" | 88°48'30" | 42.15 | 7.81 | 03:20 | 4,600 | .18 | On Wiggins uplift. |
| 2-4765 | Sowashee Creek at Meridian | 32°22'10" | 88°40'40" | 305.95 | 3.08 | 04:00 | 70 | Tr. | |
| 2-4790 | Pascagoula River at Merrill | 30°58'40" | 88°43'25" | 26.25 | 11.56 | 04:00 | 12,500 | .66 | Do. |
| 2-4825 | Pascagoula River at Cumbest Bluff | 30°25'10" | 88°34'20" | ----- | 9.28 | 04:00 | ----- | .37 | |
| 2-4793 | Red River at Vestry | 30°44'10" | 88°46'50" | 20.10 | 7.80 | 04:00 | 800 | .16 | |
| 2-4825.5 | Pearl River near Clarhage | 32°42'25" | 89°31'55" | 515.24 | ? | 04:30 | ? | .127 | No vertical scale on chart. |
| 2-4830 | Tuscolameta Creek at Walnut | 32°35' | 89°28' | 332.70 | 15.65 | 04:20 | ----- | .11 | |
| 2-4840 | Yockanookany River near Kosciusko | 33°02' | 89°35' | 374.94 | 9.33 | 04:30 | 340 | .02 | |
| 2-4845 | Yockanookany River near Ofahoma | 32°42'20" | 89°40'20" | 311.15 | 6.00 | 03:30 | 402 | .08 | A residual 0.03-ft rise in stage; on east edge of Ouachita tectonic belt. |
| 2-4860 | Pearl River at Jackson | 32°17'20" | 90°10'45" | 234.90 | 27.72 | 04:00 | 18,000 | .05 | On Jackson dome. |
| 2-4885 | Pearl River near Monticello | 31°33' | 90°05' | 155.66 | 21.77 | 04:00 | 22,500 | .90 | |
| 2-4892.4 | Pearl River near Monticello | 31°09'30" | 89°37'40" | 180 | 3.20 | 03:55 | 120 | .07 | |
| 2-4905 | Bogue Chitto near Tylertown | 31°11' | 90°17' | 227.40 | ----- | ----- | 600 | Tr. | Pen trace indistinct. |
| 7-2880 | Tallahatchie River at Etta | 34°29'00" | 89°13'30" | 273.48 | 11.45 | 05:00 | 980 | .26 | |
| 7-2830 | Skuna River at Bruce | 33°58'25" | 89°20'50" | 238.75 | 4.40 | 03:55 | 1,280 | .06 | |
| 7-2900 | Big Black River near Bovina | 32°20'51" | 90°41'48" | 84.93 | 26.85 | ? | 9,000 | .06 | |
| Missouri | | | | | | | | | |
| 5-5023 | Salt River at Hagers Grove | 39°49'40" | 92°14'10" | ----- | 4.12 | 04:30 | ----- | 0.06 | A residual 0.03-ft rise in stage. |
| 6-8990 | Weldon River at Mill Grove | 40°18' | 93°36' | 786.03 | .71 | 04:00 | 25 | .00/.02 | Bubble gage. |
| 6-8995 | Thompson River at Trenton | 40°04'45" | 93°38'35" | 721.87 | 3.83 | 03:50 | 113 | .02/.00 | Do. |
| 6-9067 | Flat Creek near Sedalia | 38°39'35" | 93°15'10" | 765 | 2.25 | 03:45 | 10 | .13 | |
| 6-9216 | South Grand River at Ulrich | 38°27'08" | 94°00'13" | 715.9 | 2.40 | 04:00 | 5 | .00/.04 | Do. |
| 6-9270 | Maries River at Westphalia | 38°25'55" | 91°59'20" | 542.74 | 2.25 | 03:45 | 75 | .00/.01 | |
| 6-9278 | Osage Fork at Dryrot | 37°38'00" | 92°27'12" | 927.85 | 3.79 | 04:30 | 90 | .01 | On southeast of Decaturville uplift. |
| 6-9280 | Gasconade River near Hazlegreen | 37°45'35" | 92°27'05" | 844.75 | 3.40 | 04:00 | 500 | .03 | Do. |
| 6-9285 | Gasconade River near Waynesville | 37°52'20" | 92°13'40" | 738.90 | 3.30 | 03:50 | 720 | .03 | Do. |
| 6-9355 | Loutre River at Mineola | 38°53'20" | 91°34'30" | 539.86 | 3.29 | 03:50 | 40 | .02 | |
| 7-0210 | Castor River at Zalma | 37°08'45" | 90°04'30" | 350.38 | 5.58 | 04:30 | 500 | .04 | On southeast of domal structure. |
| 7-0375 | St. Francis River near Patterson | 37°11'40" | 90°30'10" | 370.45 | 6.25 | 04:30 | 1,600 | .04 | Do. |
| 7-0395 | St. Francis River at Wappapello | 36°55'42" | 90°17'04" | ----- | 13.15 | 04:00 | ----- | .12 | At edge of Tertiary overlap. |
| 7-0435 | Little River Ditch 1 near Morehouse | 36°00'05" | 90°43'50" | 280.76 | 5.98 | 04:00 | 600 | .05 | Near edge of Tertiary overlap. |
| 7-0630 | Black River at Poplar Bluff | 36°45'35" | 90°23'15" | 317.38 | 8.50 | 04:15 | 760 | .87 | At edge of Tertiary overlap. |
| 7-1866 | Turkey Creek near Joplin | 37°07'15" | 94°34'55" | 848.80 | 1.96 | 04:10 | 11 | .02 | |
| ----- | Headwater Diversion Channel at Dutchtown | 37°13'54" | 89°39'31" | ----- | 8.70 | 04:30 | ----- | .26 | Seiche lasted about 40 min. On southeast of domal structure. |
| 7-1890 | Elk River near Tiff City | 36°38' | 94°35' | 750.61 | 3.28 | 03:50 | 200 | Tr. | |
| Montana | | | | | | | | | |
| 5-0145 | Swiftcurrent Creek at Many Glacier | 48°48'10" | 113°39'20" | 4,860 | 1.55 | 04:30 | 16 | 0.08 | On a thrust fault. |
| 6-0375 | Madison River near West Yellowstone | 44°39'20" | 111°04'00" | 6,650 | 1.93 | 04:10 | 378 | .07 | May lie on buried extension of thrust faults that trend northwest-southeast. This gage also recorded seiche from Lake Hebgen earthquake. |
| 6-0525 | Gallatin River at Logan | 45°53'10" | 111°28'20" | 4,082.3 | 3.33 | 04:30 | 712 | .05 | On possible extension of a thrust fault. |
| 6-1185 | South Fork of Musselshell River above Martinsdale | 46°27' | 110°23' | 4,900 | 2.47 | 03:50 | 16 | .02 | On southeast end of Little Belt uplift. |
| 6-1220 | American Fork below Lebo Creek, near Harlowtown | 46°24' | 109°46' | 4,170 | 2.25 | 03:45 | 14 | .02 | |
| 6-1235 | Musselshell River near Ryegate | 46°18' | 109°12' | 3,880 | 2.88 | 04:00 | 21 | .01 | |
| 6-1307 | Sand Creek near Jordan | 47°15' | 106°51' | 2,586.28 | 2.06 | 04:10 | ----- | .01 | South of axis of Blood Creek syncline. |
| 6-1322 | South Fork of Milk River near Babb | 48°45'20" | 113°10'00" | ----- | 2.94 | ----- | 6 | .05 | |
| 6-1975 | Boulder River near Contact | 45°33'20" | 110°12'00" | 4,930 | 1.66 | 04:00 | 56 | .015 | On extension of a small fault and on north edge of Bear-tooth uplift. |
| 6-2000 | Boulder River at Big Timber | 45°50'05" | 109°56'20" | 4,060 | 3.44 | 03:45 | 110 | .04 | On southeast end of Crazy Mountains basin. |
| 6-2890 | Little Bighorn River at State Line near Wyoala | 45°01' | 107°37' | 4,450 | 1.84 | 04:05 | 71 | .03 | On a small fault. |
| 6-3075 | Tongue River at Tongue River Dam, near Decker | 45°08' | 106°46' | 3,050 | .93 | 04:00 | 126 | .10 | On north end of Powder River basin. |
| 12-3018.5 | Kootenai River at Warland Bridge, near Libby | 48°30'00" | 115°17'10" | ----- | 5.22 | 04:00 | 2,150 | .00/.02 | Nontypical seiche with water-level decline and recovery. Bubble gage? On northeast flank of anticline. |
| 12-3235 | German Gulch Creek near Ramsey | 46°00'50" | 112°47'30" | 5,200 | 1.41 | 04:00 | 6.2 | Tr. | On a batholith. |
| 12-3588 | Middle Fork Flathead River near West Glacier | 48°29'50" | 114°00'30" | 3,130 | .90 | 04:00 | 350 | Tr. | On a normal fault. |
| 12-3895 | Thompson River near Thompson Falls | 47°35'35" | 115°13'40" | 2,410 | 1.78 | 04:05 | 115 | .04 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|---|---|-----------|------------|--------------------|------------|------------------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| Nebraska | | | | | | | | | |
| 6-4541 | Niobrara River at Agate..... | 42°25' | 103°47' | 4,440 | 2.73 | 04:10 | 23 | 0.09 | North end of Denver basin. |
| 6-6875 | North Platte River at Lewellen (North channel). | 41°19' | 102°08' | 3,284.6 | 4.20 | 04:05 | 1,200 | .085 | |
| | North Platte River at Lewellen (South channel). | 41°19' | 102°08' | 3,383.7 | 5.02 | 03:55 | | | .12 |
| 6-7635 | Lodgepole Creek at Ralton..... | 41°02'00" | 102°24'00" | 3,590 | 1.60 | 04:00 | 24 | .07 | On Cambridge arch. |
| 6-7655 | South Platte River at North Platte..... | 41°07' | 100°46' | 2,790.30 | 2.75 | 04:05 | 192 | .015 | |
| 6-7665 | Platte River at Cozad (South channel). | 40°50' | 99°59' | 2,474.07 | 4.26 | 03:55 | | .06 | |
| 6-7680 | Platte River near Overton..... | 40°41' | 99°32' | 2,299.83 | 2.78 | ----- | 1,300 | .12 | On a normal fault. |
| 6-7890 | North Loup River at Scotia..... | 41°27'30" | 98°42'40" | 1,893.13 | 2.87 | ----- | 1,100 | .08 | |
| 6-7920 | Cedar River near Fullerton..... | 41°23'45" | 98°00'15" | 1,640.40 | 2.48 | 03:50 | 330 | .05 | A residual 0.04-ft rise in stage. |
| 6-8050 | Salt Creek at Ashland..... | 41°02'50" | 96°20'30" | 1,047.04 | 2.28 | 04:05 | 236 | .10 | |
| 6-8490 | Harlan County Reservoir near Republican City. | 40°04'10" | 99°12'30" | MSL | 1,939.72 | 03:40 | 267,100 | .075 | On a dome. |
| 6-8810 | Big Blue River near Crete..... | 40°35'40" | 96°57'35" | 1,311.7 | ? | 03:50 | 132 | .025 | |
| 6-8829 | Little Blue River below Pawnee Creek near Pauling. | 40°23'50" | 98°13'20" | 1,740 | 3.52 | 04:00 | 65 | .06 | |
| 6-8830 | Little Blue River near Deweese..... | 40°20'00" | 98°04'10" | 1,632.67 | 3.35 | 04:00 | 72 | .01 | |
| Nevada | | | | | | | | | |
| No seiche was recorded at any gaging station. | | | | | | | | | |
| New Hampshire | | | | | | | | | |
| 1-0535 | Androscoggin River at Errol..... | 44°46'55" | 71°07'45" | 1,227.30 | ----- | 04:20 | 2,200 | Tr. | |
| New Jersey | | | | | | | | | |
| 1-3830 | Greenwood Lake at Awosting..... | 41°09'36" | 74°20'03" | 608.86 | 10.20 | 04:00 | 20,000 | 0.08 | In Green Pond syncline. |
| New Mexico | | | | | | | | | |
| 7-1535 | Cimarron River near Guy..... | 36°59'15" | 103°25'25" | 4,900 | 0.63 | 04:10 | 1 | 0.02 | On a normal fault. |
| 7-2050 | Six Mile Creek near Eaglenest..... | 36°31'09" | 105°16'30" | 8,195.16 | .75 | 04:10 | 3 | .01 | |
| 7-2062 | McEvoy Creek near Eaglenest..... | 36°33'00" | 105°13'30" | 8,600 | .36 | 04:10 | .1 | No seiche | |
| 7-2070 | Cimarron Creek near Cimarron..... | 36°31'00" | 104°58'35" | 6,599.58 | .79 | 03:45 | 2 | .01 | A lasting 0.002-ft drop in stage. On fault between volcanics and Precambrian. |
| 7-2085 | Rayado Creek at Sauble Ranch, near Cimarron. | 36°22' | 104°58' | 6,880 | 1.78 | 03:40 | 4 | .01 | |
| 7-2165 | Mora River near Golondrinas..... | 35°53'40" | 105°09'30" | 6,734.1 | 1.75 | ----- | 4 | .00/.03 | (On fault at contact of volcanics and Precambrian. |
| 7-2171 | Coyote Creek above Guadalupe..... | 36°10'30" | 105°13'35" | 7,700 | 1.53 | {03:55 04:40} | 3 | .01/.02 | |
| 7-2210 | Mora River near Shoemaker..... | 35°48' | 104°47' | 6,170 | .11 | 04:00 | 2 | .10 | At edge of volcanics. |
| 7-2245 | Canadian River below Conchas Dam. | 35°24'30" | 104°10'10" | 4,121.90 | 4.72 | 04:00 | 6 | .06 | |
| 8-2635 | Rio Grande near Cerro..... | 36°44'05" | 105°41'05" | 7,100 | 3.07 | 03:55 | 270 | .26 | On east edge of volcanics. |
| 8-2645 | Red River below Zwergle Dam Site, near Red River. | 36°40'25" | 105°22'50" | 8,871.88 | 1.70 | 03:50 | 4 | .02 | |
| 8-2650 | Red River near Questa..... | 36°42'10" | 105°34'03" | 7,451.92 | 2.05 | 04:10 | 12 | .03 | On volcanics near a fault. |
| 8-2675 | Rio Hondo near Valdez..... | 36°32'30" | 105°33'20" | 7,650.0 | 1.72 | 03:20 | 7 | .03 | |
| 8-2763 | Rio Pueblo de Taos below Los Cordovas. | 36°22'38" | 105°40'04" | 6,650 | 2.08 | 03:50 | 24 | .03 | On contact of Precambrian and Tertiary. |
| 8-2842 | Willow Creek above Heron Reservoir, near Park View. | 36°44'30" | 106°37'35" | 7,210 | .56 | ----- | 2 | .02 | |
| 8-2855 | Rio Chama below El Vado Dawn..... | 36°34'50" | 106°43'30" | 6,696.12 | 1.55 | 03:40 | 62 | .03 | Do. |
| 8-3145 | Rio Grande at Cochiti..... | 35°37'10" | 106°19'20" | 5,224.70 | 3.77 | 04:00 | 470 | .08 | |
| 8-3295 | Rio Grande near Bernalillo (site B)..... | 35°17' | 105°35' | 5,030.57 | 2.05 | 04:10 | 100 | .04 | A lasting 0.005-ft drop in stage. On southeast edge of volcanics. |
| 8-3320 | Bernardo Interior Drain near Bernardo. | 34°25' | 103°48' | 4,713.99 | 6.00 | 04:20 | ----- | .03 | |
| 8-3435 | Rio San Jose near Grants..... | 35°04'30" | 107°45'00" | 6,269.47 | 1.41 | 04:00 | 5 | No seiche | |
| 8-3875 | San Antonio Drain near San Marcial. | 33°44'45" | 106°55'15" | 4,489.12 | 3.74 | 03:50 | ----- | .03 | |
| 8-3810 | Gallinas River at Montezuma..... | 35°39'15" | 105°16'30" | 6,675 | 3.93 | 04:00 | 2 | .02 | |
| 8-3860 | Pecos River near Acme (auxiliary)..... | 33°32'10" | 104°22'40" | 3,500 | 3.26 | 04:20 | 8 | .01 | Do. |
| 8-3995 | Pecos River (Kaiser Channel) near Lakewood. | 32°41'22" | 104°17'53" | 3,268.53 | 1.92 | 03:45 | 22 | .04 | |
| 8-4050 | Pecos River at Carlsbad..... | 32°25'05" | 104°13'25" | 3,080.28 | 1.14 | 04:00 | 30 | .04 | |
| 8-4055 | Black River above Malaga..... | 32°13'40" | 104°09'05" | 3,070 | .66 | 03:50 | 3 | .01 | |
| 8-4085 | Delaware River near Red Bluff..... | 32°01'25" | 104°03'15" | 2,900.66 | ----- | 04:10 | 1 | .04 | |
| New York | | | | | | | | | |
| 1-3874.5 | Mahwah River near Suffern..... | 41°08'27" | 74°07'01" | 325 | ----- | 04:00 | 33 | No seiche | A lasting 0.01-ft drop in stage. In Great Pond syncline. |
| 1-3710 | Shawangunk Kill at Pine Bush..... | 41°37'05" | 74°17'40" | 305 | ----- | 04:00 | 130 | Tr. | |
| 1-4240 | Trout Creek near Rock Royal..... | 42°10'40" | 75°16'45" | 1,165.70 | ----- | 04:00 | 100 | Tr. | |
| 1-4365 | Neversink River at Woodbourne..... | 41°45'25" | 74°35'55" | 1,180 | ----- | 04:00 | 80 | Tr. | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|--------------------------------|--|-----------|-----------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| North Carolina | | | | | | | | | |
| ----- | Fontana Dam Hydro Plant head-water. | 35° | 83° | 1,689.91 | ----- | ----- | 1,000,000 | 0.05 | |
| North Dakota | | | | | | | | | |
| 5-0590 | Sheyenne River near Kindred..... | 46°37'35" | 97°00'05" | 925.55 | 3.45 | 03:00 | 47 | 0.06 | Do. |
| 6-4690 | Jamestown Reservoir near Jamestown. | 46°56'03" | 98°42'38" | MSL | 1,425.44 | 03:50 | 21,000 | No seiche | A lasting 0.08-ft drop in stage. On southeast side of Williston basin. |
| 6-4705 | Jamestown River at La Moure..... | 46°21'20" | 98°18'15" | 1,290.00 | 7.20 | ----- | 57 | Tr. | On southeast side of Williston basin. |
| Ohio | | | | | | | | | |
| 3-0865 | Mahoning River at Alliance..... | 40°55'55" | 81°05'45" | 1,037.3 | 1.75 | 04:00 | 77 | Tr. | Near edge of Pennsylvanian overlap. |
| 3-0910 | Milton Reservoir at Pricetown..... | 41°07'40" | 80°58'35" | MSL | 47.00 | 04:10 | 43,000 | 0.07 | |
| 3-0920 | Kale Creek near Pricetown..... | 41°08'25" | 80°59'45" | 914.7 | 1.10 | 03:50 | 13 | .04 | On east of 20-mgal high. |
| 3-1180 | Middle Branch Nimishillen Creek at Canton. | 40°50'30" | 81°21'20" | 1,046.6 | 1.64 | 04:20 | 25 | .03 | |
| 3-1200 | Leesville Reservoir near Leesville.... | 40°28'10" | 81°11'45" | 928.0 | 36.10 | 04:15 | 8,000 | .04 | Near top of 10-mgal high. On south edge of Michigan Basin and on northwest side of Findlay arch. |
| 3-1280 | Tappan Reservoir at Tappan..... | 40°21'35" | 81°13'35" | 870.0 | 28.55 | 04:00 | 25,000 | .06 | |
| 3-1313 | Black Fork at Melco..... | 40°41'55" | 82°21'35" | ----- | 4.63 | 04:10 | ----- | .03 | Bubble gage(?). |
| 3-1685 | Burr Oak Reservoir at Burr Oak..... | 39°32'35" | 82°03'30" | MSL | 721.40 | 03:50 | 9,400 | .10 | |
| 3-2205 | O'Shaughnessy Reservoir near Dublin. | 40°09'15" | 83°07'34" | MSL | 848.75 | 04:20 | 17,500 | .08 | |
| 3-2210 | Scioto River below O'Shaughnessy Reservoir. | 40°08'36" | 83°07'14" | 775.00 | 5.50 | 03:20 | ----- | .04 | Bubble gage(?). |
| 3-2215 | Griggs Reservoir near Columbus..... | 40°00'54" | 83°05'38" | 630.38 | ----- | 04:00 | 4,820 | .02 | |
| 3-2284 | Hoover Reservoir at Central College. | 40°06'30" | 82°53'00" | ----- | 90.20 | 03:50 | 60,000 | .03 | Bubble gage(?). |
| 3-2305 | Big Darby Creek at Darbyville..... | 39°42'05" | 83°06'35" | 713.6 | 3.00 | 03:50 | 490 | .08 | |
| 3-2340 | Paint Creek near Bourneville..... | 39°15'49" | 83°10'01" | 665.2 | 7.13 | 04:00 | 1,650 | .14 | Bubble gage(?). |
| 3-2395 | North Fork Little Miami River near Pritchlin. | 39°49'40" | 83°46'25" | 1,011.46 | 1.95 | 03:00 | ----- | .01 | |
| 3-2440 | Todd Fork near Roachester..... | 39°20'05" | 84°05'10" | 679.40 | 6.60 | 03:30 | 370 | .03 | Bubble gage(?). |
| 3-2665 | West Fork Mill Creek Reservoir near Greenhills. | 39°15'40" | 84°29'40" | 600.00 | 75.05 | 04:30 | 1,500 | .09 | |
| 3-2580 | West Fork Mill Creek at Lockland.... | 39°13'35" | 84°27'20" | 539.00 | 4.20 | 04:00 | ----- | .01 | Bubble gage(?). |
| 3-2640 | Greenville Creek near Bradford..... | 40°06'08" | 84°25'48" | 948.9 | 2.27 | 04:00 | 160 | .03 | |
| 3-2728 | Sevenmile Creek at Collinsville..... | 39°31'23" | 84°36'39" | 691.95 | 2.00 | 04:00 | 86 | .01 | Bubble gage(?). |
| 4-1920 | Miami and Erie Canal near Defiance. | 41°17'30" | 84°18'50" | 656.12 | 1.60 | 04:00 | 11 | .03 | |
| 4-1925 | Maumee River near Defiance..... | 41°17'30" | 84°18'50" | 659.12 | ----- | 03:50 | ----- | .02 | Bubble gage(?). |
| 4-1965 | Sandusky River near Upper Sandusky. | 40°51'02" | 83°15'23" | 792.8 | 2.78 | 03:50 | 520 | .03 | |
| 4-2115 | Mill Creek near Jefferson..... | 41°45'10" | 80°48'00" | 822.59 | 2.59 | 04:00 | 160 | .00/.04 | Bubble gage(?). |
| ----- | Mill Creek near Jefferson Lake gage.. | 41°45'20" | 80°48'00" | ----- | 0.62 | 03:50 | ----- | .25 | |
| Oklahoma | | | | | | | | | |
| 7-1505 | Salt Fork of Arkansas River near Jet. | 36°45' | 98°08' | 1,092.20 | 4.23 | 04:00 | 40 | 0.04 | Two seiches(?). |
| 7-1510 | Salt Fork of Arkansas River at Tonkawa. | 36°40'30" | 97°18'40" | 930.22 | 4.50 | 04:05 | 74 | .02 | |
| 7-1650 | Heyburn Reservoir near Heyburn..... | 35°57' | 96°18' | MSL | 760.33 | 03:55 | 7,100 | .20 | Bubble gage. |
| 7-1655 | Snake Creek near Bixby..... | 35°49'10" | 95°53'20" | 625 | 2.41 | 04:00 | 2 | .01 | |
| 7-1713 | Oologah Reservoir near Oologah..... | 36°25'19" | 95°40'43" | MSL | 607.06 | 04:20 | 52,730 | .06 | Bubble gage. |
| 7-1725 | Hulah Reservoir near Hulah..... | 36°56' | 96°05' | MSL | 726.40 | 04:05 | 15,450 | .055 | |
| 7-1746 | Sand Creek at Okesa..... | 36°43'10" | 96°07'58" | 689.20 | 2.88 | 03:50 | 1 | .00/.04 | Bubble gage. |
| 7-1760 | Verdigris River near Claremore..... | 36°18'30" | 95°41'40" | 538.62 | 3.90 | 04:05 | 26 | .00/.02 | |
| 7-1765 | Bird Creek at Avant..... | 36°29' | 96°04' | 615.28 | 2.46 | 03:50 | 1.1 | .06 | Unusual rise in stage 40 min before earthquake was recorded. Near Seneca Fault. |
| 7-1775 | Bird Creek near Sperry..... | 36°16'42" | 95°57'14" | 579.43 | 1.21 | 04:15 | 9.7 | .015 | |
| 7-1900 | Lake O' The Cherokees at Langley.. | 36°28' | 95°02' | MSL | 730.90 | 04:00 | 1,117,000 | .44 | |
| 7-1912.2 | Spavinaw Creek near Sycamore..... | 36°20'00" | 94°38'30" | 875 | 2.67 | 04:00 | 30 | .01 | On a normal fault. |
| 7-1930 | Fort Gibson Reservoir near Fort Gibson. | 35°52' | 95°14' | MSL | 551.70 | 04:00 | 323,000 | .12 | |
| 7-1955 | Illinois River near Watts..... | 36°07'48" | 94°34'12" | 893.78 | 2.30 | 04:00 | 126 | .11 | Do. |
| 7-1960 | Flint Creek near Kansas, Okla..... | 36°11'54" | 94°42'30" | 854.59 | 6.27 | 04:00 | 40 | .13 | |
| 7-1965 | Illinois River near Tahlequah..... | 35°55' | 94°55' | 664.14 | 4.05 | 04:30 | 320 | .11 | Do. |
| 7-1970 | Barren Fork at Eldon..... | 35°55' | 94°50' | 701.14 | 4.88 | 03:40 | 90 | .04 | |
| 7-2305 | Little River near Tecumseh..... | 35°10'25" | 96°55'55" | 898.52 | 4.46 | 04:10 | 5.4 | .03 | Do. |
| 7-2315 | Canadian River near Calvin..... | 34°58' | 96°14' | 684.72 | 1.61 | 04:00 | 63 | .00/.02 | |
| 7-2365 | Fort Supply Reservoir near Fort Supply. | 36°33' | 99°34' | MSL | 2,001.93 | 04:15 | 11,010 | .055 | Do. |
| 7-2375 | North Canadian River at Woodward. | 36°26' | 99°17' | 1,830.43 | 3.83 | 03:40 | 36 | .01 | |
| 7-2395 | North Canadian River near El Reno. | 35°33'44" | 97°57'32" | 1,299.02 | 5.12 | 03:20 | 14 | .02 | Do. |
| 7-2400 | Lake Hefner Canal near Oklahoma City. | 35°33'11" | 97°37'11" | 1,200.96 | 5.14 | 03:40 | 2 | .00/.015 | |
| 7-2410 | North Canadian River below Lake Overholser near Oklahoma City. | 35°28'44" | 97°39'47" | 1,194.66 | 10.74 | 03:40 | 1.4 | .12 | Do. |
| 7-2450 | Canadian River near Whitefield..... | 35°15'45" | 95°14'20" | 478.16 | 4.97 | 03:55 | 8.3 | .02 | |
| 7-2455 | Sallisaw Creek near Sallisaw..... | 35°28' | 94°52' | 474.78 | 2.48 | 04:10 | 35 | Tr.? | Do. |
| 7-2465 | Arkansas River near Sallisaw..... | 35°21' | 94°46' | 413.42 | ? | 04:00 | 1,870 | .05? | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|--|--|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| Oklahoma—Continued | | | | | | | | | |
| 7-2480 | Wister Reservoir near Wister..... | 34°56'10" | 94°43'10" | MSL | 471.60 | 03:50 | 30,080 | 0.13 | |
| 7-3025 | Lake Altus at Lugert..... | 34°54' | 99°18' | MSL | 1,544.85 | 04:00 | 68,480 | 2.9 | On Wichita Mountains uplift. |
| 7-3165 | Washita River near Cheyenne..... | 35°37'35" | 99°40'05" | 1,905.98 | 2.14 | 04:00 | 3.5 | .02 | |
| 7-3250 | Washita River near Clinton..... | 35°31'50" | 98°58'00" | 1,467.60 | 5.26 | 04:00 | 10 | .04 | |
| 7-3335 | Chickasaw Creek near Stringtown..... | 34°27'41" | 96°01'36" | 540.26 | 3.45 | 03:45 | 5.0 | .02 | On a thrust fault. |
| 7-3340 | Muddy Boggy Creek near Farris..... | 34°16'17" | 95°54'43" | 444.53 | 3.10 | 03:55 | 67 | No seiche | A lasting 0.06-ft drop in stage. Bubble gage. |
| 7-3342 | Byrds Mill Spring near Fittstown..... | 34°35'45" | 96°39'55" | 1,022 | 2.7 | 04:00 | 1.4 | No seiche | A lasting 0.15-ft drop in stage; after 80 min water level had recovered to preearthquake level. Float gage. On normal fault at west end of a graben. |
| 7-3375 | Little River near Wright City..... | 34°04'10" | 95°02'47" | 346.76 | 6.89 | 04:00 | 380 | No seiche | A lasting 0.01-ft drop in stage. Bubble gage. |
| 7-3379 | Glover Creek near Glover..... | 34°08'51" | 94°54'07" | 378.70 | 4.05 | 04:00 | 350 | .00/.05 | |
| | Lake Shawnee near Shawnee..... | 35°20'50" | 97°03'45" | MSL | 7733.53 | 04:00 | ? | .21 | |
| Oregon | | | | | | | | | |
| 14-0260 | Umatilla River at Yoakum..... | 45°40'40" | 119°02'00" | 768.21 | 2.58 | 04:10 | 550 | 0.03 | |
| 14-0525 | Quinn River near Lapine..... | 43°47'10" | 121°50'10" | 4,442.1 | 1.32 | 05:40 | 17 | .04 | Poor copy. |
| 14-0575 | Fall River near Lapine..... | 43°47'50" | 121°34'20" | 4,220 | 1.32 | 05:40 | 150 | .04 | |
| 14-1134 | Dog River near Parkdale..... | 45°24'30" | 121°31'10" | 4,347 | 2.45 | 03:30 | 2.8 | .02 | Near a normal fault. |
| 14-1451 | Hills Creek Reservoir near Oakridge..... | 43°42'30" | 122°25'25" | MSL | 1,508 | 03:50 | 271,600 | .11 | Seiche lasted about 80 min. |
| 14-1490 | Lookout Point Reservoir near Lowell..... | 43°54'50" | 122°45'00" | MSL | 878.8 | 03:40 | 258,000 | .06 | Seiche lasted at least 100 min. |
| 14-1530 | Cottage Grove Reservoir near Cottage Grove..... | 43°43'00" | 123°02'55" | MSL | 876.3 | 03:50 | 18,000 | .05 | Seiche lasted about 30 min. |
| 14-1550 | Dorena Reservoir near Cottage Grove..... | 43°47'10" | 122°57'15" | MSL | 810.9 | 03:40 | 41,000 | Tr. | |
| 14-1585 | McKenzie River at outlet of Clear Lake..... | 44°21'40" | 121°59'40" | 3,015.32 | 2.24 | 04:00 | 300 | .02 | |
| 14-1594 | Cougar Reservoir near Rainbow..... | 44°06'15" | 122°14'20" | MSL | 1,606.5 | 03:50 | 121,000 | .09 | Seiche lasted about 60 min. |
| 14-1680 | Fern Ridge Reservoir near Elmira..... | 44°07'15" | 123°18'00" | MSL | 369 | ? | 72,000 | Tr. | |
| 14-1700 | Long Tom River at Monroe..... | 44°18'50" | 123°17'45" | 270.57 | 4.60 | ? | 210 | Tr. | |
| 14-1735 | Calapoia River at Albany..... | 44°37'15" | 123°07'40" | 180.85 | 4.90 | 03:30 | 600 | Tr. | |
| 14-1805 | Detroit Reservoir near Detroit..... | 44°43'20" | 122°14'55" | MSL | ? | ? | 272,000 | Tr. | |
| 14-1980 | Willamette River at Wilsonville..... | 45°17'31" | 122°46'05" | MSL | 56.60 | 03:30 | 21,000 | .14 | |
| 14-2010 | Pudding River near Mount Angel..... | 45°03'47" | 122°49'45" | 119.78 | 6.84 | 03:30 | 620 | .10 | On axis of buried syncline. |
| 14-3232 | Tennille Creek near Lakeside..... | 43°34'40" | 124°11'30" | MSL | 9.55 | 03:30 | 350 | .02 | Tsunami crest arrived 4 1/2 hr after seiche. |
| Pennsylvania | | | | | | | | | |
| [Only 2 of 102 analog-recorder installations in Pennsylvania recorded the quake] | | | | | | | | | |
| 1-5520 | Loyalsock Creek at Loyalsock..... | 41°19'25" | 76°54'40" | 585.63 | 4.57 | 04:10 | 1,400 | 0.04 | On axis of anticline. |
| 3-1111.5 | Brush Run near Buffalo..... | 40°11'54" | 80°24'28" | 980 | 2.20 | 03:50 | 7.7 | .05 | |
| Puerto Rico | | | | | | | | | |
| No seiche was recorded at any gaging station. | | | | | | | | | |
| Rhode Island | | | | | | | | | |
| No seiche was recorded at any gaging station. | | | | | | | | | |
| South Carolina | | | | | | | | | |
| 2-1309.1 | Black Creek near Hartsville..... | 34°23'50" | 80°09'00" | ----- | 7.24 | 04:20 | 550 | 0.01 | Near buried southwest border of slate belt. |
| 2-1315 | Lynches River near Bishopville..... | 34°15' | 80°13' | 161 | ----- | 04:15 | 2,000 | .05 | On edge of Tertiary overlap. |
| 2-1360 | Black River at Kingstree..... | 33°39'40" | 79°50'10" | 25.66 | 10.21 | 04:40 | 2,700 | Tr. | |
| 2-1480 | Wateree River near Camden..... | 34°14'40" | 80°39'15" | 119.36 | 18.00 | 04:00 | 19,500 | .04 | On edge of Cretaceous overlap. |
| 2-1545 | North Pacolet River at Fingerville..... | 35°07'15" | 81°59'10" | 715.56 | 4.48 | 04:25 | 500 | .08 | |
| 2-1615 | Broad River at Richtex..... | 34°11'06" | 81°11'48" | 184.84 | 10.00 | 03:50 | 34,500 | .08 | Seiche lasted about 60 min. |
| 2-1705 | Lakes Marion-Moultrie diversion canal near Pineville. Auxiliary..... | 33°23'15" | 80°08'25" | MSL | 75.85 | 04:10 | 26,000 | .12 | Seiche lasted about 30 min. |
| | | 33°23' | 80°08' | 60.00 | 0.96 | 04:30 | 26,000 | .00/.02 | Bubble gage? |
| South Dakota | | | | | | | | | |
| 6-4040 | Battle Creek near Keystone..... | 43°52'18" | 103°20'08" | 3,790 | 0.88 | 03:30 | 3 | Tr. | A residual 0.005-ft rise in stage; on south edge of Williston basin. |
| 6-4100 | Castle Creek below Deerfield Dam..... | 44°01'50" | 103°46'35" | 5,805 | 1.24 | 04:15 | 2 | 0.03 | Do. |
| 6-4675 | Missouri River at Yankton..... | 42°52' | 97°24' | 1,159.68 | 1.15 | 04:00 | 24,500 | .14 | May be due to reflection from Sioux uplift. |
| 6-4730 | James River at Ashton..... | 45°00'02" | 98°28'57" | 1,244.4 | 4.58 | 03:30 | 20 | .01 | On southeast edge of Williston basin. |
| 6-4760 | James River at Huron..... | 44°21'55" | 98°11'45" | 1,223.44 | 9.04 | 03:30 | 20 | .03 | Do. |
| 6-4795 | Big Sioux River at Watertown..... | 44°56'30" | 97°08'50" | 1,710 | 5.68 | 04:15 | 3 | .04 | Do. |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|-------------------------|---|-----------|-----------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| Tennessee | | | | | | | | | |
| 3-4250 | Cumberland River at Carthage..... | 36°14'42" | 85°57'15" | 456.33 | 18.60 | 04:15 | 41,300 | 0.36 | Seiche lasted about 30 min. On Cincinnati arch. |
| 3-4265 | Cumberland River at Rome..... | 36°15'50" | 86°04'10" | 449.43 | 11.75 | 03:40 | | .21 | Do. |
| 3-4280 | Cumberland River below Old Hickory..... | 36°15'39" | 86°40'30" | 399.55 | 19.60 | 04:10 | 37,400 | .42 | On northwest side of Nashville dome. |
| 3-4280 | West Fork Stones River near Murfreesboro..... | 35°49'20" | 86°25'03" | 569.51 | 3.35 | 04:00 | 400 | .05 | On crest of Nashville dome. |
| 3-4670 | Lick Creek at Mohawk..... | 36°12'09" | 83°02'53" | 1,072.17 | 11.57 | 03:45 | 1,110 | .03 | In Bays Mountain syncline. On a thrust fault. |
| 3-4910 | Big Creek near Rogersville..... | 36°25'34" | 82°57'07" | 1,131.67 | 2.76 | 04:00 | 138 | .01 | Bubble gage; poor record. |
| 3-4955 | Holston River near Knoxville..... | 36°00'56" | 83°49'54" | 818.06 | 2.23 | 03:50 | 1,260 | ? | Between two thrust faults. |
| 3-5350 | Bullrun Creek near Halls Crossroads..... | 36°06'52" | 83°59'16" | 858.51 | 3.60 | 04:00 | 210 | .03 | Between two thrust faults. |
| 3-5359.1 | Clinch River at Melton Hill Dam (head water)..... | 35°53'04" | 84°18'13" | MSL | 793.20 | 04:00 | 54,800 | .13 | Seiche lasted about 160 min. On a thrust fault. |
| 3-5380 | Whiteoak Creek at Whiteoak Dam..... | 35°53'58" | 84°19'34" | 756.56 | 6.20 | 04:00 | 37 | .06 | Do. |
| 3-5382.25 | Poplar Creek near Oak Ridge..... | 35°59'55" | 84°20'23" | 750.59 | 6.90 | 04:00 | 416 | .04 | On a thrust fault. |
| 3-5382.75 | Bear Creek near Oak Ridge..... | 35°58'50" | 84°21'48" | 755.66 | 1.75 | 03:40 | 35 | .02 | Do. |
| 3-5396 | Daddys Creek near Hobbetsburg..... | 35°59'53" | 84°49'24" | 1,450.45 | 5.35 | 03:45 | 853 | .07 | Between two thrust faults. |
| 3-5660 | Hiwassee River at Charleston..... | 35°17'16" | 84°45'07" | 681.54 | 16.00 | 04:00 | 17,800 | .08 | Do. |
| 3-5675 | South Chickamauga Creek near Chickamauga..... | 35°00'50" | 85°12'27" | 663.41 | 12.25 | 04:00 | 5,820 | .15 | On an anticline between two thrust faults. |
| 3-5710 | Sequatchie River near Whitwell..... | 35°12'22" | 86°29'48" | 644.72 | 12.00 | 04:25 | 4,110 | .11 | Between a thrust fault and an anticline. |
| 3-5845 | Elk River near Prospect..... | 35°01'39" | 86°56'52" | 579.64 | 17.20 | 04:00 | 13,700 | .11 | |
| 3-5884 | Chisholm Creek at Westpoint..... | 35°08'04" | 87°31'45" | 603.29 | 3.08 | 03:55 | 134 | .04 | |
| 3-5935 | Tennessee River at Savannah..... | 35°13'29" | 88°15'36" | 374.82 | | 04:20 | 170,000 | .04 | On edge of cretaceous overlap. |
| 3-5995 | Duck River at Columbia..... | 35°37'05" | 87°01'56" | 549.80 | 14.30 | 04:15 | 7,460 | .14 | |
| 3-6055.5 | Trace Creek near Denver..... | 36°03'28" | 87°53'54" | 391.39 | 1.87 | 03:50 | 54 | .04 | |
| 3-6065 | Big Sandy River at Brueton..... | 36°02'19" | 88°13'42" | 385.14 | 4.38 | 04:15 | 216 | .13 | Near edge of cretaceous overlap. |
| TVA Stations | | | | | | | | | |
| ----- | Tennessee River at Chattanooga (Walnut Street)..... | 35° | 85° | 621.12 | 17.69 | 04:00 | 150,000 | .09 | Between two thrust faults. |
| ----- | Emory River at Harriman..... | 35° | 84° | MSL | 736.50 | 04:00 | 5,000 | .25 | Seiche lasted about 60 min. |
| ----- | Holston River near Morristown..... | 36° | 83° | MSL | 1,050.80 | 04:30 | 940,000 | .10 | |
| ----- | Tennessee River at Kelleys Ferry..... | ----- | ----- | MSL | 633.07 | 04:00 | 150,000 | .12/0.00 | Bubble gage. |
| ----- | Tennessee River at Dougherty's Ferry..... | ----- | ----- | MSL | ? | 04:00 | 450,000 | .14 | |
| ----- | Indian Creek at Cerro Gordo..... | 35° | 88° | 390.0 | 4.48 | 04:00 | 860 | .04 | |
| ----- | Tennessee River at Kingston..... | 35° | 84° | MSL | 736.20 | 04:15 | 800,000 | .04 | |
| ----- | Tennessee River at Clifton..... | 35° | 87° | MSL | 369.10 | 04:45 | 3,400,000 | .07 | |
| ----- | Cherokee Dam headwater..... | ----- | ----- | MSL | 1,050.74 | ----- | 940,000 | Tr. | |
| ----- | Norris Dam headwater..... | 36° | 84° | MSL | 1,000.97 | ----- | 1,450,000 | .09 | Seiche lasted about 80 min. |
| Texas | | | | | | | | | |
| 7-2996.7 | Groesbeck Creek near Quanah..... | 34°21'20" | 99°44'25" | 1,425.69 | 5.21 | 04:15 | 6.4 | 0.02 | On south side of basin. |
| 7-3121 | Wichita River near Mabelle..... | 33°45'35" | 99°08'35" | 1,062.72 | 3.79 | 04:00 | 144 | .04 | |
| 7-3150 | Little Wichita River near Henrietta..... | 33°50'00" | 98°12'30" | 831.57 | 6.19 | 03:55 | | .08 | Seiche lasted 30 min or more. On Ouachita tectonic belt. |
| 7-3315 | Lake Texoma near Denison..... | 33°49'05" | 96°34'20" | MSL | 604.13 | ----- | 1,777,200 | .00/0.04 | Bubble gage. |
| 7-3326 | Bois d'Arc Creek near Randolph..... | 33°28'30" | 96°21'50" | 564.38 | 2.25 | 04:20 | | .03 | On Ouachita tectonic belt. |
| 7-3355 | Red River at Arthur City..... | 33°52'30" | 95°30'10" | 380.07 | 8.84 | 03:55 | 3,240 | .04 | On basin in East Texas embayment. |
| 7-3368 | Pecan Bayou near Clarksville..... | 33°41'07" | 94°59'41" | 365.00 | 3.68 | 03:55 | 18 | .08 | |
| 7-3426 | South Sulphur River near Cooper..... | 32°21' | 95°36' | 374.91 | 1.09 | 04:00 | 4.5 | .02 | A residual 0.005-ft drop in stage. |
| 7-3436 | Whiteoak Creek near Talco..... | 33°19' | 96°05' | 286.45 | 3.31 | 04:00 | 12 | .02 | A residual 0.01-ft drop in stage. |
| 7-3450 | Boggy Creek near Daingerfield..... | 33°02'05" | 94°47'10" | 258.41 | 4.92 | 04:00 | 25 | .03 | |
| 7-3460.5 | Little Cypress Creek near Ore City..... | 32°40'21" | 94°45'03" | 232.67 | 4.53 | 04:00 | 84 | .05 | Seiche lasted about 45 min. On westward extension of Rodessa fault zone. |
| 7-3460.7 | Little Cypress Creek near Jefferson..... | 32°45' | 94°30' | 174.60 | 5.59 | 04:00 | 197 | .03 | On Rodessa fault zone. |
| 8-0173 | South Fork Sabine River near Quinlan..... | 32°53'52" | 96°15'11" | 461.40 | 3.27 | 04:00 | 1 | .00/0.01 | Float gage. On Ouachita tectonic belt. |
| 8-0193 | Lake Winnsboro near Winnsboro..... | 32°53'10" | 95°20'40" | MSL | 410.95 | 04:00 | 2,960 | .00/0.03 | Bubble gage. On north end of East Texas embayment. |
| 8-0195 | Big Sandy Creek near Big Sandy..... | 32°36'12" | 95°05'32" | 278.38 | 4.92 | 04:00 | 78 | No seiche | A lasting 0.005-ft rise in stage. Bubble gage. On east edge of East Texas embayment. |
| 8-0207 | Rabbit Creek at Kilgore..... | 32°23'17" | 94°54'11" | 299.80 | 2.90 | 04:00 | 20 | .03 | |
| 8-0222 | Murvaul Lake near Gary..... | 32°02'04" | 94°25'15" | MSL | 264.04 | 04:00 | 40,940 | .10 | Seiche lasted about 30 min. with 0.04 ft of motion. Between two normal faults. |
| 8-0223 | Murvaul Bayou near Gary..... | 32°01'54" | 94°22'31" | 217.82 | 3.10 | 04:00 | 7.5 | .03 | On a normal fault. |
| 8-0285 | Sabine River near Bon Weir..... | 30°45'00" | 93°38'30" | 46.42 | 5.40 | 04:00 | 3,950 | .19 | Seiche lasted about 30 min. On a normal fault. |
| 8-0305 | Sabine River near Ruliff..... | 30°18'10" | 93°44'40" | 4.08 | 11.85 | 03:50 | 6,920 | .67 | Seiche lasted about 50 min. Southeast side of East Texas embayment. |
| 8-0320 | Neches River near Neches..... | 31°53'32" | 95°25'50" | 264.06 | 6.30 | 04:00 | 204 | .11 | |
| 8-0385 | Angelina River near Zavalla..... | 31°12'41" | 94°17'40" | 104.48 | 9.89 | 04:00 | 2,010 | .63 | Seiche lasted about 50 min. |
| 8-0410 | Neches River near Evadale..... | 30°21'22" | 94°05'36" | 8.25 | 12.04 | 04:00 | 6,200 | .31 | Seiche lasted about 60 min. |
| 8-0680 | West Fork San Jacinto River near Conroe..... | 30°14'41" | 95°27'28" | 95.03 | 6.42 | 04:00 | 208 | .27 | Seiche lasted about 40 min. |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|--------------------------------|---|-----------|-------------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| Texas—Continued | | | | | | | | | |
| 8-0720 | Lake Houston near Sheldon | 29°54'58" | 95°08'28" | -0.70 | 44.61 | 03:45 | 59,600 | 0.13 | Seiche lasted about 120 min. |
| 8-0760 | Greens Bayou near Houston | 29°55'05" | 95°18'24" | - .66 | 49.71 | 04:00 | 6.4 | .07 | Seiche lasted about 30 min. |
| 8-0815 | Salt Croton Creek near Aspermont | 33°24'05" | 100°24'30" | 1,668 | 1.30 | 03:40 | .5 | .02 | |
| 8-0848 | California Creek near Stamford | 32°55'50" | 99°38'30" | | 6.21 | 04:00 | .7 | .02 | |
| 8-0873 | Clear Fork of Brazos River at Eliasville | 32°57'30" | 98°46'10" | 1,027.77 | 7.53 | 03:45 | 13 | .02/.13 | Bubble gage. |
| 8-0883 | Oak Creek near Graham | 33°12'40" | 98°37'05" | | .76 | 04:10 | 0 | .03 | |
| 8-0884 | Lake Graham near Graham | 33°08'05" | 98°36'55" | MSL | 1,072.99 | 03:50 | 48,640 | .08 | Seiche lasted about 50 min. Bubble gage. |
| 8-0953 | Middle Bosque River near McGregor | 31°30'33" | 97°21'58" | 530.51 | 2.90 | 04:00 | 27 | .04 | On Ouachita tectonic belt. |
| 8-0954 | Hog Creek near Crawford | 31°33'20" | 97°21'22" | 560.54 | 2.26 | 04:00 | 11 | .04 | Do. |
| 8-0956 | Bosque River near Waco | 31°36'04" | 97°11'38" | 365.44 | 4.04 | 04:00 | 149 | .04 | Do. |
| 8-0968 | Cow Bayou Subwatershed 4 near Bruceville | 31°20' | 97°16' | 574.46 | 10.01 | 04:00 | 58.3 | .008 | Do. |
| 8-1020 | Belton Reservoir near Belton | 31°07' | 98°28' | MSL | 589.28 | 04:00 | 213,700 | .06 | Seiche lasted about 45 min. Near a normal fault. |
| 8-1065 | Little River at Cameron | 30°50' | 96°57' | 281.89 | 7.72 | 04:00 | 1,400 | .00/03 | Float gage, near edge of tertiary overlap. |
| 8-1087 | Middle Yegua River near Dime Box | 30°20'20" | 96°54'15" | 295.4 | 1.26 | 04:00 | 1.9 | .03 | |
| 8-1100 | Yegua Creek near Somerville | 30°19'18" | 96°30'27" | 199.21 | 2.53 | 04:00 | 26 | .07 | Seiche lasted about 20 min. |
| 8-1103 | Lake Mexia near Mexia | 31°38'45" | 96°34'39" | MSL | 426.52 | 04:00 | 7,000 | .14 | Seiche lasted about 20 min. on Mexia-Talco fault zone. |
| 8-1105 | Navasota River near Easterly | 31°10'10" | 96°17'55" | 276.46 | 1.56 | 04:00 | 12 | .02 | |
| 8-1115 | Brazos River near Hempstead | 30°07'34" | 96°11'05" | 117.90 | 4.14 | 04:00 | 2,000 | .00/.12 | Bubble gage. |
| 8-1175 | San Bernard River near Bowling | 29°18'47" | 95°53'36" | 30.80 | 4.18 | 04:00 | 62 | .005/.035 | |
| 8-1180 | Lake J. B. Thomas near Vincent | 32°35'09" | 101°12'18" | MSL | 2,249.44 | 04:00 | 148,800 | .05 | |
| 8-1190 | Bluff Creek near Ira | 32°35'29" | 101°03'05" | 2,177.95 | 3.18 | 04:00 | .1 | No seiche | Slight shift downward during 20 min. |
| 8-1236 | Champion Creek Reservoir near Colorado City | 32°16'55" | 100°51'30" | MSL | 2,055.62 | 04:00 | 18,290 | .06 | Seiche lasted about 60 min. |
| 8-1270 | Elm Creek at Ballinger | 31°45'00" | 99°58'50" | 1,617.72 | 3.90 | 04:00 | 1.0 | .04 | Seiche lasted about 20 min. |
| 8-1280 | South Concho River at Christoval | 31°13' | 100°30' | 2,010.22 | 1.85 | 04:00 | 8.3 | .015/.035 | A residual 0.01-ft drop in stage. |
| 8-1365 | Concho River near Paint Rock | 31°31' | 99°55' | 1,574.43 | 12.63 | 04:00 | 1.9 | .05 | Seiche lasted about 120 min. |
| 8-1400 | Deep Creek subwatershed 8 near Mercury | 31°23'05" | 99°08'30" | 1,377.13 | 8.99 | 03:55 | 214 | .08 | A residual 0.002-ft drop in stage near a normal fault. |
| 8-1435 | Pecan Bayou at Bronwood | 31°43'54" | 98°58'25" | 1,318.58 | .52 | 04:00 | .9 | .04 | Seiche lasted about 90 min. |
| 8-1535 | Federnales River at Johnson City | 30°18' | 98°24' | 1,096.70 | 2.84 | 04:00 | 58 | .005/.000 | On north side of Llano uplift |
| 8-1610 | Colorado River at Columbus | 29°42'20" | 96°32'05" | 155.52 | 1.61 | 04:00 | 238 | .04/.06 | Float gage. On southeast side of Llano uplift. |
| 8-1676 | Rebecca Creek near Spring Branch | 29°55'08" | 98°22'09" | 985.55 | 2.14 | | 3.8 | .04 | Seiche lasted about 35 min. On northeast extension of fault. |
| 8-1713 | Blanco River near Kyle | 29°58'42" | 97°54'30" | 620.12 | 4.30 | | 20 | .05 | On Ouachita tectonic belt. |
| 8-1758 | Guadalupe River at Cuero | 29°03'57" | 97°19'16" | 128.64 | 5.16 | | 710 | .00/.39 | Seiche lasted about 30 min. On Balcones fault zone. |
| 8-1780 | San Antonio River at San Antonio | 29°24'35" | 98°29'40" | 612.26 | 1.07 | | 16 | .03 | Bubble gage. |
| 8-1790 | Medina River near Pipe Creek | 29°40' | 98°59' | 1,067.37 | 4.41 | | 66 | .03 | Seiche lasted about 30 min. Near a normal fault and on edge of Tertiary overlap. |
| 8-1824 | Calaveras Creek subwatershed 6 near Elmendorf | 29°22'53" | 98°17'34" | 516.06 | 14.85 | 04:00 | 49.6 | .018/.000 | On Ouachita tectonic belt. Water-level rise lasted about 15 min. Float gage. Near a normal fault. |
| 8-1825 | Calaveras Creek near Elmendorf | 29°15'38" | 98°17'34" | 406.45 | 4.77 | 04:00 | 1.7 | No seiche | A 0.005-ft drop in stage. |
| 8-1839 | Cibola Creek near Boerne | 29°46'25" | 98°41'52" | 1,339.61 | 2.37 | | 5.6 | .02 | On Ouachita tectonic belt. |
| 8-1875 | Escondido Creek at Kenedy | 28°49'11" | 97°51'32" | 246.40 | 8.99 | 04:00 | 1.6 | .02 | Seiche lasted about 40 min. |
| 8-1879 | Escondido Creek subwatershed 11 near Kenedy | 28°51'39" | 97°50'39" | 288.12 | 15.58 | 03:55 | 168 | .018 | Seiche lasted about 10 min. |
| 8-1893 | Media Creek near Beeville | 28°28'58" | 97°39'23" | 163.00 | 5.10 | | No flow | .02 | |
| 8-1895 | Mission River at Refugio | 28°17'30" | 97°16'44" | 1.00 | 2.07 | | 4.5 | .05 | |
| 8-2027 | Seco Creek at Cook Ranch near D'Hanis | 29°21'43" | 99°17'05" | 900.88 | 4.37 | | No flow | .03 | |
| 8-2055 | Frio River at Derby | 28°44'10" | 99°08'45" | 449.11 | .49 | | do. | .005 | |
| 8-2070 | Frio River at Callham | 28°29'30" | 98°20'45" | 153.47 | 2.84 | | 8.6 | .005 | Seiche lasted about 15 min. |
| 8-2110 | Nueces River at Mathis | 28°02'17" | 97°51'36" | 27.53 | 2.18 | | 7.3 | .00/.08 | On a normal fault. Bubble gage. |
| 8-4275 | San Solomon Springs at Toyahvale Reservoir in Bailey County | 30°56'34" | 103°47'102" | 3,311.02 | .96 | 04:00 | 30 | .07 | Seiche lasted about 30 min. |
| | | | | | | 04:10 | 16 | .5 | Miller and Reddell (1964, p. 861). |
| Utah | | | | | | | | | |
| 10-0201 | Bear River above reservoir near Woodruff | 41°26'05" | 111°01'00" | 6,455 | | 04:00 | 50 | Tr. | On north-south fault. |
| 10-0210 | Woodruff Creek near Woodruff | 41°29' | 111°16' | 6,600 | | 04:00 | 8 | Do | |
| 10-1345 | East Canyon Creek near Morgan | 40°55'20" | 111°38'20" | 5,460 | | | 14 | Do | |
| 10-1376 | Southfork Ogden River at Huntsville | 41°14'50" | 111°45'45" | 4,910 | | | 38 | Do | On a buried fault. |
| 10-1376.8 | North Fork Ogden River near Eden | 41°23'20" | 111°54'50" | 5,750 | | | 4 | Do | |
| 10-1377 | North Fork Ogden River near Huntsville | 41°17'40" | 111°49'40" | 4,903.81 | 0.55 | 04:40 | 2 | .04 | |
| 10-1705 | Surplus Canal at Salt Lake City | 40°43'40" | 111°55'35" | 4,219.02 | 1.00 | 04:10 | 70 | .06 | |
| 10-1940 | Sevier River above Clear Creek near Sevier | 38°34'20" | 112°15'25" | 5,560 | | | 90 | Tr. | Near a normal fault. |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|---|---|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| UNITED STATES—Continued | | | | | | | | | |
| Vermont | | | | | | | | | |
| 4-2835 | East Barre Detention Reservoir at East Barre. | 44°09'20" | 72°26'40" | MSL | 1,130.67 | 04:00 | 8,600 | 0.06 | Near axis of north-south syncline. |
| 4-2850 | Wrightsville Detention Reservoir at Wrightsville. | 44°18'35" | 72°34'30" | MSL | 618.72 | 04:00 | 29,000 | .23 | |
| Virginia | | | | | | | | | |
| No seiche was recorded at any gaging station. | | | | | | | | | |
| Washington | | | | | | | | | |
| 12-1555 | Snohomish River at Snohomish..... | 47°54'45" | 122°06'30" | -9.86 | 3.49 | 03:45 | <10,000 | <0.45 | Seiche superimposed on tidal curve. Seiche lasted about 30 min. On small structural complex. |
| 12-3971 | Outlet Creek near Metaline Falls..... | 48°50'45" | 117°17'15" | 2,550 | 9.18 | 04:15 | 17 | No seiche | Temporary drop in stage of 0.005 ft. |
| 12-3980.9 | Pend Oreille River at Metaline Falls. | 48°51'55" | 117°22'20" | ----- | 11.80 | 03:45 | ? | .16 | On a fault. |
| 12-4087 | Mill Creek at mouth near Colville.... | 48°34'25" | 117°56'40" | 1,540 | 1.36 | 03:50 | 27 | .03 | |
| 12-4360 | Franklin D. Roosevelt Lake at Grand Coulee Dam. | 47°57'20" | 118°59'10" | MSL | 1,253.30 | 03:45 | 6,800,000 | 1.04 | Seiche lasted at least 2 hr and perhaps about 12 hr on Colville batholith. |
| 12-4390 | Osoyoos Lake near Oroville..... | 48°59'15" | 119°27'15" | MSL | 911.15 | 04:00 | ----- | Tr. | Near north edge of Columbia River Basalt. |
| 12-4395 | Okanogan River at Oroville..... | 48°55'55" | 119°25'05" | 899.77 | 3.55 | 03:45 | 575 | Tr. | Do. |
| 12-4440 | Whitestone Lake near Tonasket..... | 48°47'15" | 119°27'50" | ----- | 4.35 | 03:30 | ----- | .13 | Do. A 0.03-ft rise in stage. |
| 12-4500 | Alta Lake near Pateras..... | 48°01'30" | 119°56'30" | 1,175 | 8.03 | 04:00 | ----- | .13 | Seiche was recorded during 60 min. |
| 12-4545 | Wenatchee Lake near Plain..... | 47°49'50" | 120°46'30" | MSL | 1,870.10 | 04:10 | ----- | No seiche | Slight temporary rise in water level on axis of anticline. |
| 12-4670 | Crab Creek near Moses Lake..... | 47°11'25" | 119°16'00" | 1,070.39 | 1.40 | 03:00 | 6 | No seiche | A lasting 0.005-ft rise in stage. In Quincy basin. |
| 12-4690 | Blue Lake near Coulee City..... | 47°34'25" | 119°25'15" | MSL | 1,063.27 | 03:50 | ----- | .04 | On axis of syncline. |
| 12-4695 | Lenore Lake near Soaplake..... | 47°31' | 119°30' | MSL | 1,078.20 | 04:00 | ----- | Tr. | Pen trace became darker. On axis of syncline. |
| <i>U.S. Corps of Engineers</i> | | | | | | | | | |
| ----- | McNary Reservoir at Port Kelly..... | 46° | 118° | MSL | 337.38 | 03:45 | ----- | .69 | Bubble gage. |
| ----- | McNary Reservoir at Wallula Junction. | 46° | 118° | MSL | 337.39 | 04:00 | ----- | .15 | Stevens A-35 recorder. |
| ----- | McNary Reservoir at Union Pacific RR bridge near Kennewick. | 46° | 119° | MSL | 337.26 | 03:45 | ----- | .08 | Do. |
| ----- | McNary Reservoir at Snake River Bridge near Burbank. | 46° | 119° | MSL | 337.30 | 03:45 | ----- | .12 | Do. |
| ----- | McNary Reservoir at Pasco-Kennewick Highway bridge. | 46° | 119° | MSL | 337.40 | 03:45 | ----- | .22 (est.) | Do. |
| ----- | McNary Reservoir at Richland Pumping Plant. | 46° | 119° | MSL | 337.82 | 03:45 | ----- | .10 | Do. |
| ----- | Ice Harbor Reservoir Navigation Lock. | 46° | 119° | MSL | 437.56 | 03:45 | ----- | .20 | Preexisting wind seiches were amplified by seismic waves. |
| ----- | Ice Harbor Reservoir near Page..... | 46° | 119° | MSL | 437.58 | 03:45 | ----- | .30 | Bubble gage. |
| West Virginia | | | | | | | | | |
| No seiche was recorded at any gaging station. | | | | | | | | | |
| Wisconsin | | | | | | | | | |
| 4-0790 | Wolf River at New London..... | 44°23'30" | 88°44'25" | 749.37 | ----- | 03:50 | 710 | 0.01 | On south edge of Precambrian felsic intrusive body. |
| 4-0800 | Little Wolf River at Royalton..... | 44°24'45" | 88°51'55" | 774.00 | 1.28 | 03:50 | 140 | .02 | Do. |
| 5-3360 | St. Croix River at Grantsburg..... | 45°55'25" | 92°38'20" | 848.98 | ----- | 03:40 | 1,300 | .01 | On axis of syncline. |
| 5-4050 | Baraboo River near Baraboo..... | 43°28'55" | 89°38'00" | 788.21 | ----- | 03:50 | 170 | .01 | |
| 5-4240 | East Branch Rock River near Mayville. | 43°31'45" | 88°34'00" | 857.20 | ----- | 04:00 | 50 | .01 | |
| 5-4330 | East Branch Pecatonica River near Blanchardville. | 42°47'10" | 89°51'40" | 796.8 | ----- | 04:00 | 64 | .01 | |
| Wyoming | | | | | | | | | |
| 6-2316 | Middle Popo Agie below the Sinks, near Lander. | 42°45'25" | 108°47'50" | 6,150 | 2.00 | 04:20 | 18 | Tr.? | On west side of Wind River basin. |
| 6-2355 | Little Wind River near Riverton..... | 42°59'51" | 108°22'29" | 4,901.84 | 3.24 | 03:35 | 270 | .01 | |
| 6-2445 | Fivemile Creek above Wyoming Canal near Pavillion. | 43°18'04" | 108°42'04" | 5,495 | 1.95 | 04:00 | 4 | .02 | Do. |
| 6-2765 | Greybull River at Meeteetse..... | 44°09'20" | 108°52'35" | 5,739.42 | ----- | 04:15 | 68 | Tr. | On west side of Big Horn basin. |
| 6-2785 | Shell Creek near Shell..... | 44°34' | 107°42' | 4,367.20 | ----- | 03:30 | 35 | .08 | |
| 6-2803 | South Fork Shoshone River near Valley. | 44°12'30" | 109°33'15" | 6,200 | 2.47 | 04:00 | 59 | .02 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|------------------------------------|---|--------------|---------------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| UNITED STATES—Continued | | | | | | | | | |
| Wyoming—Continued | | | | | | | | | |
| 6-2844 | Shoshone River near Garland..... | 44°44' | 108°36' | 4, 100 | 4. 74 | 04:00 | 660 | 0. 08 | On possible extension of a thrust fault. |
| 6-6377.5 | Rock Creek above Rock Creek Reservoir. | 42°32'59" | 108°46'26" | 8, 330 | 4. 43 | 04:00 | 1 | . 01 | |
| 9-1985 | Pole Creek below Little Half Moon Lake near Pinedale. | 42°53' | 109°43' | 7, 350 | 2. 80 | 04:20 | 11 | . 07 | On buried thrust fault. |
| 9-2105 | Fontenelle Creek near Herschler Ranch, near Fontenelle. | 42°05'45" | 110°25'10" | 6, 950 | 3. 25 | 04:10 | 32 | Tr. | On axis of an anticline. |
| 9-2230 | Hams Fork near Elk Creek Ranger Station. | 42°06'40" | 110°42'40" | 7, 455 | 3. 94 | 03:30 | 23 | . 02 | In area of thrust faults. |
| 13-0110 | Snake River at Moran..... | 43°51' | 110°35' | 6, 727. 84 | ----- | 04:00 | 408 | . 005 | Lake Hebgen earthquake was also recorded by this gage. Near end of a thrust fault. |
| AUSTRALIA | | | | | | | | | |
| Australia Capital Territory | | | | | | | | | |
| ----- | O'Conner Reservoir at Canberra.... | 35° S. | 149° E. | ----- | ----- | 04:45 | 21 | Tr. | Previous earthquakes in Kurile Islands (Oct. 13, 1963), Banda Sea (Nov. 4, 1963), and New Hebrides were recorded on this reservoir (Robert Underwood, written commun., Sept. 20, 1965). |
| New South Wales | | | | | | | | | |
| ----- | Tantangara Reservoir..... | 35°47'53" S. | 148°39'44" E. | MSL | 3, 971. 51 | 04:40 | 25, 680 | 0. 02 | Recorder is near dam. |
| Northern Territory | | | | | | | | | |
| 113A..... | Victoria River..... | 16°22' S. | 131°06' E. | ----- | ----- | 04:45 | ----- | 0. 00/. 02 | Servomanometer recorder. |
| Victoria | | | | | | | | | |
| M17..... | Melicke Munjie River..... | 37°14'40" S. | 148°08'30" E. | 2, 100 | ----- | 04:00 | ----- | 0. 02 | |
| CANADA | | | | | | | | | |
| Alberta | | | | | | | | | |
| 5-0130 | Waterton River near Waterton Park.. | 49°07' | 113°50' | ----- | 0. 84 | 04:00 | ----- | 0. 03 | |
| 6-1345 | Milk River at Milk River..... | 49°09' | 112°05' | ----- | 2. 45 | 03:50 | ----- | . 02 | |
| 6-1355 | Sage Creek at "Q" Ranch near Wild Horse. | 49°08' | 110°13' | ----- | 2. 25 | 04:00 | ----- | . 09 | |
| ----- | Athabasca River near Hinton..... | 53°25' | 117°35' | ----- | 7. 02 | 03:55 | ----- | . 05 | |
| ----- | Belly-St. Mary Diversion Canal..... | 49°20' | 113°32' | ----- | 3. 55 | 05:00 | ----- | . 01 | |
| ----- | Bow River at Calgary..... | 51°03' | 114°03' | ----- | ----- | 04:00 | ----- | . 03 | |
| ----- | Clearwater River at Draper..... | 56°41' | 111°15' | ----- | ----- | 03:45 | ----- | . 00/. 05 | A sudden 0.13-ft rise in stage. Bubble gage. |
| ----- | Clearwater River near Rocky Mountain House. | 52°21' | 114°56' | ----- | 3. 84 | ----- | ----- | . 07 | |
| ----- | Elbow River at Bragg Creek..... | 50°57' | 114°34' | ----- | 5. 40 | 03:45 | ----- | . 03 | |
| ----- | Highwood River near Aldersyde..... | 50°42' | 113°51' | ----- | 4. 61 | ----- | ----- | . 01 | |
| ----- | Lesser Slave River at Highway 2..... | 55°18' | 114°35' | ----- | 86. 60 | 04:00 | ----- | No seiche | A lasting 0.02-ft rise in stage. Bubble gage. |
| ----- | Little Smokey River near Guy..... | 55°27' | 117°10' | ----- | 9. 73 | 04:20 | ----- | . 03/. 045 | A residual 0.01-ft drop in stage. Bubble gage. |
| ----- | Oldman River at Lethbridge..... | 49°42' | 112°52' | ----- | 2. 32 | 04:20 | ----- | . 02/. 04 | Bubble gage. |
| ----- | Peace River at Fort Vermilion..... | 58°24' | 116°00' | ----- | 57. 95 | 03:45 | ----- | . 08/. 10 | Do. |
| ----- | Peace River at Peace Point..... | 59°07' | 112°26' | ----- | 58. 79 | 04:10 | ----- | . 03 | Do. |
| ----- | Peace River at Peace River..... | 56°15' | 117°19' | ----- | 21. 33 | 04:30 | ----- | . 025/. 05 | Do. |
| ----- | Prairie Creek near Rocky Mountain House. | 52°16' | 114°56' | ----- | 3. 06 | 03:00 | ----- | . 02/. 00 | |
| ----- | Red Deer River at Drumheller..... | 51°28' | 112°42' | ----- | ----- | 04:15 | ----- | . 31 | |
| ----- | Sheep River at Aldersyde..... | 50°43' | 113°53' | ----- | 5. 00 | 04:00 | ----- | . 00/. 04 | Bubble gage. |
| ----- | Slave River at Fitzgerald..... | 59°52' | 111°35' | ----- | 687. 37 | 04:10 | ----- | . 00/. 10 | Do. |
| ----- | South Saskatchewan River at Medicine Hat. | 50°03' | 110°41' | ----- | 7. 35 | 05:00 | ----- | . 00/. 07 | Do. |
| ----- | Stimson Creek near Pekisko..... | 50°26' | 114°10' | ----- | ----- | 03:45 | ----- | . 03 | |
| ----- | Twin Creek near Seebe..... | 50°58' | 115°10' | ----- | ----- | ----- | ----- | . 025 | |
| ----- | Middle Creek near Alberta Boundary. | 49°28' | 110°03' | ----- | 3. 10 | ----- | ----- | . 01 | |
| 6-1340 | North Fork Milk River near International Boundary. | 49°01'20" | 112°58'20" | 4, 120 | 3. 45 | 03:20 | 8. 3 | . 03 | Stage rose 0.03 ft after seiche was recorded. |
| 6-1330 | Milk River at Western Crossing of International Boundary. | 49°00' | 112°33' | 3, 820 | 3. 96 | 03:20 | 20 | . 01 | |
| 6-1360 | Sage Creek at International Boundary. | 49°00'10" | 110°12'30" | 2, 800 | 2. 63 | 04:00 | 6 | . 08 | |
| 5-0205 | Saint Mary River near International Boundary. | 49°00' | 113°18'50" | 4, 120 | 5. 06 | 03:50 | 69 | . 02 | |

TABLE 3.—Seismic effects from the Alaska earthquake at surface-water gages—Continued

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|------------------------------|---|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|--|
| CANADA—Continued | | | | | | | | | |
| British Columbia | | | | | | | | | |
| | Prince Rupert | 54°19' | 130°20' | | | 03:45 | | 0.25 | Data from Wigen and White (1964). |
| | Bella Bella | 52°10' | 128°08' | | | 03:45 | | .35 | Do. |
| | Tasu | 52°45' | 132°01' | | | 03:45 | | 1.10 | Do. |
| | Victoria | 48°25' | 123°24' | | | 03:45 | | .15 | Do. |
| | Point Atkinson | 49°20' | 123°15' | | | 04:00 | | .40 | Do. |
| | Vancouver | 49°17' | 123°07' | | | 03:45 | | .40 | Do. |
| | Port Moody | 49°17' | 122°52' | | | | | .35 | Do. |
| | Balenas Island | 49°20' | 124°09' | | | | | .40 | Do. |
| | Frazer River at New Westminster | 49°11'52" | 122°54'42" | | | 03:45 | | .15 | Do. |
| | Link Lake near Ocean Falls | 52°21' | 127°41' | | | | | .26 | Do. |
| 8BB-1 | Taku River at Tulsequah | 53°38'20" | 133°32'25" | | 3.30 | 03:50 | | .05 | 20 min after seiche, water level began rise of 0.34 ft in 2 hr. |
| 8EG-14 | Rainbow Lake near Prince Rupert | 54°11'36" | 130°04'50" | | 2.35 | 02:40 | | .20 | |
| 8FA-7 | Owikeno Lake near Wadhams | 51°40'40" | 127°10'30" | | 4.66 | 03:40 | | .12 | |
| 8KB-1 | Fraser River at Shelley | 54°00'40" | 122°37'00" | 1,859.67 | 10.30 | 04:00 | | .05 | Seiche lasted about 4 hr. Trace of upward shift. |
| 8LA-10 | Mahood Lake near Clearwater Station | 51°56'18" | 120°14'28" | | 3.05 | 04:00 | | .10 | Wind seiche amplified by seismic seiche. |
| 8LA-12 | Clearwater Lake near Clearwater Station | 52°07'55" | 120°11'10" | | 4.40 | 03:45 | | .15 | |
| 8LE-53 | Shuswap Lake at Sicamous | 50°51'05" | 119°00'43" | 1,131.93 | 1.90 | 04:20 | | .14 | |
| 8ME-17 | Seton Lake near Shalath | 50°43'40" | 122°14'00" | 0.36 | 774.18 | 04:00 | | .55/.00 | Seiche lasted about 10 hr. Maximum observed seiche was about 3 ft. |
| 8MH-16 | Chilliwack River at outlet Chilliwack Lake near Vedder Crossing | 49°05'02" | 121°27'24" | | 1.70 | 03:50 | | .00/.10 | 30 min required for water level to recover, but did not rise to previous level. |
| 8MH-52 | Pitt Lake near Alvin | 49°28'10" | 122°30'45" | | 5.50 | 03:45 | | .46 | Pitt Lake is tidal. |
| 8MH-62 | Pitt Lake near outlet near Pitt Meadows | 49°21'27" | 122°34'38" | | 6.60 | 03:50 | | .22 | Do. |
| 8NE-45 | Upper Arrow Lake at Nakusp | 50°14'12" | 117°48'07" | 1,374.07 | 1.70 | 04:00 | | 1.25 | Seiche lasted about 12 hr. Lake highly resonant. Exponential decay well defined. |
| 8NH-64 | Kootenay Lake at Queen's Bay | 49°39'16" | 116°55'47" | 0.38 | 1,739.20 | 03:45 | | .06 | |
| 8NH-67 | Kootenay Lake at Kuskanook | 49°17'56" | 116°39'31" | 1,735.20 | 4.62 | 03:45 | | .10 | |
| Manitoba | | | | | | | | | |
| | Nelson River at Cross Lake | 54°36' | 97°47' | | | 03:35 | | 0.29 | |
| | Lake Winnipeg at Pine Dock | 51°38'30" | 96°47'45" | | | 03:50 | | .05 | |
| | Lake Manitoba at the Narrows | 51°05'00" | 98°47'45" | | | 04:10 | | .03 | |
| | Deloraine Reservoir near Deloraine | 49°08'50" | 100°24'40" | | | 03:50 | | .44 | P. W. Strilaeff (written commun., 1964). |
| Northwest Territories | | | | | | | | | |
| | Cambridge Bay | 69°07' | 105°04' | | | | | 0.30 | Seiche lasted 15 min. (Wigen and White, 1964). |
| | Talston River at outlet Tsu Lake | 60°39' | 111°57' | | 85.20 | 04:00 | | .00/.15 | Bubble gage. |
| | Willowlake River near the mouth | 62°39' | 122°55' | | 62.20 | 03:50 | | .00/.03 | Water level rose 0.01 ft. Bubble gage. |
| | Great Bear Lake at Port Radium | 66°04' | 117°52' | | 389.53 | 03:50 | | .00/.22 | Bubble gage. |
| | Lockhart River at outlet Artillery Lake | 62°53' | 108°28' | | 96.08 | 03:40 | | .055/.035 | Do. |
| | Hay River above Hay River | 60°45' | 115°21' | | 65.73 | 03:50 | | .00/.09 | Do. |
| | Mackenzie River at Wrigley | 63°16' | 123°38' | | 70.94 | 03:40 | | .00/.10 | Do. |
| Ontario | | | | | | | | | |
| | English River at Sioux Lookout | 50°04'15" | 91°56'40" | | | 03:50 | | 0.14 | Two maximums of equal size about 12 min. apart. |
| | Lake of the Woods at Clearwater Bay | 49°43'06" | 94°48'10" | | | 03:45 | | .09/.03 | Bubble gage. |
| | Gull River at Norland | 44°43'55" | 78°49'08" | | 61.47 | 04:00 | | .03 | |
| | Skootamata River at Actinolite | 44°32'39" | 77°19'35" | | 10.90 | 04:00 | | .055 | |
| | Wanapitell-Wanup River | 46°21' | 80°50' | | 708.36 | 04:00 | | .02 | Water level began decline of 0.05 ft after seiche recorded. |
| | Lac la Croix at Campbell's Camps | 48°21'20" | 92°12'50" | | | 03:30 | | .13 | |
| | Mississagi River | 46°54' | 83°14' | | 4.85 | | | .03/.04 | Bubble gage. |
| | French River-Dry Pine Bay | 46°03'01" | 80°34'28" | | 593.12 | 04:00 | | .03 | |
| Saskatchewan | | | | | | | | | |
| | Buffalo Pound Lake at Pumping Station | 50°35' | 105°23' | | 71.85 | 03:30 | | 0.075 | |
| | Fond du Lac River at outlet Black Lake | 59°09' | 105°33' | | 93.16 | 04:00 | | .00/.075 | Bubble gage. |
| | South Saskatchewan River near Lemsford | 51°01' | 109°08' | | 4.24 | 04:20 | | Tr. | |
| | Spruce River below Anglin Lake Reservoir | 53°40' | 106°00' | | 2.88 | 04:00 | | .03 | |

TABLE 3.—*Seismic effects from the Alaska earthquake at surface-water gages—Continued*

| Station number | Station name and location | Latitude | Longitude | Datum of gage (ft) | Stage (ft) | Time | Discharge (cfs) or storage (acre ft) | Seiche double amplitude (ft) | Remarks |
|-------------------------------|--|-----------|------------|--------------------|------------|-------|--------------------------------------|------------------------------|---|
| CANADA—Continued | | | | | | | | | |
| Saskatchewan—Continued | | | | | | | | | |
| 6-1495 | Battle Creek near International Boundary. | 49°00'10" | 109°25'20" | 2,729.8 | 2.22 | 03:50 | 4 | 0.09/.00 | |
| 6-1580 | Frenchman River above Eastend Reservoir near Ravenscrag. | 49°29' | 109°00' | 3,040 | 1.76 | 03:45 | 12 | .19 | |
| 6-1785 | East Poplar River at International Boundary. | 49°00'00" | 105°24'30" | 2,410.92 | 2.65 | 04:00 | 4.5 | .16 | |
| ----- | Long Creek below Boundary Reservoir. | 49°06'43" | 102°59'42" | ----- | ----- | 03:35 | ----- | .30 | P. W. Strlaeff (1964, written commun.). |
| ----- | Weyburn Reservoir near Weyburn... | 49°38'28" | 103°49'24" | ----- | ----- | 03:50 | ----- | .04 | Do. |

The Alaska Earthquake
March 27, 1964:
Effects on the
Hydrologic Regimen

*This volume was published
as separate chapters A-E*

UNITED STATES DEPARTMENT OF THE INTERIOR

STEWART L. UDALL, *Secretary*

GEOLOGICAL SURVEY

William T. Pecora, *Director*



CONTENTS

[Letters designate the separately published chapters]

- (A) Effects of the March 1964 Alaska earthquake on the hydrology of south-central Alaska, by Roger M. Waller.
- (B) Effects of the March 1964 Alaska earthquake on the hydrology of the Anchorage area, by Roger M. Waller.
- (C) Hydrologic effects of the earthquake of March 27, 1964, outside Alaska, by Robert C. Vorhis, *with sections on* Hydroseismograms from the Nunn-Bush Shoe Co. well, Wisconsin, by Elmer E. Rexin and Robert C. Vorhis, *and* Alaska earthquake effects on ground water in Iowa, by R. W. Coble.
- (D) Effects of the March 1964 Alaska earthquake on glaciers, by Austin Post.
- (E) Seismic seiches from the March 1964 Alaska earthquake, by Arthur McGarr and Robert C. Vorhis.

