Public-data File 91-22mm

NATIONAL URANIUM RESOURCE EVALUATION GEOCHEMICAL DATA FOR STREAM- AND LAKE-SEDIMENT SAMPLES IN THE TANANA QUADRANGLE, ALASKA

by

M.A. Wiltse

Alaska Division of Geological & Geophysical Surveys

December 1990

THIS REPORT HAS NOT BEEN REVIEWED FOR TECHNICAL CONTENT (EXCEPT AS NOTED IN TEXT) OR FOR CONFORMITY TO THE EDITORIAL STANDARDS OF DGGS.

794 University Avenue, Suite 200 Fairbanks, Alaska 99709-3645

NATIONAL URANIUM RESOURCE EVALUATION GEOCHEMICAL DATA FOR STREAM- AND LAKE-SEDIMENT SAMPLES IN THE TANANA QUADRANGLE, ALASKA

by M.A. Wiltse

INTRODUCTION

Purpose:

In December of 1990 the Alaska Division of Geological and Geophysical Surveys (ADGGS) began a mineral resource evaluation of those lands still available for state selection under the Alaska Statehood Act. As part of that process ADGGS is reviewing the stream- and lake-sediment geochemical data generated during the U.S. Department of Energy, National Uranium Resource Evaluation (NURE) program.

This Public-data File has been released so that a summary of that data is available to interested persons. This publication has not been formally reviewed for technical accuracy or for conformity to the editorial standards of ADGGS.

Scope of data:

ADGGS has reviewed NURE geochemical data for the following 1:250,000 quadrangles:

Anchorage	ANC
Baird Mountains	XBM
Beaver	BVR
Bendeleben	BEN
Bering Glacier	XBG
Bettles	BET
Big Delta	XBD
Black River	BLR
Candle	CAN
Chandalar	CHN
Charley River	CHR
Circle	CIR
Eagle	EAG
Gulkana	GUL
Healy	HEA
Hughes	HUG
Iditarod	IDT
Kateel River	KAT
Lime Hills	LIM
Livengood	LIV
Medfra	MED
Melozitna	MLZ
Misheguk Mountain	MIS
Mount Hayes	XMH
Nabesna	NAB

Nome	NOM
Norton Bay	NOB
Nulato	NUT
Phillip Smith Mountains	PSM
Point Hope	XPH
Point Lay	XPL
Ruby	RUB
Selawik	SLK
Shungnak	SHU
Sleetmute	SLT
Solomon	SOL
Talkeetna Mountains	TLM
Tanacross	TNX
Тапапа	TAN
Teller	TEL
Umiat	UMI
Unalakleet	UKT
Utukok River	XUR
Valdez	VAL
Wiseman	WIS

Limitations of data:

Our review has been limited to the following elements: Ag, As, Au, Ba, Be, Bi, Cd, Co, Cr, Cu, Fe, La, Mn, Mo, Ni, Pb, Sb, Sn, Ti, U, U/Th, V, W, Zn

The Nure data set also contains analyses for: Al, Ca, Ce, Cl, Cs, Dy, Eu, Hf, K, Li, Lu, Mg, Na, Nb, Rb, Sc, Sm, Sr, Ta, Tb, Th, Yb, Zr. These data have not been analyzed in the present study.

Because of the procedures used in generating the initial chemical analyses, the NURE geochemical data set has severe limitations. Many elements were determined at only a few sample sites resulting in many samples having incomplete data coverage. The detection limit for many elements is high, making those data of limited effectiveness in delineating mineral resources. Regardless of these and other shortcomings, however, the NURE data do provide information concerning mineralization in many poorly accessible parts of Alaska.

Contents:

This Public-data File (PDF), and the PDF's for the above listed quadrangles, contain a columnar ASCII file on a 5 1/4" high density floppy disk that includes: sample number, replicate code, sample type code, latitude, longitude, and the complete set of elemental analyses available for each sample in the quadrangle. These data are consistently ordered in the file as shown below:

Sample number, Replicate code, Latitude, Longitude, Sample-type code, U Ag Bi Cd Cu Nb Ni Pb Sn W As Zr Mo Be Li Al Au Ba Ca Ce Cl Co Cr Cs Dy Eu Fe Hf K La Lu Mn Mg Na Rb Sb Sc Sm Sr Ta Tb Th Ti V Yb Zn U/Th

All values are entered as parts per million. The data file bears a three letter identification and the extension "ASC" (eg. EAG.ASC for the Eagle Quadrangle data file).

A sample replicate code of "0" indicates that the sample is the initial sample taken at a site and is the code found for most samples. Subsequent samples collected from the same site have successively

higher integer designations. Sample type codes range from "01" to "99". The definition of these codes is found in Appendix A "Key to Sample Types".

Within the elemental analysis fields of a sample, values of -999 indicate that no analyses was attempted for that element. Other negative numbers (eg. -5) in an elemental analysis field of a record indicate that the element was not detected at a level equal to the absolute value of the negative number tabulated.

TREATMENT OF DATA

Elements:

Although all the elemental NURE data available for a quadrangle is included in the digital ASCII file supplied with this PDF, only a 24 element subset of data was analyzed for this PDF: Ag, As, Au, Ba, Be, Bi, Cd, Co, Cr, Cu, Fe, La, Mn, Mo, Ni, Pb, Sb, Sn, Ti, U, U/Th, V, W, Zn.

Grouping of data:

The majority of the Alaska Nure geochemical data is derived from stream sediment or lake sediment samples. Many data sets, however, have a few samples that are subtypes of these two fundamental sample groups. For the purpose of the data review released in this PDF, all subtype samples have been recoded to either the stream sediment type or the lake sediment type, whichever type they most closely resembled. We estimate that less than 1 percent of the samples encountered in this review were recoded.

Following sample-type recoding, brief summary statistics were calculated separately for the stream sediment samples (type=12) and for the lake sediment samples (type=13). These statistics provide a quick reference to the number of samples that have analytical values exceeding the detection limit and provide an indication of the geochemical dispersion of the elements for each sample type.

Single-element Pseudomaps of the data have been made that show the location of all samples having analytical values greater than the mean. This was accomplished by separately standardizing the data for each sample type, recoding all standard scores that were less-than-or-equal-to-zero to zero and then plotting a symbol at each sample site, the size of which is proportional to the elemental standardized value (Z-score) at that sample site. Because Z-scores are measures of standard deviation, this procedure results in a psuedomap with varying symbol size that directly reflects how far a sample's element content is above the mean. The larger symbols correspond to element values that are farthest above the mean value for the element in question. A Symbol-size key is provided in figure 1 which indicates the symbol size for element abundances from 1 to 6 standard deviations above the mean.

THE FOLLOWING RESULTS ARE FOR:
TYPE = 12.000

TOTAL OBSERVATIONS: 583

	Ü	AG	BI	CD	CU
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	583 0.370 80.140 5.989 7.239	4 5.000 9.000 6.000 2.000	99 5.000 25.000 8.818 4.190	5.000 11.000 6.750 2.872	547 10.000 144.000 34.678 16.725
	NI	ВЯ	SN	Ŵ	AS
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	515 15.000 172.000 41.592 22.892	480 5.000 141.000 11.942 8.075	46 10.000 330.000 39.239 59.864	98 15.000 203.000 27.898 22.492	553 5.000 492.000 19.669 31.787
	MO	BE	AU	ва	CO
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	· · · · · · · · · · · · · · · · · · ·	0	7 0.120 0.410 0.213 0.106	572 344.000 2818.000 887.524 304.667	583 6.300 186.600 24.937 14.644
	CR	FE	MN	SB	TI
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	577 20.000 2891.000 122.101 175.987	582 9005.000 148500.000 34406.108 11907.438	583 115.000 11270.000 908.211 824.681	15 2.000 9.000 4.600 2.230	572 2017.000 17060.000 5073.598 1470.054
	V	ZN	UTH	LA	
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	582 25.000 277.000 128.730 31.276	218 31.000 709.000 149.528 82.115	577 0.109 2.337 0.415 0.213	569 17.000 390.000 46.012 32.503	

THE FOLLOWING RESULTS ARE FOR:

TYPE = 13.000

TOTAL OBSERVATIONS: 87

	U	AG	BI	CD	CU
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	87 0.770 5.290 2.787 0.617	•	5 5.000 7.000 6.200 0.837	· · ·	86 21.000 94.000 50.244 15.942
	NI	PB	SN	W	AS
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	86 20.000 87.000 44.349 14.753	79 5.000 19.000 10.430 3.525	1 16.000 16.000 16.000	27 15.000 35.000 20.889 5.542	80 5.000 949.000 25.725 104.848
	МО	BE	AU	ВА	СО
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV		· · · · · · · · · · · · · · · · · · ·	0	84 511.000 1851.000 971.881 202.333	87 8.300 83.600 25.600 15.505
	CR	FE	MN	SB	TI
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	86 29.000 126.000 83.453 17.433	86 9980.000 223000.000 31356.628 22021.317	87 192.000 8089.000 553.230 828.913	· · ·	85 1623.000 5527.000 4396.635 764.497
	Δ	ZN	UTH	LA	
N OF CASES MINIMUM MAXIMUM MEAN STANDARD DEV	86 36.000 178.000 124.535 23.486	42 60.000 261.000 165.452 59.893	85 0.253 1.763 0.380 0.167	78 18.000 39.000 30.987 4.468	

THESE RESULTS ARE FOR tank NURE DATA


TOTAL OBSERVATIONS: 670

Figure 1. Symbol-size key for single element pseudomaps indicating the size of plotted symbols for values that are from 1 to 6 standard deviations above the mean.




1 2 3 4 5 6
STANDARD DEVIATIONS ABOVE THE MEAN


TANZMAP NURE DATA FOR A




TANZMAP NURE DATA FOR AU




TANZMAP NURE DATA FOR BE




TANZMAP NURE DATA FOR CD




TANZMAP NURE DATA FOR CO




TANZMAP NURE DATA FOR UTH




TANZMAP NURE DATA FOR U




TANZMAP NURE DATA FOR PB




TANZMAP NURE DATA FOR NI




TANZMAP NURE DATA FOR LA




TANZMAP NURE DATA FOR BA




TANZMAP NURE DATA FOR MN



TANZMAP NURE DATA FOR MO




TANZMAP NURE DATA FOR FE




+ ~


TANZMAP NURE DATA FOR CU




TANZMAP NURE DATA FOR CR




TANZMAP NURE DATA FOR TI



TANZMAP NURE DATA FOR AG




TANZMAP NURE DATA FOR SB




, -


TANZMAP NURE DATA FOR SN




TANZMAP NURE DATA FOR V




TANZMAP NURE DATA FOR W



TANZMAP NURE DATA FOR ZN



TANZMAP NURE DATA FOR BI



APPENDIX A

KEY TO SAMPLE TYPES

This numerical key provides the necessary tie between the specific type or form of each sample taken and each individual suite of field and laboratory data to which the sample relates. It defines the various sample types collected by the LASL in the DOE HSSR for uranium.

The two-digit <u>key number</u> assigned to each <u>sample type</u> designates three distinct <u>properties</u> of the samples taken. These properties are: (a) The general sample <u>source</u> (spring, stream, dry stream, etc.); (b) The sample <u>medium</u> (water or sediment); and (c) The <u>treatment</u> given the sample in the field or laboratory prior to its analysis by the LASL.

The <u>key numbers</u> are inserted in the sample type columns of the specially formatted DOE sample numbering system to positively identify the sample type for all LASL sample data submitted.

KEY NO.

SOURCE / MEDIUM / TREATMENT

- 01 Spring water sample untreated.
- 02 Stream water sample untreated.
- 03 Well water sample untreated.
- 04 Natural pond water sample untreated.
- 05 Artificial pond water sample untreated.
- 06 Spring water sample filtered through a 0.45-µ membrane filter and acidified to a pH of ≤1 with reagent-grade nitric acid (HNO₃).
- 07 Stream water sample filtered through a 0.45-µ membrane filter and acidified to a pH of <1 with reagent-grade nitric acid (HNO₂).
- 08 Well water sample filtered through a 0.45- μ membrane filter and acidified to a pH of ≤ 1 with reagent-grade nitric acid (HNO₃).
- 09 Natural pond water sample filtered through a 0.45-µ membrane filter and acidified to a pH of ≤1 with reagent-grade nitric acid (HNO₃).
- 10 Artificial pond water sample filtered through a 0.45-μ membrane filter and acidified to a pH of <1 with reagent-grade nitric acid (HNO₃).
- 11 Wet spring sediment sample dried at <100°C and sieved to -100 mesh through stainless steel sieves.
- 12 Wet stream sediment sample dried at <100°C and sieved to -100 mesh through stainless steel sieves.
- 13 Wet natural pond sediment sample dried at ≤100°C and sieved to -100 mesh through stainless steel sieves.

- 14 Wet artificial pond sediment sample dried at <100°C and sieved to -100 mesh through stainless steel sieves.
- 15 Dry stream sediment sample <u>dried</u> at ≤100°C (if necessary) <u>and sieved to</u> -100 mesh through stainless steel sieves.
- 26 Spring water sample acidified to a pH of ≤1 with reagent-grade nitric acid (HNO3).
- 27 Stream water sample acidified to a pH of ≤ 1 with reagent-grade nitric acid (HNO3).
- 29 Natural pond or lake water sample acidified to a pH of ≤ 1 with reagent-grade nitric acid (HNO₃).
- 31- Wet spring sediment sample dried at <100°C and sieved to -40 mesh through stainless steel sieves.
- Wet stream sediment sample dried at <10 10 and sieved to -40 mesh through stainless steel sieves.
- 33- <u>Wet natural lake sediment</u> sample <u>dried</u> at <100°C <u>and sieved to -40 mesh</u> through stainless steel sieves.
- Dry stream sediment sample dried at ≤100°C and sieved to -40 mesh through stainless steel sieves.
- 41- Wet spring sediment sample dried at ≤100°C and sieved to -80 mesh through stainless steel sieves.
- 42- Wet stream sediment sample dried at <100°C and sieved to -80 mesh through stainless steel sieves.
- 43- Wet natural lake sediment sample dried at ≤100°C and sieved to -80 mesh through stainless steel sieves.
- 45- Dry stream sediment sample dried at <100°C and sieved to -80 mesh through stainless steel sieves.
- 51- <u>Wet spring sediment</u> sample <u>dried</u> at ≤100°C <u>and sieved to -170 mesh</u> through stainless steel sieves.
- 52- Wet stream sediment sample dried at ≤100°C and sieved to -170 mesh through stainless steel sieves.
- 53- Wet natural lake sediment sample dried at <100°C and sieved to -170 mesh through stainless steel sieves.
- 55- Dry stream sediment sample dried at ≤100°C and sieved to -170 mesh through stainless steel sieves.
- 61- Wet spring sediment sample dried at -100°C and sieved to -230 mesh through stainless steel sieves.
- 62- Wet stream sediment sample dried at -100°C and sieved to -230 mesh through stainless steel sieves.
- 63- Wet natural lake sediment sample dried at -100°C and sieved to -230 mesh through stainless steel sieves.

- 65- Dry stream sediment sample dried at -100°C and sieved to -230 mesh through stainless steel sieves.
- 71- Sediment sample collected from the stream bank, dried at <100°C, and sieved to -40 mesh through stainless steel sieves.
- 72- Sediment sample collected from the stream bank, dried at ≤100°C, and sieved to -80 mesh through stainless steel sieves.
- 73- Sediment sample collected from the stream bank, dried at $\leq 100^{\circ}$ C, and sieved to = 100 mesh through stainless steel sieves.
- 74- Sediment sample collected from the stream bank, dried at <100°C, and sieved to -170 mesh through stainless steel sieves.
- 75- Sediment sample collected from the stream bank, dried at ≤100°C, and sieved to -230 mesh through stainless steel sieves.
- 96 Dry natural pond sediment sample <u>dried</u> at <100°C (if necessary) <u>and</u> <u>sieved to -100 mesh</u> through stainless steel sieves.
- 97 Dry artificial pond sediment sample dried at <100°C (if necessary) and sieved to -100 mesh through stainless steel sieves.
- 98 Other water These key numbers are to be used only for water (98) or sediment (99) samples coming from a special source and/
- 99 Other sediment or given a special treatment not described for any of the types of samples above.

PDF 91 - 22: ERRATA

ONE ELEMENT WAS OMITTED FROM LISTING OF ELEMENTS ON DISK. THAT ELEMENT WASSE.....IT SHOULD GO BETWEEN AS AND ZR.

CORRECT ORDER OF ELEMENTS ON FILE IS:

U AG BI CD CU NB NI PB SN
W AS SE ZR MO BE LI AL AU
BA CA CE CL CO CR CS DY EU
FE HF K LA LU MN MG NA RB
SB SC SM SR TA TB TH TI V
YB ZN and U/TH

PDF 91-22 - UPDATE

The section of PDF 91-22 which describes the format of the NURE data as it is available on computer disk has changed. Instead of all data on one file/quadrangle in columnar format separated by blanks, it is now split into 3 files/quadrangle with commas and blanks separating the fields. The new files are named "NXXX#.ASC". N is for NURE data, XXX is the 3 character quadrangle identification, and # is 1, 2, or 3. This new version will make it easier for users to input the data directly into Quatro-Pro, Lotus, or other spread sheets with a 250 character limit on record length. In Quatro-Pro use IMPORT option, ASCII file, QUOTE & COMMA delimited. Two records were added in front of the data:

1.) a header record which says

"Part <n>, Quadrangle: <name>"

2.) a record with column headings so users can tell which elements are in the file and the order. The column headings are comma and blank delimited too. The data is still in ASCII format and the commas can be eliminated by using a variety of text editors.

Following are the formats of the 3 files. Column I was left blank for all records so that all data in the files could be printed even when the first item is interpreted as a carriage control character.

Record 1: 55 Characters of text. - starts col 2 and length depends on length of quadrangle name. It is enclosed in quotes.

Record 2: col 2-39

"Samp-Id","RC", "Lat.", "Long.", "ST" (Sample Type--see main text)

Starting in col 40, 14 groups of: "Xx" which are the elements names for the columns. For this record they are: U, Ag, Bi, Cd, Cu, Nb, Ni, Pb, Sn, W, As, Se, Zr, and Mo. NOTE: There is NOT a comma after the last item and all items are enclosed in quotes.

Record 3 to end:

col 2-8. 7 digit sample number.

col 9-10 ", " - a comma followed by a blank

col 11-13 replicate code - 3 digits allowed, most values will be 0 or 1 digit.

col 14-24", " followed by Latitude in decimal degrees with 5 decimal places col 25-35", " followed by Longitude in decimal degrees with 5 decimal places

col 36-39", " followed by 2 digit sample type

Starting in col 40, 17 groups of "," (comma) followed by 8 digit value of element in ppm. Decimal point is present. None of the values require all 8 digits so that leaves a blank space after the comma. NOTE: no comma after the last item.

FILE 2:

Record 1: - Same as for file 1

Record 2: col 2 to 8 - "Samp-Id"

Starting in col 9, 17 groups of: "Xx", which are element names for the columns. For this file they are: Be, Li, Al, Au, Ba, Ca, Cc, Cl, Co, Cr, Cs, Dy, Eu, Fe, Hf, K, and La. NOTE: no comma at end and items enclosed in quotes.

Record 3 to end:

Col 2-8 7 digit sample number.

Starting in col 9, 17 groups of ",", followed by 8 eight digit value of element. As in File 1.

FILE 3: Same format as file 2 with different elements. The elements are: Lu, Mn, Mg, Na, Rb, Sb, Sc, Sm, Sr, Ta, Tb, Th, Ti, V, Yb, Zn, and U/Th

NOTE: In the original listing of the elements, Se was accidently left out. It goes between As and Zr.

A copy of this file is included on disk. It is labeled README, NUR. If there are any problems reading this data contact Shirley Liss at DGGS. (907) - 474 - 7147.