




Base from U.S. Geological Survey, 1975.
Geology based on field reconnaissance and
interpretation of NASA Ames photos 2420-
2423, 2487-2489, accession 2786, July 1979,
and 9766-9768, accession 3101, July 1982.

A 65KA

SCALE 1:63,360

1 0 2 3 4 MILES

1 0 1 2 3 4 5 KILOMETERS

CONTOUR INTERVAL 50 FEET

1:250,000 SCALE, 1:50,000

TRUE NORTH
MAGNETIC N

Geology by S.E. Rawlinson, 1981-85, assisted
by S.B. Hardy, 1981-82, and D.R. Hickmott
1983-85. Reviewed by R.D. Reger.

SURFICIAL-GEOLOGIC MAP OF THE BEECHEY POINT A-3 QUADRANGLE, ALASKA

By
Stuart E. Rawlinson
1986

MAP SYMBOLS

- Pingo
 - Broad-based mound (Walker and others, 1985)
 - Measured section
- _____ Contact
- — — — — Terrace scarp, often poorly defined; tick marks on low side; separates levels within units
- - - - - Boundary of gravel pit
- Qf-Qfi Combination unit; dominant unit listed first
- or Qs/Qt Indicates Qs overlies Qt
- Facility pad or disturbed area or both

DESCRIPTION OF MAP UNITS

Qs SAND COVER DEPOSITS - Fine to medium sand derived from dune deposits (Qsd) deposited downward in a sheet. Deposits proximal to the dunes are several meters thick and thin downward from the dunes. As seen on aerial photographs, the deposits infill low-center ice-wedge polygons, thaw ponds, and small thaw lakes. The deposits are arbitrarily mapped to the boundaries of underlying units because actual boundaries cannot be distinguished. Surfaces support grasses, sedges, and tundra.

Qsd SAND-DUNE DEPOSITS - Fine to medium sand derived from barren flood-plain, delta, and beach deposits and deposited as dunes by wind. Dune deposits are as thick as 6 m, but generally range in thickness from 1 to 4 m. Dunes are most commonly longitudinal and trend northeast-southwest. Unvegetated dunes are being reworked.

Q1 THAW-LAKE DEPOSITS - Peat and pebbly silt or fine sand, or mixtures or interbeds of all three, deposited in basins of thaw lakes by lacustrine and eolian processes. A topmost and often a basal bed of peat bound interbeds of peat and silt or fine sand. These deposits rarely exceed 2 m total thickness and are generally less than 1 m thick; superposition of deposits accounts for thicker deposits. Deposits over the Qap are otherwise featureless. Surface disturbance is common. Surfaces are very wet and are featureless or marked by indistinct large-diameter ice-vedge polygons. Surfaces support grasses and sedges.

ICE-RICH THIN-LAKE DEPOSITS - Peat and pebbly silt or fine sand, or mixtures or interbeds of all three, deposited in basins of thaw lakes by lacustrine and eolian processes. A topmost and often a basal bed of peat bound interbeds of peat and pebbly silt or fine sand. These deposits rarely exceed 2 m total thickness and are generally less than 1 m thick; superposition of deposits accounts for thicker deposits. Deposits are cap unless otherwise indicated. Cryoturbation is common. Surfaces range from wet to dry and are marked by low-center ice-wedge polygons. Surfaces support grasses, sedges, and tundra.


ACTIVE FLOOD-PLAIN DEPOSITS - Fine to medium sand or sandy gravel, or both, deposited in channels of modern flood plains by streams. Sandy gravel is dominant. Where present, sand deposits are moderately sorted, generally 0.5 m or less thick, and overlie poorly sorted sandy gravel with a sharp contact. Subrounded to rounded pebbles and cobbles are common in the sandy gravel. Deposits extend up to 1 m above the mean stream surface. Surfaces of sandy gravel are unvegetated; sandy surfaces are unvegetated or support sparse grasses, sedges, and willow.

INACTIVE FLOOD-PLAIN DEPOSITS - Peat and pebbly silt or fine to medium sand, and gravel interbeds of all three, deposited in flood-plain overbank environments by fluvial, eolian, and lacustrine processes and underlying sandy gravel, or gravel, or interbeds of all three, deposited in channels by fluvial processes. The sandy gravel and gravel layers may be up to 0.5 m thick and generally consist of a topmost bed of peat up to 0.5 m thick and underlying pebbly silt or sand about 0.5 m thick. These deposits are in sharp contact with channel channel deposits. The pebbly silt or sand is rounded, occasional interbeds of pebbles and cobbles are common in channel deposits. These deposits rarely exceed 2 m high and are generally less than 1 m high. Surfaces support grasses, sedges, and tundra.

Q9a **ABANDONED FLOOD-PLAIN DEPOSITS** - Peat and pebbly silt or fine to medium sand, or mixtures or interbeds of all three, deposited in flood-plain channels and on the flood-plain surface by fluvial, lacustrine processes, and in underlying sand, sandy gravel, or gravel, or interbeds of all three, deposited in channels by streams; sandy gravel is the dominant underlying deposit. Overbank deposits are generally consist of a topmost bed of peat up to 1 m thick, cryoturbated and vertically foliated and underlying, interbedded pebbly, silt, pebbly fine sand, and peat all up to 2 m thick. Overbank deposits are in sharp contrast with underlying deposits. Solonchaks are common in the peat and are imbricated pebbles and cobbles are common in channel deposits. Especially in the peat, these deposits rarely exceed 3 m high and are generally less than 2 m high. They are not bounded by any surface, but they are bounded by the water table and ice wedge polygons. Surfaces support tundra; centers of polygons support grasses and sedges.

Qat ALLUVIAL-TERRACE DEPOSITS - Peat and pebbly silt or fine to medium sand, or mixtures or interbeds of all three, deposited in flood-plain overbank and channel deposits. The pebbles are rounded to subangular, and may be in sandy gravel, or gravel, or interbeds of all three, deposited in channels by stream; sand/gravel is the dominant underlying deposit. Overbank deposits consist of the topmost bed of peat up to 1 m thick that is often cryoturbated and vertically foliated and underlying, interbedded pebbly silt and fine sand and peat up to 2 m thick. The overbank deposits are in sharp contact with the underlying channel deposits. The channel deposits are pebbly silt and fine sand and peat, and are commonly 1 to 2 m thick. Scarp faces are common in the channel deposits; these clasts are sometimes imbricated. Exposures of these deposits rarely exceed 4 m high and are generally less than 3 m wide. Scarps that bound the deposits are commonly 1 to 2 m high and are low-center ice-wedge polygons. Surfaces support tundra; centers of polygons support grasses and sedges.

CORRELATION OF MAP UNITS



Q. UNDIFFERENTIATED ALLUVIUM. - Peat and silt or fine to medium sand, or mixtures or interbeds of all three, deposited in flood-plain, overbank environments by fluvial, eolian, and lacustrine processes and underlying sand, sandy gravel, or gravel, or interbeds of all three, deposited in channels of flood plain, or by streamflow in gravel channels. The deposits are poorly sorted. The deposits are up to 5 m thick but average 2.5 m thick. The top of these deposits is composed of peat and peat-rich pebbly silt up to 4.5 m thick; average pebbly silt is 1 m. The deposits are composed of peat, peat-rich pebbly silt, pebbly silt, silty sand, and fine sand compose up to 3.5 m of the bottom of these deposits; average thickness is 1.5 m. Channel deposits are poorly to moderately pebbly and pebbles and cobbles are abundant and abundant. The deposits that bound the unit are indistinct. The surfaces are generally wet and marked by low-center ice-wedge polygons. Surfaces support grasses, sedges, and tundra.

QAP ALLUVIAL-PLAIN DEPOSITS - Pebble fine sand and silt deposited by wind; underlying, interbedded pebbly fine to medium sand and sandy gravel; and sandy gravel deposited by wind. The pebbly sand and silt is overlain by an alluvial plain section often consists of peat (0.3 to 1.3 m thick) with an interbed of pebbly sandy silt (0.1 to 0.4 m thick) and sometimes underlying thaw-lake deposits. The pebbly sandy silt is dark grayish brown when wet and light gray when dry. The pebbly, is shaped somewhat like a flattened sphere. The pebbly sand silt is probably loose with pebbles because its distribution is extensive. Thaw-lake deposits often overlie the pebbly fine sand and silt, but are not mapped because they cannot be recognized outside lake-basin boundaries and are photographed. The pebbly sand silt is composed of fine to medium oxidized brownish yellow. Pebbles in the sand are most commonly brown and black chert; all are polished and some are wind scoured. These deposits are up to 4 m thick, including the peat, loess, and thaw-lake deposits.

QAP ALLUVIAL-PLAIN DEPOSITS - Pebble fine sand and silt overlain by interbedded pebbly fine to medium sand and sandy gravel. When present, the interbedded interval is up to 6 m thick, and averages 3 m thick. The modal diameter of the gravel is 3 cm; pebbles are subrounded to rounded. The lower part of the alluvial-plain deposit is dominated by sand and silt, although thin layers of pebbly sand, organic- and wood-rich silt are present. Alluvial-plain deposits have been measured as deep as 16 m below the ground surface in gravel pits. The modal diameter of the gravel varies from 2 to 5 cm; cobbles are common and boulders are sometimes present near the surface. The pebbly sand and silt is organic- and wood-rich silt sometimes mark changes in the clast mode.

Discontinuously along the coast and up to 9 km inland, pebbly marine mud, interbedded sand and gravelly sand, and cobbles and boulders of the Flaxman Member of the Gubik Formation (Dinter, 1985) crop out at or near sea level. These deposits are generally less than 1 m thick and are overlain by the wind-deposited pebbly fine sand. Surfaces are moist to dry and marked by low- and high-center ice-wedge polygons.

Tsg SAND AND GRAVEL. - Unconsolidated sand and gravel deposited by fluvial and eolian processes and underlying thinly laminated, gray to brown clay and silt beds deposited in the Salt River delta. The sand and gravel beds are composed of quartz pebbles, and others, the clay and silt beds commonly contain organic matter, small chert pebbles, and mud lumps along the bedding planes. Most pebbles are on the upper surface of the clay units and of a size and shape that would be expected from a stream bed. Desiccation cracks are preserved in a few beds. Limonite staining is common. The sand and gravel unit is pink, tan, orange, and yellow. Limestone zones several centimeters thick locally cement the sand and gravel into sandstone and rock. The gravel is composed primarily of light-colored quartz pebbles and cobbles of volcanic ash. The sand and gravel is commonly found in common some sand units. Beds of volcanic ash are interlayered with the sand and gravel. Coarse gravel deposited by river processes often caps the unit. The gravel cap is composed primarily of pebbles and cobbles of light-colored quartz, and green tuffaceous sandstone, and minor amounts of limestone, which give surfaces a characteristic white appearance. The age of this capping gravel is uncertain. Surfaces support tundra.

REFERENCES CITED

- Detterman, R.L., Reiser, H.N., Brosge, W.P., and Dutro, J.T. Jr., 1975, Post-Carboniferous stratigraphy, northeastern Alaska: U. S. Geological Survey Professional Paper 886, p. 57-39.
- Dinier, D.A., 1985, Quaternary sedimentation of the Alaskan Beaufort Shelf; influence of regional tectonics, fluctuating sea levels, and glacial sediment sources: *Tectonophysics*, v. 114, no.1-4, p. 133-161.
- Walker, D.A., Walker, M.D., Everett, K.R., and Webber, P.J., 1985, Pingos of the Prudhoe Bay region, Alaska: Arctic and Alpine Research, v. 17, no. 3, p. 321-

AGE-DATE INFORMATION

Station	Coordinates	Sample	Type	Material	Age (y. B.P.)
BPA3-2	N 70° 6'35" W148°37'58"	BPA3-2-3	C-14	Sandy peat	10,540 ± 310
BPA3-10	N 70°14'33" W148°15'47"	BPA3-10-3	C-14	Peat	4,640 ± 90
BPA3-10	N 70°14'33" W148°15'47"	BPA3-10-5	C-14	Wood	>38,400
BPA3-13	N 70° 0'24" W148°27'30"	BPA3-13-4	C-14	Peat	>34,800
BPA3-13	N 70° 0'24" W148°27'30"	BPA3-13-7	TL	Silt	3,140 ± 500

Stratigraphy and significance

0.7 m below surface; basal peat; minimum age of earliest alluviation associated with drainage Sagavanirktok River through Putuligayuk River basin.

5.1 m below surface; paleosol that suggests sea level was several meters lower at present probably represents base of Holocene alluvium.

8.7 m below surface; minimum age of deposits deeper than 8.7 m.


7.4 m below surface; peat in topmost silt of sequence of alternating channel and overbank alluvium; minimum age of this sequence.

8.1 m below surface; basal sample of silt bed (see BPA3-13-4); date is inconsistent with C-

EXPLANATION OF STRATIGRAPHIC SECTIONS

- Figure 1 is a stratigraphic column for BPB3-00. The column shows depth from 0.0 to 3.0 meters. The stratigraphy includes: 0.0-0.5m: Vegetative mat; 0.5-1.0m: Peat; 1.0-1.5m: Silty peat; 1.5-2.0m: Sandy peat; 2.0-2.5m: Silt; 2.5-3.0m: Sandy silt; 3.0-3.5m: Peaty silt; 3.5-4.0m: Fine sand. The column is divided into two sections: the upper section (0.0-2.0m) is labeled 'Medium sand' and the lower section (2.0-4.0m) is labeled 'Coarse sand'. The column is also divided into two sections: the upper section (0.0-2.0m) is labeled 'Ice-wedge pseudomorph' and the lower section (2.0-4.0m) is labeled 'Ice-wedge'. The column is also divided into two sections: the upper section (0.0-2.0m) is labeled 'Ice pods or lenses' and the lower section (2.0-4.0m) is labeled 'Organic stringer'. The column is also divided into two sections: the upper section (0.0-2.0m) is labeled 'Wood' and the lower section (2.0-4.0m) is labeled 'Gastropod (fresh water)'.

STRATIGRAPHIC SECTIONS



PALYNOLOGY ANALYSIS

Station	Coordinates	Sample	Palynomorphs
BPA3-3-13	N 70° 0'24" W 148°27'30"	BPA3-13-3	Betulaceae (F) <u>Lycoodiumsporites</u> sp. (R) Ericaceae (V)

F = frequent (6-15 specimens), R = rare (2-5 specimens),
V = very rare (1 specimen)

FAUNA ANALYSIS

Station	Coordinates	Sample	Fauna
BPA3-3-13	N 70° 0'24" W148°27'30"	BPA3-13-3	shell fragments (F) <u>Cyclammina</u> cf. <u>pacifica</u> (V) reworked

F = frequent (11-32 specimens), V = very rare (1 specimen)

ENGINEERING CHARACTERISTICS

PERMAFROST

- | | |
|-----|---|
| Qs | Continuously frozen, but seasonally thawed to about 1 m deep; may contain pore ice. |
| Qsd | Continuously frozen, but seasonally thawed to about 1 m deep; may contain pore ice. |
| Q1 | Continuously frozen, but seasonally thawed to about 0.6 m deep; massive ice may be present, especially below shallow lake basins. |
| Qti | Continuously frozen, but seasonally thawed to about 0.6 m deep; pore, segregated, and massive ice are present. |
| Qf | Continuously frozen, but seasonally thawed to about 1 m deep; also thawed in zones below channels with water greater than 2 m deep; pore ice may be present. |
| Qfi | Continuously frozen, but seasonally thawed to about 0.6 m deep; pore and segregated ice are present in top fine-grained sediments; some massive ice may be present. |
| Qfa | Continuously frozen, but seasonally thawed to about 0.6 m deep; pore, segregated, and massive ice are present in top fine-grained sediments. |
| Qat | Continuously frozen, but seasonally thawed to about 0.6 m deep; pore, segregated, and massive ice are present in top fine-grained sediments. |
| Qal | Continuously frozen, but seasonally thawed to about 0.6 m deep; pore, segregated, and massive ice are present in top fine-grained sediments. |
| Qap | Continuously frozen, but seasonally thawed to about 0.6 m deep; pore, segregated, and massive ice are present in top fine-grained sediments. |
| Tsg | Continuously frozen, but seasonally thawed to about 0.6 m deep; pore, segregated, and possibly massive ice are present in top fine-grained sediments. |

FROST-ACTION SUSCEPTIBILITY

- Not frost susceptible except where silt included.
- Not frost susceptible except where silt included.
- Moderately to highly frost susceptible; susceptibility increases with increase in silt content.
- Moderately to highly frost susceptible; susceptibility increases with increase in silt content.
- Not frost susceptible.
- Low to moderately frost susceptible; susceptibility increases with increase in silt content.
- Low to moderately frost susceptible; susceptibility increases with increase in silt content; highly frost susceptible in channel fills where silt content is high.
- Low to moderately frost susceptible; susceptibility increases with increase in silt content; highly frost susceptible in channel fills where silt content is high.
- Highly frost susceptible especially in channel fills; susceptibility increases with increase in silt content.
- Moderately to highly frost susceptible; susceptibility increases with increase in silt content.
- Fine-grained sediments; highly frost susceptible.

CONSTRUCTION SUITABILITY

- Qs Good as fill with addition of binder; limited quantity; good foundation base except where very thin and/or silt is included.
- Qsd Good as fill with addition of binder; good foundation base except where silt is included.
- Qt Not good as fill; not good for foundations because of frost susceptibility and possible massive ice.
- Qti Not good as fill; not good for foundations because of frost susceptibility, presence of massive ice, and poor drainage.
- Qf Good as fill; good foundation base except that potential of flooding is high.
- Qfi Good as fill, although thin, fine-grained cover must be removed; acceptable for foundations except where silt content is high, although potential of flooding is high and drainage is poor.
- Qfa Good as fill, although thin, fine-grained cover must be removed; acceptable for foundations except where silt content is high, although potential of flooding is moderate to high and drainage is poor.
- Qai Good as fill, although thin, fine-grained cover must be removed; acceptable for foundations except where silt content is high.
- Qal Good as fill, although thin, fine-grained cover must be removed; not good for foundations because of frost susceptibility, presence of massive ice, and poor drainage.
- Qcp Coarse-grained deposits may be good as fill, although clay size may be too large and/or percent of fine-grained clasts

POTENTIAL PROBLEMS

- Easily eroded when vegetation is removed; often very thin so that settlement of underlying deposits may occur upon thawing.
- Easily eroded when vegetation is removed or not present.
- Differential settlement upon thawing; very poor drainage; possible pingo formation, especially in recently drained basins.
- Severe differential settlement upon thawing; very poor drainage; possible pingo formation.
- Frequent flooding, ice jams, and scour.
- Frequent flooding, ice jams, and scour; differential settlement upon thawing; poor drainage.
- Frequent flooding in low areas, ice jams, and scour; occasional extensive flooding; severe differential settlement upon thawing; poor drainage.
- Severe differential settlement upon thawing; poor drainage.
- Severe differential settlement upon thawing; poor drainage.
- Severe differential settlement upon thawing; poor drainage.