EXPLANATION

This map shows the locations of DGGS and other samples associated with middle to late Quaternary deposits in the Tyonek Quadrangle. Map index numbers are keyed to sites where stratigraphic sections are described and where radiocarbon, tephra, magnetic-susceptibility, pebble, and fossil samples were collected (table 1). Maximum and recessional limits of the late-Wisconsin Naptowne glaciation were identified by photointerpretation and field investigations conducted intermittently from 1977 through 1994. Former glacial extents differ slightly from models proposed by previous workers (Miller and Dobrovolny, 1959; Karlstrom, 1964; Reger and Updike, 1983; Schmoll and others, 1984; Schmoll and Yehle, 1986). These differences reflect the normal advance of knowledge that results from improved analytical methods, better aerial photographs and topographic maps, and more information from hundreds of new exposures and subsurface borings, as well as different interpretations of the available data. Morainal limits shown on this map correlate with type Naptowne moraines in the Skilak Lake (Kenai Quadrangle) and Anchorage (Anchorage Quadrangle) areas (Reger and Pinney, 1996).

MAP SYMBOLS

- Glacial limit during Elmendorf stade of Naptowne glaciation, dashed where approximate, queried where uncertain
- Glacial limit during Skilak stade of Naptowne glaciation, dashed where approximate, queried where uncertain
- Glacial limit during Killey stade of Naptowne glaciation, dashed where approximate, queried where uncertain
- Glacial limit during Moosehorn stade of Naptowne glaciation, dashed where approximate, queried where uncertain
- TY0-5 — Sample locality

REFERENCES CITED

Karlstrom, T.N.V., 1964, Quaternary geology of the Kenai Lowland and glacial history of the Cook Inlet region, Alaska: U.S. Geological Survey Professional Paper 443, 69 p., 2 pls., 5 map sheets, scales 1:63,360, 1:250,000, 1:3,168,000, and 1:5,000,000.

Miller, R.D., and Dobrovolny, Ernest, 1959, Surficial geology of Anchorage and vicinity, Alaska: U.S. Geological Survey Bulletin 1093, 128 p., 1 sheet, scale 1:63,360.

Reger, R.D., and Pinney, D.S., 1996, Late Wisconsin glaciation of the Cook Inlet region with emphasis on Kenai Lowland and implications for early peopling, *in* Davis, N.Y., and Davis, W.E., eds., Adventures through time: Readings in the anthropology of Cook Inlet, Alaska: Anchorage, Cook Inlet Historical Society, p. 15-35.

Reger, R.D., and Updike, R.G., 1983, Upper Cook Inlet region and the Matanuska Valley, *in* Péwé, T.L., and Reger, R.D., eds., Guidebook to permafrost and Quaternary geology along the Richardson and Glenn Highways between Fairbanks and Anchorage, Alaska: Alaska Division of Geological & Geophysical Surveys Guidebook 1, p. 185-263, 1 sheet, scale 1:250,000.

Schmoll, H.R., and Yehle, L.A., 1986, Pleistocene glaciation of the upper Cook Inlet basin, *in* Hamilton T.D., Reed, K.M., and Thorson, R.M., eds., Glaciation in Alaska - the geologic record: Anchorage, Alaska Geological Society, p. 193-218.

Schmoll, H.R., Yehle, L.A., Gardner, C.A., and Odum, J.K., 1984, Guide to surficial geology and glacial stratigraphy in the upper Cook Inlet basin: Anchorage, Alaska Geological Society guidebook, 89 p.

SCALE 1:250,000

5 0 5 10 15 20 MILES

5 0 5 10 15 20 25 30 KILOMETERS

SAMPLE LOCALITIES IN THE TYONEK QUADRANGLE, ALASKA

By

Richard D. Reger, DeAnne S. Pinney, Raymond M. Burke, and Milton A. Wiltse

1996

ALASKA

QUADRANGLE LOCATION

Date: 11/25/96 /files4/deanne/tunal/tyoesamples.grs