MAP AND TABLE SHOWING METAMORPHIC ROCKS OF ALASKA

[References for information are given either in text or in appropriate regional metamorphic facies map (shown in Fig. 2.)]

Unit No.	Metamorphic unit [1,2]	Lithotectonic terrane(s) in which unit occurs[3]	Inclusive age range[4] and protolith[5]	Metamorphic rock type[6] key metamorphic mineral[7]; structural characteristics if relevant	Direct evidence for age of metamorphism[8] and evidence used[9]	Maximum age of metamorphism[10] and evidence used[11]	Minimum age of metamorphism[12] and evidence used[13]
NORTHERN ALASKA							
Western and central Brooks Range							
1	AMP (A) / GNH (A)		AAH	Z and older?; cont. ords., carb., marl. viz., int. & maf. plut.	Qu-mica schist, epidz, marble, metabasite, metatuff.; M_2: lit. gp., bh: M_2: ch, ar, bar ampb, gl, ph	M_2: Z; K-Ar ms and bi ages between 729±22 and 594±18 Ma and Rb-Sr mineral-whole rock isochron age of 698±16 Ma	M_2: 362 Ma; same evidence as for unit 3
2	AMP (DK) / GNH (LDK)		AAH, AAC	mD and older; same protolith and gneiss gives for unit 3	Same rock types as those given for unit 1; M_2: ai, tk, gt, bi; M_3: ab, ep, ch, ser, pumpellyite, lawsonite	----	M_2: E; metamorphism is younger than most affected rocks whose protolith may be -in age. M_3: mD, same evidence as for unit 3
3	GNH ↔ L (KJ)		AAH, AAD AAE	Z(?)--; cont. rocks ... carb., felsic and maf. vol.	Qz-mica schist, calc-silicate, marble, Gr. metagraywacke, metaconglomerate; in early part of high-P phase: lw, glps, gl; late part of high-P phase: ep, ab, ch, ser, ta, ca-amp; in early intermediate- to low-P phase: ab, ch, ar, bi; polydeformed, isoclinally folded	----	m3: K-Ar ages of 172±5 and 167±5 Ma on lib from gt amphibolite at side of ultramafic intrusion; interpreted as cooling ages during late tectonic history prior to emplacement of oceanic terrane on North America; 167±5 Ma = 163 to 172 Ma
4	LPP (KJ)		AAN	-C-Si; cont. rocks, vol., carb.	Metasedi., argil., phyl., metaarch., metabasite, metarch.; ps, ch, ser, ta, ca-amp, bi; polydeformed; bimodal unit; P structure is thin unit appear to correlate with those in unit	----	m3, same evidence as for unit 3	mK; same evidence as for unit 3
5	GNS (KJ)		AAN	-C-D; cont. rocks, vol., carb.	Same rock types as those given for unit 4; vein, ss, ali; same structure characteristics as those given for unit 4	----	m3, same evidence as for unit 3	mK; same evidence as for unit 3
6	LPP (K)		AAD, AAE	Metased., metaarch., metabasite, metabasite; vm, ps, ch, ac	Metased., metaarch., metabasite, metasilite; vm, ps, ch, ac	----	sK; age of youngest protolith	mK; probable correlation of metamorphic history with that in unit 4
7	GNS (K)		AAE	D-M; cont. rocks., carb.	Phyl., -gnitic, metaarch., metased.; ----	----	sK; correlation of metamorphic history with that of unit 6	mK; correlation of metamorphic history with that of unit 6
8	LPP (Kv3) ?	?	AM	D-J; maf. vol. & plut., -ss, -sedi., -clast	Metabasite, metaarch., argil., phyl., metased.; vm, ps, ep, ac; lower contact fault-bounded; relict igneous texture locally preserved in upper parts of thrust	----	m3, same evidence as for unit 3	mK; age of overlying conglomerate that contains pebbles of this unit
9	GNS (Kv3) ?	?	AM, RB Vf(N?)	D-S; mdx., carb., chert	Metagraywacke, phyl., metaarch., metabasite; ch, ab, vm; rarely bi; singly deformed, semipenetrative cleavage	----	m3, same evidence as for unit 3	mK; same evidence as for unit 3
10	LPP/GNS (TV)	?	K	IK; cont. rocks., vol.	Metaarch., slate, phyl., metabasite; vm, ch, ep, bi(?); highest plate of Brooks Range; polydeformed, bending foliation; kinematic indicator data down-to-the-S movement on Kobuk fault zone; S-dipping normal faults; degree of microtallization and shearing increases near faults suggesting metamorphism contemporaneous faulting	----	IK; age of youngest protolith of this unit	sT; correlation of retensional faulting with that of extensional faulting with uplift of Brooks Range
Eastern Brooks Range								
11	LPP (MD)	?	AAN	E-to-C(?)D; cont. rocks., carb., vol.	Semischist, phyl., argil., metaarch., metased.; psc, ps, ch, wm, ch, ab, ar; rarely ep; metamorphic grade increases to S	----	m(?)D; age of youngest protolith	M; age of immediately overlying unmetamorphosed rocks
12	GNS (MD)	?	AAN	Same rock types as those of unit 11; ss, ch, wm, ab, ar, ol	----	----	m(?)D; same evidence as for unit 11	M; same evidence as for unit 11
13	GNS (S) ?	?	AAN	D(?)-DD; cont. rocks., maf. vol.	Pelitic schist, slate, quitte, mafic schist, phyl., greenstone, metaconglomerate; characteristics of plates of this age suggests rocks, were associated with and slightly preceded plutonism	ID; increase in metamorphic grade from chl to ep-bi zone assemblages with decreasing distance from overarching Late Devonian plutons	ID; age of youngest protolith	ID; age of crosscutting plutons (pC?) plutons
14	LPP (Ck)		AAE, AAN, MDN, TN, DHS	Argil., metaarch., metaarkf., metaclast., metabasite, ch, ps, pa, wm	----	----	K; age of youngest protolith	sK; age of youngest crosscutting intermediate-composition plutons
15	LPP (Kd2)		TZ	D(?) in southern two areas, D-sK in northern area; cont. rocks.	Phyl., -gnitic, slate; wm, ch	----	D, older limit on age range of youngest protolith	sK, younger limit on age range of youngest protolith
Seward Peninsula								
16	GNH ↔ I (Kf)		SD	B-D; cont. rocks., carb., marl. sid. igneous rocks	Pelitic schist, quitte, marble, mafic schist, metabasite, orthogneiss; early part of High-P phase: lw, gl, ep, ab, ch, ser, ta, ca-amp, bi; late part of low-P phase: ab, ch, ar, bi, ep, ch, gl; intermediate-P phase: bar ampb, ep, ch, ab, ep; rhythmically deformed, shallow-dipping transposition foliation	High-P part of cycle: ca. 21 Rb-Sr ages on mica from mica schists and orthogneiss; ages of about 150 Ma; intermediate-P phase: bar ampb, ep, ch, ab, ep; ca. 12 ages; Rb-Sr and K-Ar mica ages about 150-130 Ma.	----	----
17	GNH (Kf)		SD	mMu(?)-J(?); maf. vol., ultramafic rocks	Metabasite and minor serpentinite; ol, tw, px in ?t, ep, ep, ac in S; metamorphic grade of this unit increases to S; original igneous grain and gabbroic fabric; unit comprises narrow, fault-bounded belt	----	m2; possible correlation of metamorphic history with that of unit 8	sK; possible correlation of metamorphic history with that of unit 8
18	AMI (K)		SD	B-sMu; cont. rocks., carb., maf. and fel. igneous rocks	Pelitic, calcareous, and quartzofeldspathic schist, marble, quitte, amphibolite; gt, ky, ri, sil, bi, ep; locally metamorphic minerals and structures of unit 16 are preserved in this unit	8-sK; 100- to 80-Ma K-Ar ages on plutons probably from emplacement of unit 16	sK; approximate 100-Ma upper age limit for previous intermediate-P phase of metamorphism of unit 16	IK; 80-Ma K-Ar ages on plutons that crosscut metamorphic effects overprint fabric and mineral assemblages of this unit
19	gPx (K)		SD	B-sMu; same protolith as those of unit 18	----	----	----	----
20	GNS (Kfk)	?	?	mMu(?); cont. rocks., carb., vol.	Qz-mica schist, quitte, metaarch., calcareous, graphitic, and amphibolitic schist	----	mMu; probable age of protolith of metaarch.	K; age of post-metamorphic granitoids in the region
21	AMP (Kf)		?	B(?)-fMu(?); cont. rocks., carb., vol.	Hb gneiss, pelitic gneiss, calc-silicate gneiss, marble, hb, qtz, gt, ri, bi, ol	----	E; possible age of protolith	K; age of plutons whose contact metamorphic effects overprint the fabric and mineral assemblages of this unit
CENTRAL ALASKA								
22	GNS (pC)		NK	Z-pC; fel., and mid. vol., cont. rocks.	Metased., pelitic schist, calc. schist, semischist, quitte, phyl., argil., marble; ch, ctd, ep, ab, ser, ta, ab, and rarely ta and (or) gt	$C(?)$; 514 Ma oldest K-Ar age on mica from ch schist	Z, protolith age	O, age of oldest metamorphic assemblages
23	AMP (Pz) ?	AMP (Kd)	NK	Y-pC; cont. rocks, maf. and fel. vol., plut.	Pelitic schist, calc-schist, metaarch., metased., metaarkf., metaplut.; M1 in pelitic schist: bi, gt, ac, ctd, hz in pelitic schist: ab, ep, ch, ser, ta; locally developed hbphornblende may have accompanied M_2	M_2: 71(?); 921-Ma Rb-Sr whole-rock isochron age (greenish age) on gneiss from metased. M_3: 227(?); 665-Ma Rb-Sr whole-rock isochron age (greenish age) on gneiss from metased.	Y; 1265-Ma U-Pb zircon age (intercept age (greenish age) on gneiss from metased.	mMu; 3990±6-Ma U-Pb zircon age (intercept age (metamorphic age(?)) on zircons from metased.
24	GNS/LAMP (pK) ?	?	RB, NK, BRY	B(?)-mMu; cont. rocks., vol., carb., chert, fel. plut.	Pelitic, quartzofeldspathic, calcareous, and mafic schist, quitte, marble, amphibolite, metaarch., metaplut.; wm, ab, ch, ctd, ep, ac, bi, rarely ana and or; polydeformed; bimodal unit; P structure is thin unit appear to correlate with those in unit	-----	mMu(?); probable age of youngest protolith	sK; 110-Ma K-Ar and U-Pb discordant ages of crosscutting, unmetamorphosed plutons
25	AMP (pK) ?	?	RB	B(?)-mC(?)b; cont. rock., plut., vol.	Pelitic schist, quitte, felsic gneiss, amphibolite, metaarch.; bi, gt, ac, ol; ky in central area; ca in southern area	----	mMu(?); same evidence as for unit 24	sK; same evidence as for unit 24
26	AMP (pK)	GNS (Kd) ?	RB	B(?)-D; cont. rocks., carb., igneous rocks, fel. plut., carb.	Quartzofeldspathic paragneiss, calcareous, pelitic schist, quitte, marble, metaplut., amphibolite, calcareous schist and gneiss, marble, gryllenite; M1: ?t, gt, mmSt(?); synkinematic with distal deformation and isoclinal folding. M_2: ch, wm, ctd, and gt at one locality; synkinematic with second deformation; locally with axial planar lineation parallel to that in some of intense cataclasis	M_2: 178-sK; probable correlation of metamorphism with that of plutons of the age suggests rocks, were associated with and slightly preceded plutonism	D; age of youngest protolith	sK; 104-Ma K-Ar age of crosscutting, unmetamorphosed phased plutons
27	AMP (pK) ?	?	AMP (Kf)	RB	Same age range as those of unit 26	----	M_2: pD(?); probable D age of youngest protolith; age of protolith of granitoid gneiss; M_3: mD; probable age of protolith of granitoid gneiss	sK; 111-Ma K-Ar age of crosscutting, unmetamorphic plutons
28	GNH (pGk)		RB	B(?)-D; cont. rocks., carb., marl. vol., chert	Pelitic schist, quitte, phyl., slate, metaarch., metabasite, metaarkf., metachert; wm, ch, ctd, ep, and minor gt in upper parts of unit; bi and gt occur rarely	IK-sK; same evidence as for unit 26	D; age of youngest protolith	sK; same evidence as for unit 27

Unit No.	Metamorphic unit [1,2]	Lithotectonic terrane(s) in which unit occurs[3]	Inclusive age range[4] and protolith[5]	Metamorphic rock type[6] key metamorphic mineral[7]; structural characteristics if relevant	Direct evidence for age of metamorphism[8] and evidence used[9]	Maximum age of metamorphism[10] and evidence used[11]	Minimum age of metamorphism[12] and evidence used[13]
CENTR ...ALASKA (Continued)							
29	LPP (pCk)	TZ, IN	ID-fE; maf. vol. and plut., chert, ss. ands., carb.	Greenstone, metabasalt, metachert, argil., slate, metaarkf., metased., metagraywacke, phyl.; wm, ch, ep, ac, ep, gl; occurs at base of unit in one area; kinematic indicators in Tanana quad. indicate this unit was thrust from the SE	?E-sK; probable association of metamorphism with obduction of this unit from the SE	?E; age of youngest protolith	sK; age of oldest post-metamorphic plutons that intrude this region
30	AMP (KA)	RB	K_2; maf. igneous, fel. plut., cont. rocks.	Amphibolite, orthogneiss, including augen gneiss, pelitic schist, quitte, bi, ms, hb, ep	Y and sK-K-Ar ages of 1.22 Ga, 1.08 Ga, and 126 Ma on hb and amphibolite; and 113 Ma on bi from amphibolite	K_2; U-Pb zircon protolith age of orthogneiss	sK; youngest K-Ar age from amphibolite
SOUTHWESTERN ALASKA							
31	AMP (K) / GNH (K)	KIL	K_1; plut., cont. rocks., maf. vol., carb.	Orthogneiss, pelitic schist, amphibolite, rare marble; hb, ab, gt, epz, bT, and ky at one locality; M_2 assemblages overprinted and largely obliterated by younger evidence; NE strike of foliation and trend of lineation approximately parallel to trend of unit	M_2: K; 1.77-Ga U-Pb age of epz from orthogneiss; K-Ar ages of 1.75 Ga on hb from amphibolite, and Rb-Sr whole-rock isochron. M_3: 1.6; 1.56- to 1.50-Ma K-Ar ages on hb from this unit	K_2; age of protolith	M_2: Y; 1.2-Ga $^{40}Ar/^{39}Ar$ plateau age of hb from gt amphibolite. M_3: 1.6; 1.56; overlying unmetamorphosed plutons
32	GNS (JN)	GD	P(?)-TR(?); vol., oc. sed., maf. and fel. plut., metamorphosed equivalents	Metabasite, metavol., metasands., metaarch., serpentinite; ch, ep, ab, ser, ta; schistose fabric developed in metasands. and metaarch. NE strike of foliation and trend of lineation approximately parallel to trend of unit	----	----	IK; probable age of youngest protolith
33	GNH (JN)	GD	P(?)-TR(?); same protolith as those given for unit 32	Same rock types as those given for unit 32; ch, bi, ep, gt, ac, ep; primary textures and minerals partly preserved; kinematic indicators, mal. mélange-like fabric	TR(?); 231±5-Ma K-Ar age on musc from schist	IK; same evidence as for unit 32	mJ; same evidence as for unit 32
34	DN(J) (JN)	GD	O/S-D(?); basalt, turf., chert, limestone, etc. seds.	Mafic schist, metachert, metalimestone, phyl., quartzose, graphitic, and calcareous schist; ch, ep, ac, and locally gt	m(?)-eK(?); 146±1.5-Ma K-Ar age on musc from mafic schist	TR(?); maximum age for low-grade metamorphism in the region	sK; minimum of age range for 146±1.5-Ma K-Ar age on ms from mafic schist
35	LPP (pK)	GD, NY, TG, TH, RB, KY	sPk-sMz; maf. igneous, vol., ocn. rocks., limestone	Metabasite, metavol., metaarkf., metachert, metalimestone, metased.; ch, ep, ab, ser, ta; schistose fabric developed in metasands. and metaarkf.	None; metamorphic age may differ in various parts of unit	----	sK; age of youngest metamorphosed plutons in the region
EAST- AND SOUTH-CENTRAL ALASKA							
Yukon-Tanana upland							
36	AMP (CD) / AMP (pK) (K)	YT	D(?)-sPk(?); mafic vol., igneous	Eclogite, mafic, calc-magnesian, quartzofeldspathic, and pelitic schist, gneiss; marble; eclogite: gt, jdpx, bar ampb, gln, ep, atspk, ab, ep; pelitic schist: ky, st, gt; polydeformed-to-penetrative fabric; local preserving of above mineralogy in lens, ath, gn, bi, and ch; unit occurs as two klippen; M_2 associated with N- to NW-trending folds and M_3 with NE-trending folds	TR-sE; 212±2-Ma $^{40}Ar/^{39}Ar$ plateau age on ampb from eclogite. M_2: 455(?); 115- to 103-Ma K-Ar ages on mica from schist	----	sK; 115- to 103-Ma K-Ar ages on mica from schist
37	AMI (J?k)	YT (but probably at better related to ST)	Pz-TR(?); mafic vol., limestone	Bt gneiss and schist, amphibolite, marble, quitte, pelitic schist; gt, bi, ep, hz, ms, st, ky; wcl foliated and lineated; M_2 assemblages overprinted by M_3 assemblages; M_3 associated with NE-trending folds and late N-trending open folds	TR-sE; 211±2-Ma $^{40}Ar/^{39}Ar$ plateau age on ampb from metagabbro. 205±3-Ma $^{40}Ar/^{39}Ar$ plateau age on ampb from synmetamorphic plutons (J?g)	----	----
38	J?-SVG (J?k)	YT, SV	P-Si; maf. vol., ocn. seds. and vol., limestone, and minor ultramafic rocks	Greenstone, argil., metaarkf., marble, quitte, gneiss, metabasite, metachert, metased., metaarkf., metavol.; ab, ep, ch, ser, ta, ca-amp, bi, gt, st, ky, wm, ac, and ol at rare locality; consists of thrust remnants which are internally broken by thrust faults	sE; 201±5-Ma $^{40}Ar/^{39}Ar$ plateau age on ms from greenstone	TR; Triassic age of youngest protolith	----
39	GNS (Jk ?)	YT	Be(?); maf., int. and fel. vol., ---, seds.	Quartzose mafic, calcareous, and pelitic schist, phyl., quitte; ch, wm, ep, ab, ser, ta, ca-amp, bi, gt, st, ky, and rarely ol; metased. and metavol. commonly mylonitic and blastomylonitic	?E(?)-eK(?); 175±14-Ma K-Ar age on ms from mylonitic schist (Rb-Sr whole-rock age from correlative(?) Kiondike Schist in Canada	----	----
40	LPP (SG)	MNK, WS, LO, WHM, MS, NX, GL	-C(?)-O-D; seds., carb., vol., chert	Argil., slate, semischist, quitte, metaarch., metavol., metachert; wm, ch, ab, ep, ac, ep, gl, ta, and ca occur in one sample, and gl occurs in one other sample; rarely preserved; locally mylonitic or highly foliated	----	D; age of youngest protolith	IK; 91-Ma age of crosscutting, unmetamorphosed plutons that probably intrudes this unit
41	GNI (SN)	YT	E(?)-Be; cont., vol., minor carb.	Quitte, pelitic, feldspathic, and metaarkf. schist, gneiss, and marble; ab, ep, ch, ser, ta, ca-amp, bi, gt, st, ky, and ol; metased. and metaarkf. fill-ward increase in metamorphic grade from ch zone to bi, wm, ac, ep, gt, st zone; ky, ol, mEx polydeformed, mylonitic fabric developed near the NW contact, shown as a thrust fault	sE(?); 138- to 100-Ma K-Ar age on mica from greenschist	Rx; probable age of youngest protolith	sK; 115- to 90-Myr-old crosscutting ages of other unmetamorphosed plutons in the Yukon-Tanana region
42	AMI (SN)	YT	Z-M; cont. rocks., carb., plut., carb.	Pelitic schist and gneiss, quitte, metaarkf., amphibolite, calc-silicate rock, amphibolite, marble, and minor metachert; ab, ep, ch, ser, ta, ca-amp, bi, gt, st, ky, ol, wm, ac, and ep; polydeformed and generally mylonitic; augen gneiss forms basketlike concordant sheets	sE(?); 115-Ma U-Pb age on zircon from augen gneiss; 112-Ma Rb-Sr mineral isochron for augen gneiss and 124-Ma K-Ar age on bt from gt gneiss	Rx; probable age of youngest protolith	sK; same evidence as for unit 41
43	LPP (SN)	YT	E(?)-Be(?); cont. rocks.	Carbonaceous quitte and argil., pelitic schist, metaarkf., marble, wm, ch, clastic textures well preserved	----	Rx; probable age of youngest protolith	sK; same evidence as for unit 41
44	LPP (SNB) (SN)	YT	E(?)-Be(?); same protolith as those given for unit 43	Same rock types as those given for unit 43; wm, ch, and rarely bi, gt, and traces of ep and ac; thrusted; upgraded zone contrast with Tertiary plutons	sE(?); 125-Ma K-Ar whole-rock age on ms-schist and 140-Ma K-Ar age on bi from calc-schist	Rx; probable age of youngest protolith	sK; same evidence as for unit 41
45	GNS (SN)	YT	Be; vol., seds., carb.	Qz-ch-ser. schist, sylonitic schist, metasandst., metaarkf., metachert, metacarb.; wm, ch, ab, ep, ac	----	Rx; probable age of youngest protolith	sK; same evidence as for unit 41
46	AMI (SN)	YT	E(?)-M; seds., vol., plut., minor carb.	Qz-mica schist, bt and hb gneiss, augen gneiss, quitte, amphibolite, marble, and minor metachert; wm, ch, ab, ep, ac, ep, gt, st, ky, and ol; polydeformed and generally mylonitic; augen gneiss forms basketlike concordant sheets	sE; 136-Ma U-Pb age on sph from augen gneiss; 135.5-Ma Rb-Sr mineral isochron for augen gneiss and hb from schist and amphibolite; limited structural and metamorphic data suggest a possible M episode as well, synch. with deformation in the augen gneiss protolith	Mz; 342-Ma U-Pb age on zircon from augen gneiss which dates the intrusive age of the youngest known protolith	sE; 115-Ma U-Pb age on zircon from augen gneiss that intrudes this unit
Alaska Range north of the McKinley and Denali faults							
47	GNS (NS)	GNS (N)	YT, PN, MK	D and older; IB, in some areas, mid. and fel. plut., minor carb and clast(?)	Metabasite, gryite, phyl., calc-phyl., mafic schist, greenstone, metasands., metaarkf., metaconglomerate, metasands. M_2: metasomatic gneiss increases N-ward; ch, wm, ep, ac, ep, gt, st, ky, ol, and cc; M_3: ab, ch, ep, ser, ta, ca-amp, bi; Metadiorite, metagabbro, and possibly Triassic metasands. singly deformed and metamorphosed; other rock types polydeformed and mylonitic; M_2 assemblages overprinted metamorphic history of unit poorly known and may have differed in various parts	M_2(?); sK; 107-Ma K-Ar age on ms from phyl.	M_2: D age of youngest protolith known in here here metamorphosed	M_2: sK (in unit); tentative correlation with synchronous age of metamorphism of unit 46; 107(?)-Ma ms-mica schist (?)94-Ma(?) K-Ar age (?)90-Ma(?) age of amphibolite in unit 48. M_3: sK; 64-Ma K-Ar age (?)56-Ma on mica from unit 48; IK; age of crosscutting plutons in SE Mt. Hayes quad.
48	AMP (KS)	GNS (N)	YT	D and older; vol., seds., fel. plut.	Pelitic schist, gryite, amphibolite, gneiss; garnet: bt; ep, ts, mm, gt, ilm, YSW-ward increase in metamorphic grade; associated with M_2 and M_3; M_2: ab, ch, ep, bi, ac, gt; associated with NE-trending folds	----	D; age of youngest protolith	M_2; same ages as those given for unit 47; M_3; sK; oldest K-Ar age (?)56-Ma on mica from schist
49	LPP(GNS, (K) ?	?	YT	O(?)-M; seds., vol., carb.	Quartzose and quartzofeldspathic semischist, metabasite, metasands., carbonaceous schist, phyl., slate, metaconglomerate; wm, ch, ab, ep, gt, ep, ac; associated with NE-trending folds	sE(?); 110±3-Ma K-Ar whole-rock age on metarhyolite	M; age of youngest protolith	----
50	LPP(GNS (KS)?	?	PN, MK, GL, WY	mPk-mK; seds., vol., carb., marl. vol., chert	Phyl., metasands., metaconglomerate, metavol., metasiltstone; wm, ch, ab, ep, ac	----	mK; age of youngest protolith	sK; Paleocene age of overlying unmetamorphosed Cantwell Fm.
51	GNI-I (N) ?	?	ML	J-K; seds., int. and fel. vol., carb., gabbro	Phyl., -gnite, metaconglomerate, metavol., metasiltstone; greenstone; wm, ch, ab, ac, ep, ac	----	sE; proposed age of youngest protolith	sT; same evidence as for unit 50
52	AMG-I (N) ?	?	ML	K-K; seds., carb., coarse, plut., mal. igneous	Pelitic schist, calc-silicate, marble, gryite, amphibolite, weakly metamorphosed granitic rocks; E-ward increase in metamorphic grade from ch zone to bi, wm, ac, ep, gt zone; ab, ep, ch, ser, ta, ca-amp, bi, gt, st, ky, ol, wm, ac, ep, and gt; unit bounded near the SW contact, shown as a thrust fault	sE(?); 105-Ma $^{40}Ar/^{39}Ar$ plateau age on hb from synmetamorphic gneiss; 100-Ma Rb-Sr whole-rock isochron age for 18 samples of unit. U-Pb protolith ages of 1.8 Ga and older for weakly metamorphosed granite phases	----	----
53	GNS (KD)	YT	E(?)-Be(?); seds., vol.	Slate, phyl., semischist, -gnite, metaarkf.; wm, ch, ab, ep, ac	----	D; age of youngest known protolith in nearby units 47, 48 whose metamorphic history may, in part, correlate with that of this unit	K; minimum or actual age of metamorphic units 47, 48 whose metamorphic history may, in part, correlate with that of this unit
54	GNS (KD)	WY, MK?	Be(?), mD; seds., int. and maf. vol., carb.	Phyl., metaconglomerate, pelitic, calcareous, and mafic schist, gryite, metabasite, greenstone; wm, ts, ch, ab, ac, ep; tectonically flaked, mylonitic, and polydeformed by later kink bands	IK(?); 54- to 49-Ma K-Ar ages on synmetamorphose(?) plutons whose metamorphic effects overprint fabric and concordancy with bedding in gneiss is indicated; ages may be synmetamorphic	----	IK; age of plutons discussed in "Direct evidence..." column

Unit No.	Metamorphic unit [1,2]	Lithotectonic terrane(s) in which unit occurs[3]	Inclusive age range[4] and protolith[5]	Metamorphic rock type[6] key metamorphic mineral[7]; structural characteristics if relevant	Direct evidence for age of metamorphism[8] and evidence used[9]	Maximum age of metamorphism[10] and evidence used[11]	Minimum age of metamorphism[12] and evidence used[13]
AREA OF SOUTHERN ...LASKA BETW...EN THE McKIN...LEY AND DENALI FAULTS AND T...HE BORDER RANGES FAULT SYSTEM							
55	AMP (DH)		PE	B(?); fSt(?)-sMz(?); int. and maf. vol. and plut., seds., carb., chert	Meta-ols. and metaplut., especially amphibolite, metadiorite, and gneissose, quitte or metachert, pelitic and calcareous schist, marble, and minor metavol.; M1 and M2 occurs in pelitic schist from two localities on Alaska Peninsula, metamorphic fabrics range from imperfectly schistose to mineral-segregation gneissose; metamorphic grade increases toward E-ward; batholiths whose contacts and internal structures are discordant to that within this metamorphic unit	M_2; proposed age of metamorphism is probably younger than 65 Ma	TF-sD(?); 177- to 155-Ma K-Ar ages of batholiths intruding unmetamorphosed and non-sch associated with metamorphism
56	AMP (J) / GNS (PK)		PE	B(?)-J(?); cc(?) seds.	M_2: m-sJ(?); assumed correlation with metamorphism of unit 55. M_3: sK; correlation of ages with ok plutonism and possible metamorphism in unit 61 and (or) with K and eT metamorphism of units 72 and ?? of the Chugach terrane	----	eT; Paleocene age of overlying nonmetamorphosed plutons
57	GNS (J)		PE	fSt(?), sMz(?); int. and maf. vol., seds., carb., int. plut.	Ch. schist, greenstone, metaarch., marble, metagraywacke, meta-quartz diorite; ab, ch, ep, ab, ms; metamorphic grade increases S-ward; ky, st, gt, and ol in some areas; deeply dipping, steeply dipping, sylonitic schistosity locally developed parallel to trend of beds and major faults that bound unit	M_2; K; 1.77-Ga U-Pb age of epz from orthogneiss, K-Ar ages of 1.75 Ga on hb from gt amphibolite and Rb-Sr whole-rock isochron. M_3: 1.6; 1.56- to 1.50-Ma K-Ar ages on hb from this unit	K_2; age of protolith	M_2; Y; 1.2-Ga $^{40}Ar/^{39}Ar$ plateau age of hb from gt amphibolite. M_3: 1.6; 1.56; overlying unmetamorphosed plutons
58	GNS (MP) ?	GNS (Tk)	PE (more likely WS)	B-sMz(?); maf. vol., ---, seds., carb., chert, int. plut.	Metavol., metadiorite, amphibolite, metametadiorite, amphibolite, metasands., marble, lenses of ---, ch, ep, ab, ser, ta, ca-amp, bi, wm, epz; near horizontal E-W-striking stretching lineation in zone, vertical E-W-striking mylonitic foliation; metamorphism especially well developed in the Tk and Mt. Foraker plutons; ab, ep, ac, bi; (late phase) ch, ab, evolved from shelfs to brittle conditions	M_2; 187(?); 154-Ma $^{40}Ar/^{39}Ar$ plateau age of hb from metadiorite; 160-Ma ages either on hb or bi from intrusive or metamorphic age of the synmetamorphic Mt. Foraker pluton. M_3: 65-47; correlation of retrograde metamorphism with undeformed to slightly deformed Tk and Mt. Foraker plutons; M_3 assemblages in this area may have been produced during emplacement of an inferred large subsurface pluton	----	eT; Eocene age of post-metamorphic plutons
59	GNS (MP) ?	LPP (Tk)	WR	sW, P, (?)t, seds., vol. and plut., marble, chert	Schist, phyl., amphibolite, marble, metaarch., metavol., metachert, metaplut.; ab, ch, ep, ab, ser, ta, ca-amp, bi, gt, st, ky, ol, wm, ac, ep, and gt; metamorphic grade generally increases S-ward; ab, ch, ep, ab, ser, ta, ca-amp, bi, gt, st, ky, ol, wm, ac, ep, and gt; foliated, multiply deformed, and locally sheared and mylonitized	M_2; TR; age of metamorphism probably M_2; m(?)-J(?) dates of the Mt. Foraker pluton are affected by this	----	eT; Eocene age of post-metamorphic plutons in the region
60	GNS (AMP) ?		WR	P-; int. to maf. vol. and plut., minor oc. seds., chert	Greenschist, metagabbro, metasands., metaarkf., metaconglomerate; ab, ch, ep, ab, ms, ep, ac, bi, gt; fabrics of metagenesis rocks range from massive to gneissic and schistose; metamorphic rocks intensely folded and faulted	----	----	----
61	GNS/AMP (K?) ?		PE	P-(?); P, Mz(?); ---, seds., chert, maf. and int. vol., carb.	Argillite, metachert, quartzose or pelitic schist, amphibolite, calc-silicate rock, marble, blocks of metagraywacke and of foliated argillaceous matrix; ch, ep, bi, hz, ms, gt, epz; variable metamorphic grade; foliated rocks occur as discrete bodies, as slices in broken formation, and as rafted pendants of J-age plutons	e-mJ and (or) sK; age of plutons whose spatial relationships, age, and shared deformation with this unit may indicate that they were intruded during the waning stages of an as-sK and an eK synkinematic metamorphism episode	P; protolith age of marble	sK; age of youngest crosscutting plutons
62	AMP (TK)	?		B(?)-MzC(?); int. to maf. vol.(?), seds.	Metagraywacke and metaclastic rocks, amphibolite, schist, phyl.; hb, ch, gt, unit occurs within the Denali fault zone and probably is a slice far removed from the rest of the Yukon-Tanana terrane as part of the Wrangellia or Alexander terrane	----	Rx; probable age of youngest protolith	IT, 20- and 17-Ma K-Ar ages on crosscutting, unmetamorphosed plutons
63	GNS/AMP (TK)		PET	O(?)-fS; silicidentic and calcareous seds., maf. to int. plut. and vol., ultramafic rocks	Pelitic and calcareous schist, phyl., quitte, metabasite, metachert, mafic schist, metased., pyroxenite; ch, ens, ts, gt, ep, ac, ep, and ol; metamorphic grade increases S-ward; unit is a thick, fault-bounded, penetratively deformed, shallow-dipping, isoclinally folded sequence; structural contacts are faults; little lateral continuity due to tectonic mixing; degree of metamorphism (mostly greenschist facies with some blueschist facies in the Chelmic) differs between nearby fault blocks indicating postmetamorphic movement; ages of metamorphism unknown and adjacent to belt of IK and T batholiths	----	Rx; probable age of youngest protolith	IK-eT; metamorphism probably younger than or associated with(?) the intrusive ages of K- to Paleocene or late Eocene to Paleocene plutons
64	GNS (WS)		AXC	-(?)-D(?); ---, seds., carb., int. to maf. vol., int. plut.	Marble, mica schist, phyl., metaconglomerate, minor amphibolite, weakly metamorphosed diorite to gabbro; metamorphic grade of ms, ch, and rarely bi and gt; metamorphic grade decreases NE-ward; unit is a thick, fault-bounded, penetratively deformed sequence and are architecturally similar to belt and slip cleavage locally developed	IK; age of youngest protolith (of Mt. Foraker pluton); unit may be younger than 65 Ma	sK; age of metamorphic-plutonic	----
65	LPP (TK)		PE	fR; maf. vol., volcaniclastic seds.	Greenstone, metasands.; py, px	----	IK; age of protolith; 57(-6) of alteration in crosscutting 2 plutons that cut unit; 57-Ma age of metamorphism	eT; 60-Ma K-Ar age from metamorphosed plutons that intrude this unit
66	LPP (TK)	KH, CH, WF, BF, SU		Metagraywacke, semischist, argil., metachert, greenstone, metasandstone, volcaniclastic metasandst., metavol.; px, mx; metasands. show only incipient flattening of grains, but semischistose or protomylonitic fabrics present locally	m-IK; probable association of metamorphism (of at least Pyschocal rocks) with accretion of the Chugach terrane and forming for that event	----	----
67	LPP (TK)	?	KH	JK; Pyschocal seds., gabbro	Argil., metagraywacke, metasandstone, metaconglomerate, volcaniclastic metasands.; px, ps, ch, wm, ep, ac; primary depositional features generally preserved; duty cleavage in highest grade (N-most) zone; cleavage-related quartz veins; faults; unit forms lowest grade part of N-dipping inverted metamorphic sequence (Mackenzie metamorphic belt)	IK-eT; evidence consists of relations of plutons of the age to the Eocene Sanak batholith; 56-Ma U-Pb ages on epz and bt K-Ar ages on hb from gneissic phase of the batholith	----	----
68	LPP (TK)	?	KH	JK; Pyschocal seds., gabbro	Argil., metagraywacke, metasandstone, metaconglomerate, volcaniclastic metasands.; px, ps, ch, wm, ep, ac; primary depositional features generally preserved; duty cleavage in highest grade (N-most) zone; cleavage-related quartz veins; faults; unit forms lowest grade part of N-dipping inverted metamorphic sequence (Mackenzie metamorphic belt)	IK-eT; evidence consists of relations of plutons of the age to the Eocene Sanak batholith; 56-Ma U-Pb ages on epz and bt K-Ar ages on hb from gneissic phase of the batholith	----	----
69	GNI (TK)		ML, KH, NN	S-IC; Pyschocal seds., carb., marl. igneous	Phyl., metagraywacke, carbonaceous marble, quitte, metaconglomerate, metasandstone, metavol.; wm, ch, ab, ep, ac, and locally bi; penetrative deformation and metamorphism of unit S-ward increase in grade part of N-dipping, inverted metamorphic sequence (Mackenzie metamorphic belt)	----	----	----
70	AMI