

Base from U.S. Geological Survey, 1965
Geology generalized by MacKevett, 1976

Background information for this folio is published as U.S. Geological Survey Circular 739, available free of charge from the U.S. Geological Survey, Reston, Va. 22092.




Table showing linear correlation coefficients between logarithmic values of the concentration of selected elements versus arsenic, McCarthy quadrangle, Alaska. (Leads to -- indicate insignificant data.)

Table with 24 columns for elements (Fe, Mg, Ca, Ti, Mn, Ag, B, Ba, Be, Co, Cr, Cu, Mo, Nb, Ni, Pb, Sc, Sr, V, Y, Zn, Zr, Au, Cu, Pb, Zn, Hg, As) and 2 rows for Correlation Coefficient(X100) and Number of pairs.

Au, Cu, Pb and Zn by atomic absorption analysis
Hg by flameless atomic absorption analysis
As by colorimetric analysis

Table showing linear correlation coefficients between logarithmic values of the concentration of selected elements versus mercury, McCarthy quadrangle, Alaska. (Leads to -- indicate insignificant data.)

Table with 24 columns for elements (Fe, Mg, Ca, Ti, Mn, Ag, B, Ba, Be, Co, Cr, Cu, Mo, Nb, Ni, Pb, Sc, Sr, V, Y, Zn, Zr, Au, Cu, Pb, Zn, Hg, As) and 2 rows for Correlation Coefficient(X100) and Number of pairs.

Au, Cu, Pb, and Zn by atomic absorption analysis
Hg by flameless atomic absorption analysis
As by colorimetric analysis

DISTRIBUTION AND ABUNDANCE OF ARSENIC AND MERCURY IN STREAM SEDIMENTS AND MORAINIC DEBRIS, MCCARTHY QUADRANGLE, ALASKA




By Keith Robinson, R. M. O'Leary, C. M. McDougal, and Theodore Billings

DISCUSSION


A geochemical survey was conducted in the McCarthy quadrangle, Alaska, to identify areas containing anomalous concentrations of various metallic and nonmetallic elements. This study incorporated the results of geochemical and mercury from 426 and 424 stream sediment and glacial moraine debris samples collected in the McCarthy quadrangle, and analyzed by the U.S. Geological Survey between 1961 and 1976 using colorimetric and flameless atomic absorption spectrophotometry respectively. No analytical results for arsenic and mercury are available for stream sediment samples from the White River area, located in the northeastern part of the quadrangle.

Highly positive arsenic and mercury anomalies were detected in samples of stream sediments collected from the west slope of Bonanza Ridge (T. 4 S., R. 14 E.), and some weak arsenic and mercury anomalies in sediments from the McCarthy and Nikolai Creeks (T. 5 S., R. 15 E.). These anomalies may result from contamination by the Nikolai mines, respectively. It is significant, however, that arsenic and mercury seem to exist in association with rocks containing Kameo-type copper deposits. Stream sediments collected from the area west of Bonanza Ridge, extending to Castle Peak (T. 3 S., R. 11 E.), and southeast of Bonanza Ridge in the Nikolai Butte area (T. 6 S., R. 17 E.) also have anomalously high arsenic and mercury contents.

South and southeast of the University Peak (T. 6 S., R. 20 E.) in an area encompassing the Bartlett, Barnard, and Anderson Glaciers, located in the southeastern corner of the quadrangle, a large number of arsenic and a few mercury anomalies were detected in samples of glacial moraine and stream sediment. These samples originate in a region containing rocks of the Kaskawishak Group and some metamorphosed rocks of the Sojak Group that have been intruded by a monzonitic granitic complex of Pennsylvanian age.



Histogram showing frequency distribution of arsenic in stream sediments and glacial debris, McCarthy quadrangle, Alaska.



Histogram showing frequency distribution of mercury in stream sediments and glacial debris, McCarthy quadrangle, Alaska.

REFERENCES

Knabel, Jeff, 1970, Geochemical survey and geological reconnaissance of the White River area, southeastern Alaska: Alaska Div. Mines and Geology, Geol. Rept. 21, 60 p.