

PROPERTY OF DGGG LIBRARY

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

TO ACCOMPANY MAP MF-1480

STRUCTURE AND PETROLEUM POTENTIAL OF THE YAKUTAT SEGMENT OF THE NORTHERN GULF OF ALASKA CONTINENTAL MARGIN

By

Terry R. Bruns

ABSTRACT

Major structural features of the Yakutat segment of the continental margin, between Cross Sound and Icy Bay in the northern Gulf of Alaska, are delineated by multichannel seismic reflection data. A large structural high centered on Fairweather Ground lies at the edge of the shelf from Cross Sound to west of the Alsek Valley. A basement uplift, the Dangerous River zone, along which the seismic acoustic basement shallows by up to 2 km, extends north from the western edge of Fairweather Ground toward the mouth of the Dangerous River. The Dangerous River zone separates the Yakutat segment into two distinct subbasins. The eastern subbasin has a maximum sediment thickness of about 4 km, and the axis of the basin is near and parallel to the coast. Strata in this basin are largely of late Cenozoic age (Neogene and Quaternary) and approximately correlate with the onshore Yakataga Formation. The western subbasin has a maximum of at least 9 km of sediment, with about 4.5 km of Paleogene strata overlain by late Cenozoic strata. The Paleogene strata are truncated along the Dangerous River zone by a combination of erosion, faulting, and onlap onto the acoustic basement. Within the western subbasin, the late Cenozoic basin axis is near and parallel to the coast, but the Paleogene basin axis appears to trend northwest, diagonally across the shelf. Sedimentary strata throughout the Yakutat segment show regional subsidence and only minor deformation except in the vicinity of the Fairweather Ground structural high, near and along the Dangerous River zone, and at the shoreline near Lituya Bay.

Seismic data across the continental slope and adjacent deep ocean show truncation at the continental slope of the shelf Paleogene strata. Up to 6 km of undeformed or mildly deformed abyssal sedimentary section are present at the base of the slope, and in part onlap the slope. Faulting may have occurred along a relatively narrow zone along the slope or at the base of the slope. Observed deformation at the base of the slope is primarily related to the late Cenozoic uplift of Fairweather Ground and to Quaternary folding perpendicular to the Pacific-North America relative-convergence vector. No accretionary section or major deformation is observed along the continental slope. The absence of these features suggests that no major subduction of the Pacific plate beneath the Yakutat margin has occurred during the late Cenozoic. However, transform faulting has most likely occurred along the base of the slope, because probable Oligocene oceanic basement is juxtaposed against Mesozoic and Paleogene sedimentary strata of the Yakutat slope. This juxtaposition most likely occurred during late Oligocene and Miocene time. During much of the late Cenozoic, and especially during Pliocene and Pleistocene time, the Yakutat segment has apparently been moving northward with the Pacific plate.

Dredge samples from the continental slope recovered potential petroleum source and reservoir rocks from the Paleogene sedimentary sequences. Most of the organic matter from these samples is immature to marginally mature. However, if similar rocks are present beneath the shelf, they are likely to be thermally mature at a depth of 4 to 5 km and deeper. Thus, the Paleogene strata may have significant resource potential below the shelf. The Paleogene strata are contained primarily within the western subbasin; strata in the eastern subbasin appear to have little resource potential.

Structural traps are apparently present in parts of the basin near and along the Dangerous River zone. These traps are updip from potentially mature strata of the western subbasin, and could hold commercial accumulations of hydrocarbons, if sufficient hydrocarbon generation and migration has occurred to fill such traps and if sufficient reservoir rock is present.

INTRODUCTION

This report discusses the structure, geologic history, and petroleum potential of the Yakutat segment, the part of the continental margin between Cross Sound and Icy Bay, northern Gulf of Alaska (figs. 1, 2, and 3). As part of a program of geological and geophysical investigations of the continental margin in the northern Gulf of Alaska, the U.S. Geological Survey collected multichannel seismic reflection data along about 2,000 km of tracklines in the study area during 1975, 1977, and 1978 (fig. 3). In addition, dredge samples from the continental slope were acquired during the 1977, 1978, and 1979 field seasons. Geologic results of the dredge cruises are reported by Plafker and others (1978c, 1979, 1980). The first part of this paper presents an interpretation of the seismic reflection and refraction data, including structure contour maps, isopach maps, and interpreted seismic sections; the second part is a discussion of the implications for petroleum potential. The primary area of interest is the continental shelf and slope, but some data from strata at the base of the slope are also included. Some of these data were presented in previous reports by Plafker and others (1978a), Bruns (1979), and Bruns and Plafker (1982). The seismic data for the 1975 cruise are reported by Bruns and Bayer (1977).

REGIONAL SETTING

The Yakutat segment of the continental margin lies in a zone of transition between transform and convergent motion (Rogers, 1977; Plafker and others, 1978b; von Huene and others, 1979; Bruns, 1979; Perez and Jacob, 1980; Lahr and Plafker, 1980). To the east of the study area, the Pacific plate is moving laterally past the North America plate along the Queen Charlotte-Fairweather fault system (fig. 2). To the west, in the Kodiak Island and Alaska Peninsula regions, plate motion is accommodated by underthrusting of the Pacific plate beneath the continental margin, resulting in the formation of the Aleutian Trench and volcanic arc. The northernmost extent of compressional deformation associated with this convergence lies onshore in the complex folds and faults of the Yakataga and Malaspina districts and offshore within the structures of the Yakataga segment of the continental margin along what has been informally termed the Pamplona zone (Plafker, 1971; Plafker and others, 1978b, Bruns, 1979; Lahr and others, 1980; Bruns and Schwab, 1983). The Yakutat segment is part of a small crustal plate, informally termed the Yakutat block (Rogers, 1977; Plafker and others, 1978b; von Huene and others (1979), Bruns, 1979; Lahr and Plafker 1980; Bruns and Schwab, 1983), that is bounded by Kayak Island on the west, the Queen Charlotte-Fairweather fault system on the north and northeast, and the base of the continental slope on the south (fig. 2).

REGIONAL GEOLOGY

The study area is bordered onshore by rocks ranging from Mesozoic through Cenozoic age that decrease in age seaward (fig. 2). Rocks of Mesozoic and older age are present north of the study area in the high Chugach, St. Elias and Fairweather mountains, and are exposed on the seaward islands of southeastern Alaska. These rocks are characteristically highly deformed, locally metamorphosed and intruded sedimentary and volcanic rocks (Stoneley, 1967; Plafker 1967, 1971; Plafker and others 1973, 1978a; Belkman, 1980).

Cenozoic rocks onshore consist of bedded terrigenous clastic sedimentary rocks that were deposited primarily in a marine environment. The rocks are broadly divisible into three sequences that correspond to major changes in the tectonic and depositional environment of the basin. (1) A lower Tertiary sequence (Paleocene through early Oligocene) is characteristically hard, dense, and variably deformed and faulted; it is composed of deep-sea fan deposits and associated volcanic rocks of Paleocene age, and continental to shallow-marine coal-bearing clastic rocks of Eocene and early Oligocene age. (2) A middle Tertiary sequence (middle Oligocene through early Miocene) is composed primarily of mudstone and siltstone. (3) An upper Tertiary (Miocene through Holocene) sequence, the Yakataga Formation, is a marine diamictite characterized by abundant glacial detritus and composed of mudstone, muddy sandstone and conglomeratic sandy mudstone (Plafker, 1971; Plafker and Addicott, 1976; Plafker and others, 1975, 1978a; and Winkler, 1976). The Yakataga Formation reflects deposition adjacent to the intensely glaciated, high St. Elias and Fairweather Mountains, and deposition of these strata probably correlates to uplift of these mountains (Plafker and Addicott, 1978).

Offshore, strata of Mesozoic through Pliocene age that underlie the continental slope of the Yakutat segment have been sampled by dredging (sheets 1 and 2; Plafker and others, 1978c, 1979, 1980). These strata can be generalized in much the same way as the onshore strata from oldest to youngest as follows (unit designations are from Plafker and others, 1980): (1) An inferred Upper Cretaceous sequence consists of hard graywacke, argillite and possibly intrusive rocks (unit A). This sequence is found on the continental slope off Fairweather Ground and probably underlies much of Fairweather Ground. (2) A lower Tertiary sequence consists of upper Paleocene(?) and lower Eocene(?) sandstone, conglomerate, and shale (unit B), lower Eocene basalt flows and pyroclastic rocks (unit C), lower and middle Eocene sandstone, conglomerate, siltstone, and shale (unit D), and upper Eocene and lower Oligocene(?) shale, tuffaceous shale, siltstone, and sandstone (unit E). This sequence is found along parts of the continental slope from Fairweather Ground to about Pamplona Spur. (3) A middle Tertiary sequence consisting of upper Oligocene siltstone (unit F) crops out locally along the upper slope. (4) A Miocene and younger clastic sedimentary sequence includes abundant glaciomarine deposits (unit G), approximately equivalent to the onshore Yakataga Formation. The Paleogene sedimentary strata (units B and D through F) are probably continentally derived, with younger rocks deposited in progressively shallower water. In addition, these rocks differ strikingly in lithology from coeval rocks exposed in outcrops onshore or penetrated in exploratory wells onshore or offshore (Plafker and others, 1980).

GEOPHYSICAL DATA

This study is based primarily on 24-fold multichannel seismic-reflection data gathered in 1975 from the Geophysical Services, Inc. vessel M/V Cecil H. Green under contract to the U.S. Geological Survey (Bruns and Bayer,

1977), and in 1977 and 1978 from the U.S. Geological Survey research vessel R/V S.P. Lee. Tracklines for these three cruises are shown on sheets 1 and 2, and in figure 3. Single-channel seismic-reflection data across the continental slope south of Icy Bay (von Huene and others, 1975) provide subsurface information between the multichannel lines.

The seismic system on the M/V Cecil H. Green consisted of a tuned array of 22 air guns with a total capacity of 19.6 liters (1,200 in.³), a 2,400-m 48-group streamer and DSF IV¹ digital-recording instruments. These data were processed by Petty-Ray Geophysical, Inc. (Bruns and Bayer, 1977). The seismic system on the R/V S. P. Lee consisted of a tuned array of five air guns with a capacity of 21.7 liters (1,328 in.³), a 2,400-m 24-group streamer, and GUS 4300 digital-recording instruments. These data were processed by the U.S. Geological Survey. In all surveys, navigation was by means of an integrated satellite, Loran C, and doppler-sonar navigation system.

Bathymetric, gravity, and magnetic data were also acquired during these cruises (Schwab and Bruns, 1979; Schwab and others, 1980; Burkhard and others, 1980a, b; Atwood and others, 1981; and Bruns and others, 1981a, b).

GEOPHYSICAL INTERPRETATION

Shelf and Slope

Contour and isopach maps of strata beneath the continental shelf and slope (sheets 1 and 2) are based on mapping of seismic horizons D and F throughout the seismic grid. Horizon D correlates with the D horizon in the Yakataga segment to the west (Bruns and Schwab, 1983). An additional horizon, horizon E, is mapped along parts of a few seismic lines. These horizons are all at least local unconformities and strong seismic reflectors. In areas where the strata are conformable, mapping is on a seismic horizon interpreted as correlative with the unconformities. Since seismic mapping is on unconformities or the correlative conformity, the mapped horizons have time-stratigraphic significance. Interpretation problems in following the seismic horizons are created by the wide spacing of the data, by poor reflector continuity, and by limited acoustic penetration, especially in the deep parts of the basin. Several factors that limit acoustic penetration and disrupt reflector continuity can be identified or inferred at least locally. These factors include persistent water-bottom and interbed multiples that obscure primary reflectors, coarse or bouldery near-surface and subsurface glacially derived sediments that may scatter or limit penetration of the acoustic signal, and local unconformities and rapid facies or bedding changes within the Yakataga Formation. Because of these problems, and because of the wide spacing of the seismic data, correlation errors may be present in some areas; however, these errors should not seriously change the general basin form shown in the structural contour maps.

The character of the mapped horizons and the seismic reflectors between these horizons is best seen on the slope portions of the seismic sections (sheet 3). The mapped seismic horizons and the reflection character between these horizons are discussed from shallowest to deepest in the following section. The seismic sequence, strata, or time interval between any two horizons is referred to by the two horizon letters; for example strata between horizons F and D is the F-D strata. The seismic sections presented in this report (fig. 3) are shown on sheet three.

Horizon D is mapped on a prominent continuous reflector or group of reflectors best observed on the parts of the seismic lines extending across the continental slope south of Icy Bay (seismic sections 404, 916, 906; sheet 3) and on lines near and over Fairweather Ground (seismic sections 403, 911, 400; sheet 3). The horizon is moderately well defined

¹Use of trade names in this report is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey.

throughout the seismic grid, except on the landward ends of the seismic sections where the sedimentary section is thickest.

The relation of seismic reflectors between horizon D and the sea floor (D-SF reflectors) to horizon D appears to be primarily one of concordance. However, on the gently dipping continental slope south of Icy Bay, strata immediately above horizon D show very gentle downlap onto the D reflector, indicating a prograding slope depositional environment (seismic sections 404, 916, 906; sheet 3). Onlap of the D-SF reflections onto the horizon D reflector is also seen on seismic sections over Fairweather Ground (seismic section 400, sheet 3). On the steep continental slope between Yakobl Valley and Yakutat Valley, strata of the D-SF sequence are very thin to absent (seismic sections 403, 909; sheet 3).

In general, seismic reflectors between horizon D and the sea floor show variable continuity and are parallel or slightly divergent. Well-defined high-amplitude reflections within the sequence are separated both vertically and laterally by low-amplitude to almost reflection-free zones. The reflector configuration generally suggests discontinuous lithologic units and a variable lithology. The D-SF sequence thickens landward from the shelf edge, indicating regional subsidence of the continental shelf (seismic sections 404, 906, 403, 909, 400; sheet 3).

Horizon E is mapped on a prominent, local unconformity on the seaward part of seismic section 403 (sheet 3), but the unconformity cannot be traced with confidence landward or into adjacent seismic lines. Reflectors in the E-D sequence are parallel or subparallel to reflectors in the overlying D-SF sequence. On the seaward part of the line, E-D reflectors onlap horizon E at the base of the sequence; elsewhere these reflectors are concordant with horizon E.

Horizon F is mapped on a strong seismic reflector that is acoustic basement on the seismic lines. The acoustic basement is best seen on section 403 and lines to the east (sheet 3), and can only be discontinuously mapped west of line 403. The acoustic basement is seen only on the slope portion of lines west of line 906, most likely because the acoustic energy was insufficient to penetrate through the thick sedimentary sequence in the area south of Icy Bay. Refraction data (Bayer and others, 1978) give an estimate of total sediment thickness in this area and are used to infer the approximate position of horizon F contours (sheet 1). Age assignments based on foraminiferal data from the Colorado Oil and Gas wells near the town of Yakutat (wells 83-85, sheets 1 and 2; Rau and others, 1977) indicate the depth to strata correlated to horizons F and D, and are used to estimate the position of the horizons in the area of Yakutat Bay. East of line 909, horizons F and D merge, as the F-D reflectors are truncated against a structural high.

Seismic reflections below the D horizon can best be seen on the continental slope segments of seismic sections 404, 916, 906, and 403 (sheet 3); beneath the shelf, the reflectors are partly or totally obscured by multiples generated within the overlying section. Where best seen, reflections in the upper part of the F-D sequence show moderate to poor continuity, and in some places, the sequence is almost reflection free. Reflections are divergent with seaward thickening (sections 906, 403; sheet 3). The reflection configuration suggests a well-bedded, uniform lithology. On section 906 the lower part of the sequence is markedly different, with reflections showing good continuity and high amplitude; this difference may reflect the complex character of the basement unit described in the following section on geologic correlation of the seismic horizons.

The character of reflections below horizon D suggests two interpretation problems. First, these reflections are low amplitude and, beneath the continental shelf, are usually obscured by multiple reflections. Therefore, the stratigraphic character and internal structure of the F-D strata cannot be well determined even though the top and bottom of the F-D sequence are moderately well defined. Second, the lower part of the F-D sequence on the seaward end of seismic section 906 shows a set of strong reflectors which thicken seaward. The same set of reflectors beneath the shelf would likely not be resolvable, except for the initial

reflector at the top of the strong reflector set. Thus, the "acoustic basement" picked on the seismic lines may exclude some strata, and the resulting sediment thickness must be considered a minimum.

Base of Slope

Seismic data indicate that a well bedded sequence of sedimentary strata is present at the base of the slope (sections 404, 906, 403, 909, 967, 400; sheet 3). Three horizons are mapped and designated as horizons A1, A2, and A3. A correlation of multichannel seismic data to Deep Sea Drilling Project (DSDP) hole 178 east of Kodiak Island indicates correlations of basal Pleistocene(?), basal Pliocene(?), and top oceanic basalt respectively for the horizons; the ages are queried due to the distance to the drill hole. Horizon A3 (oceanic basalt) is characterized by strong, moderately discontinuous, hyperbolic reflectors. Where sediment overlying the basalt is thick, the reflector is fainter than in other areas and more continuous. The reflection configuration between horizons A3 and A1 is highly variable and in part area dependent, ranging from faint and discontinuous, to strong and continuous. In general, reflectors are parallel to subparallel throughout the interval. This section may include both pelagic units and varying amounts of turbidite deposits. The reflection configuration above horizon A1 is commonly complex and variable, and includes wavy or mounded configurations with low to high reflector continuity. The reflections are suggestive of turbidite fans, with surface and buried channels and levees common. Near the base of the slope, particularly from Yakutat Valley to the west, the sequence is characterized by chaotic reflectors, suggesting rapid deposition.

VELOCITY DATA

To make structural and isopach maps, seismic time was converted to depth using two types of velocity information: stacking velocities obtained during processing of the multichannel reflection data, and velocities from seismic refraction data. An area-average stacking-velocity curve was derived from all the stacking-velocity picks from individual seismic lines. Good velocity picks are concentrated in the upper 2 to 3 seconds of the seismic data, in general, corresponding to the data above horizon D. Thus, the stacking-velocity picks give good velocity estimates for the upper 2 to 3 seconds, but only poor velocity estimates below about 3 seconds.

Refraction data helps constrain the velocity in the deeper sedimentary section. Four unreversed-refraction sonobuoys were obtained during the 1977 cruise, using the air gun as a source. All four sonobuoys are in areas of gentle or no dip (fig. 3). Two of the sonobuoys (3 and 6) are over an area of relatively shallow acoustic basement, and the basement picks were excluded from the calculation for an average time-depth curve. The other two sonobuoys (4 and 5) are over an area with a thick sedimentary sequence (fig. 4).

The area-average stacking-velocity curve was converted to interval velocities by use of the Dix formula (Dix, 1955), and the interval curve was used to construct a time-depth curve. This curve was combined with the refraction velocity data, and fitted with a least-squared-error routine to yield a single curve, $z = + 0.895t + 0.296t^2 - 0.020t^3$, where z is the depth in kilometers and t is the two-way traveltime in seconds (fig. 5). This curve was used for all time-to-depth conversions. An approximate error range of the resulting curve (fig. 5) was derived by estimating a maximum and minimum limit for the stacking velocities; these limits were then converted to time-depth picks in the manner given above.

The final area-average time-depth curve is very similar to an independently derived curve developed for the Yakataga shelf to the west (Bruns and Schwab, 1983), and closely fits velocity data derived from sonic logs of the Standard Oil Company of California Rios Bay No. 1 well in Icy Bay (well 58, sheets 1 and 2), and the Continental Offshore Stratigraphic Test No. 1 well, drilled southeast of Icy Bay on the continental shelf (not shown; Boim and others,

1976; Bruns, 1979; Bruns and Schwab, 1983). Both of these wells are in structurally simple areas. The method of derivation of the curve gives a good depth estimate for the sedimentary section underlying the continental shelf, approximately the section above horizon F, but it is not valid for depth conversion below horizon F or for times greater than about 5 seconds two-way reflection time. In areas of deep water, examination of stacking-velocity data indicates that the time-depth curve applied from the seafloor is almost identical with the curve developed for the shelf. Using a single velocity function introduces errors in the depth conversions of the contoured horizons and of the seismic lines, but the errors should not affect the basic conclusions of this report.

GEOLOGIC CORRELATION OF SEISMIC HORIZONS

The correlation of the mapped horizons with the regional geology (fig. 6) is determined primarily from dredge data from the continental slope (Plafker and others, 1980). Horizon D is approximately correlative with the base of unit G of Plafker and others (1980), with correlation of the seismic data best defined on seismic sections 404 and 908 (sheet 3). Unit G is in turn correlative with the onshore Yakataga Formation. The horizon can also be correlated with the Riou Bay well at Icy Bay (well 58, sheets 1 and 2), where the horizon is approximately at the base of the Yakataga Formation (Bruns and Schwab, in press).

Onshore well data (Rau and others, 1977) suggest that much or most of the Yakataga Formation within the study area could be as young as Pliocene and Pleistocene. In the Riou Bay well, the lower part of the drilled section is Pliocene(?). In the Malaspina wells (wells 81 and 62, sheets 1 and 2), the Yakataga Formation is of Pliocene and younger age and overlies upper Eocene(?) strata. In three Colorado Oil and Gas wells near Yakutat (wells 63, 64 and 65, sheets 1 and 2), the lower Tertiary section is truncated updip and the Yakataga Formation unconformably overlies Eocene (Narizian) strata in the two seaward wells, and Cretaceous strata in the landward well. Finally, most of the Yakataga Formation at Lituya Bay is of Pliocene and younger age. The Yakataga Formation was not extensively sampled during the dredging cruises, so the age correlation of the offshore strata (D-SF strata) with the onshore Yakataga Formation is not known, but may be similar to ages known in the wells. Thus, as a possible end-member case for horizon D, it could be a major time-stratigraphic break and an erosional unconformity, with Oligocene and older strata below horizon D, and Pliocene and younger strata above the horizon. Alternatively, at the other extreme, the section across horizon D could be continuous with no major breaks in deposition. Both of these cases will be considered later in the report. It is likely that horizon D is intermediate between these cases, with the strata above the horizon comprising a seaward prograding sedimentary sequence, and with the magnitude of the unconformity diminishing toward the coast.

Horizon E is correlated with the dredge data on and near section 403 (sheet 3). On section 403, the horizon is a prominent unconformity, separating dipping F-E reflectors from overlying more gently dipping E-D reflectors. The horizon is therefore approximately of middle Oligocene age. The unconformity is not well defined in adjacent seismic lines, and strata appear to be conformable across the horizon over much of the shelf. Dredge samples from the continental slope suggest lower or middle Oligocene strata are unconformably overlain by upper Oligocene strata in the vicinity of line 403 (Plafker and others, 1980). These samples reflect what may be the greatest age difference across horizon E, since they were dredged in an area where uplifted strata in the F-E sequence are overlain by flat-lying, overlapping strata in the upper part of the E-D sequence (section 403). The observed unconformity is interpreted to reflect local deformation of F-E strata during about middle Oligocene time, followed by onlap of E-D strata during late Oligocene time.

Horizon F is best correlated with the dredge data on seismic sections 908 and 403 (sheet 3). On these lines the horizon is correlative with the top of unit C of Plafker and

others (1980), a basalt unit of early Eocene age. Thus, in the area south of Yakutat Bay, the section between horizons F and D includes strata of middle and late Eocene and early and middle Oligocene(?) age (units D, E, and F of Plafker and others, 1980).

Horizon F is not correlative everywhere with the top of the basalt. East of line 403, the section between seismic horizons F and D is truncated in the vicinity of the Alsek Canyon against a basement high, termed the Dangerous River zone in this report (figs. 4-7). Thus, in the area east of the zone, and generally bounded by the shoreline and Fairweather Ground, horizons F and D coincide. Horizon F in this area coincides at least in part with the Cretaceous strata sampled on Fairweather Ground, although a lower Tertiary section may still be present, but not distinguishable on the seismic sections.

In the area west of line 908, primarily in the vicinity of line 404 (fig. 3), the relation of seismic horizon F with the dredge data is not clear, as the acoustic basement cannot be clearly mapped beneath the continental shelf. The seismic-refraction data of Bayer and others (1978) give an approximate position for the acoustic basement, based on a change of refraction velocity from around 4 km/s to 7 km/s. If horizon F is simply extended seaward, the resulting interpretation is that geologic units B and C lie above horizon F. This conclusion is considered unlikely because (1) the units elsewhere form acoustic basement and (2) this conclusion would require uplift and truncation of units D, E, and F, and probably much of unit C below the shelf and upper slope between lines 403 and 404, whereas the seismic data do not suggest that such massive truncation occurs. An alternative interpretation is that a major fault is present in the lower Tertiary strata along the lower slope and that a wedge of units B and C has been uplifted seaward of the fault. A further study of this interpretation using modeling studies of gravity and magnetic data is in progress.

In summary (fig. 6) horizon D is correlated with the base of unit G of Plafker and others (1980). It corresponds to the unconformity at the base of the Yakataga Formation, which in the study area separates Oligocene from Miocene(?), Pliocene, and younger strata. Horizon E appears to separate units E and F of Plafker and others (1980) and is approximately middle Oligocene in age. The horizon is mappable only in the vicinity of line 403 and cannot be mapped throughout the rest of the shelf area. The geologic significance of horizon F is variable. East of the Dangerous River zone, horizon F largely corresponds to horizon D and is on the top of probable unit A strata, approximately equivalent to the top of the Jurassic(?) and Cretaceous Yakutat Group. To the west, horizon F is at the top of unit C, an Eocene basalt. In either case, the depth to horizon F gives a minimum thickness for Cenozoic sedimentary rocks for the shelf and upper slope; additional sedimentary strata could be present within or below units C and B.

SHELF STRUCTURE

The structure of the Yakutat segment of the continental shelf (sheets 1 and 2) is characterized by three major elements: (1) a large structural high at the shelf edge centered on Fairweather Ground, (2) a basement uplift, referred to herein as the Dangerous River zone, extending north from the western edge of Fairweather Ground toward the mouth of the Dangerous River, and (3) two sedimentary basins, separated by the Dangerous River zone. These features, and seismic data across these features, are discussed below.

Fairweather Ground

The Fairweather Ground high lies at the edge of the continental shelf, roughly between Alsek Valley and Yakobi Valley (sheets 1 and 2). Dredge hauls (Plafker and others, 1980) and magnetic data (Schwab and others, 1980) suggest that the core of the high and the eastern part of the upper continental slope south of Fairweather Ground consist of pre-Tertiary rocks typical of the Yakutat Group on the adjacent mainland (unit A of Plafker and others, 1980). Tertiary rocks

of Eocene through Oligocene age (units C, D and E of Plafker and others, 1980) are present along the western part of the upper continental slope bordering the high (sheets 1 and 2). The lower Tertiary strata are also present in the subsurface on the western and northwestern flanks of the high (sections 909, 911, 912; sheet 3). Deep-sea deposits, probably hemipelagic sediments and turbidites, onlap the lower continental slope seaward of the high (sections 967, 400; sheet 3). Sedimentary rocks of late Cenozoic age similar to the Yakutat Formation (unit G of Plafker and others, 1980) occur in isolated basins on the high (Plafker and others, 1980).

On the landward side of Fairweather Ground, seismic data show that Neogene and Quaternary sedimentary strata (D-SF strata) onlap the high and dip toward the coast into the eastern subbasin of the Yakutat shelf. These strata may have surrounded and covered the older rocks, and are uplifted and truncated at the sea floor over at least part of the high (sections 909, 911 and 912; sheet 3). Sections 909 and 911 (sheet 3) and depth sections of these lines (figs. 7 and 8) show that much of the uplift occurred late during the D-SF time interval. Little or no thinning of strata is evident in F-D strata or in the lower part of the D-SF strata along these lines, whereas more pronounced thinning is present in the upper part of the D-SF sequence. On section 911 (sheet 3 and fig. 8) the D-SF sequence can be divided into two subsequences at horizon U. Strata below horizon U show only minor seaward thinning, but strata above horizon U show much greater seaward thinning. Thus, horizon U marks a change from no or relatively slow uplift of Fairweather Ground to more rapid uplift. The amount of uplift can be estimated by flattening on horizon U, resulting in the restored section of figure 8. In this section, strata are flat or dip gently seaward, suggesting that little uplift occurred prior to horizon U time. The minimum amount of uplift is estimated from the difference between the restored depth to horizon F and the current depth to the horizon, indicating uplift slightly less than 1 km along section 911. A similar study along section 909 suggests uplift of about 2 km (fig. 7). Uplift may be greater because a substantial thickness of D-SF strata may have been eroded from the top of the high, and the original dip of horizon U may have been slightly seaward; the uplift estimate does not include the difference between horizon U and the sea level at horizon U time.

Dangerous River Zone

The Dangerous River zone is an area where the acoustic basement on the seismic data becomes markedly shallower, with structural relief on the acoustic basement horizon of 2 km or more (sheets 1 and 2). Seismic data show an abrupt geologic change across the zone. The thick Paleogene section present in the area west of the zone is truncated at horizon D; the section onlaps and is in part truncated by faulting against acoustic basement along the trend of the zone (lines 903, 909, 911, 912, 913, 914; sheet 3). Paleogene strata are either thin or absent east of the zone, or have markedly different acoustic properties and form the acoustic basement. The southern extension of the Dangerous River zone cannot be seismically mapped across the Fairweather Ground high. However, the zone trends into an area on the continental slope where, on the basis of dredge data, Paleogene strata are in close proximity to pre-Tertiary strata and may be in fault contact (sheets 1 and 2; also see fig. 4 of Plafker and others, 1980). The north or northwest extension of the zone is inferred to pass beneath the Colorado Oil and Gas Yakutat wells (wells 63-65, sheets 1 and 2), where a thick Paleogene section is cut out by truncation or faulting (Rau and others, 1977).

The character of the structural and stratigraphic changes across the Dangerous River zone is variable. Along sections 903 and 912 (sheet 3) the Paleogene section appears to thin primarily by erosion at about horizon D time, although some onlap is also apparent. On lines 911 and 913 (sheet 3 and fig. 7), the section onlaps the acoustic basement. Finally, on line 909 (figs. 13, 23), the Paleogene strata both onlap the gently dipping acoustic basement (horizon F), and in part buttress against faulted strata of the acoustic basement (below location 1, fig. 7). Some additional minor deformation

is also present in the early Tertiary section and extends into the upper part of the section (below location 2, fig. 7). Strata do not clearly thin landward, as might be expected if faulting occurred prior to or during deposition, and some of the strata in the upper part of the F-D sequence are unfaulted over the fault zone. The structure along the Dangerous River zone may be largely a result of uplift prior to Eocene time, with Eocene and Oligocene strata onlapping a paleoslope formed at the fault zone. Strata beneath the F/D horizon east of the Dangerous River zone must be largely of Cretaceous through early Eocene age, since middle and upper Eocene strata onlap the acoustic basement horizon.

The Dangerous River zone marks the subsurface expression of the Paleogene basin edge that Plafker and others (1980) delineated on the continental slope from dredge data. Initial development of the basin started prior to horizon F time. Seismic reflectors in the lower part of the F-D sequence onlap horizon F. Foraminifers from dredged rocks of geologic units D and E show decreasing water depths, from middle to upper bathyal and outer-shelf depths for unit E. Further, considerable transport of shallow-marine sands into deeper marine sediments occurred in unit D (Plafker and others, 1980). Thus, seismic and geologic data indicate deposition into a preexisting structural low in approximately early and middle Eocene time and infilling of the low during middle and late Eocene and early Oligocene time. Faulting along the zone occurred largely prior to the early stages of deposition of the F-D sequence. Additional deformation occurred during late Oligocene and early Miocene time and is discussed below.

Sedimentary Subbasins

The Dangerous River zone separates the Yakutat shelf into two distinct subbasins (sheets 1 and 2). The eastern subbasin is bounded by the shoreline, the Dangerous River zone, the Fairweather Ground high, and the offshore extension of the Fairweather fault system. Strata in this basin that are resolvable on the seismic data are of late Cenozoic age, and the lower Tertiary strata are missing, thin, or form acoustic basement. On the isopach map of upper Cenozoic strata in the eastern subbasin the isopach contours virtually coincide with the structure contours.

The upper Cenozoic section onlaps the Fairweather Ground high and is uplifted and truncated at the sea floor along at least parts of the high. The sedimentary section dips toward the coast, with a maximum thickness of about 4 km east of Dry Bay. The axis of the basin is near and parallels the coast. Marked uplift and folding of the basin strata are present along the shoreline of the onshore Lituya Bay district, where the Yakutat and Topsy Formations crop out with almost vertical dip (Plafker, 1967, 1971). Offshore seismic data indicate flat-lying sediment within 3 km of the coast near Lituya Bay, suggesting that a fault or sharp fold with significant deformation and vertical displacement lies near the shoreline. High-resolution seismic data (von Huene and others, 1979; Carlson and others, 1979), and section 400 (sheet 3) indicate that this fault extends to the southeast where it merges with the offshore Fairweather fault system, and it could be a strike-slip fault. The fault may extend onshore to the northwest beneath the thick alluvium of the Yakutat district either along or near the shoreline, or trending into upper Yakutat Bay. In general, the upper Cenozoic strata within the eastern subbasin show only regional dip toward the basin axis and are deformed only along the fault zone near Lituya Bay and its southern extension, and over or around the Fairweather Ground high.

The western subbasin lies between the Dangerous River zone and the compressional folds of the Pamplona zone (sheets 1 and 2). The structure and isopach maps show that the Tertiary sedimentary section thickens markedly west of the Dangerous River zone to more than 9 km south of Joy Bay and includes a thick Paleogene section (F-D strata, sheets 1 and 2). South of Yakutat Bay, roughly between the shelf break and the coast, this Paleogene section has a maximum thickness greater than 5 km. In the western part of the subbasin, the thickness and extent of this section is not well defined, since seismic reflection and refraction data give only

an approximate position for horizon F, but the section is at least 4.5 km thick. The thickest part of the Paleogene section, as seen in the isopach map (sheet 2), trends roughly northwest and may define the axis of the early Tertiary basin. The section is truncated at the continental slope and thins out over and onlaps the acoustic basement along the Dangerous River zone.

Strata in the Paleogene section were at least locally deformed prior to horizon D time. On seismic section 403 (sheet 3), uplift of strata is evident at what is now the shelf edge (location 1) and a gentle anticline is present in Paleogene strata in the center of the line (location 3). A depth section of this line (fig. 9), with reconstructed flattening on horizons D, E and an intermediate U horizon, suggests the following history. (1) Strata were initially deposited into an existing basin or onto a subsiding platform, with minor uplift at location 1 prior to U time (fig. 9). Paleontologic data from dredged rocks from the F-D sequence suggest that the sediments were deposited in a middle to upper bathyal environments, and that shallow marine sediments were transported into an existing basin (Plafker and others, 1980). (2) Between horizon E and horizon U times, differential subsidence resulted in the formation of a local structural high (location 3, fig. 9). Prior to horizon D time, uplift of about 1 km at location 1 truncated part of the E-D strata. (3) Finally, deposition of D-SF strata and regional subsidence buried the Paleogene section. Uplift again occurred during the middle and late D-SF time interval, as strata from this sequence are truncated at the sea floor near the shelf edge. The seismic data suggest that major erosion did not occur at horizon E time, since no marked truncation of seismic reflectors in the F-E sequence is seen at horizon E in the deformed zones on section 403.

Strata in the F-E sequence are unconformably overlain by strata in the E-D sequence. The age difference across horizon E is variable, as successively younger strata in the E-D sequence onlap horizon E on both the anticline and on the uplifted strata at the shelf edge. The angular unconformity may not be present in other areas of the shelf, as little structure is seen in F-D strata except along section 403 and at the shelf edge between Alsek Valley and Yakutat Valley. Seismic data are insufficient to define the extent of the anticline seen on section 403, but blocks leased during OCS Sale 55, 1980, presumably are on this high and provide some indication of its extent and the location of some nearby smaller structures (fig. 3).

The structure and isopach maps of strata above horizon D (D-SF strata, sheet 2) show that the depositional axis of the upper Cenozoic strata in the western subbasin trends east to west and lies near the coast. The section increases uniformly in thickness from the shelf edge to the basin axis, with a thickness greater than 5.5 km in the deepest part of the basin. These strata are relatively continuous across the Dangerous River zone into the eastern subbasin, with faulting and folding seen nearshore only along the northern extent of the zone (section 914, sheet 3). This deformation may be a result of reactivation of faulting along the Dangerous River zone, or could be due to the westward continuation of the fault off Lituya Bay.

Adjacent to the structures of the Pamplona zone, the structural contours on horizon D show that the D-SF strata are bowed down into a structural low beneath the continental slope. However, the isopach map shows that the section maintains the same general strike and uniform increase in thickness toward the basin axis as is seen in the rest of the subbasin, instead of showing an increase in thickness in the structural low. Thus, the upper Cenozoic section in this area, rather than being deposited into a preexisting structural low, is apparently being bowed down in response to compressional deformation across the Pamplona zone. Elsewhere in the western subbasin the D-SF strata show regional subsidence toward the basin axis, but are otherwise undeformed.

SLOPE AND BASE OF SLOPE STRUCTURE

Seismic data across the continental slope and adjacent deep ocean show three major features. (1) The lower Tertiary strata (F-D strata) are truncated at the continental slope, and

strata above horizon D (D-SF strata) in part prograde seaward over the older rocks. (2) A thick, undeformed or mildly deformed abyssal sedimentary section is present at the base of the slope and in some areas onlaps the slope. (3) Faults are present along a relatively narrow zone along the continental slope or at the base of the slope, in part recent high-angle reverse or normal faults and possibly strike-slip faults. Most of these features can readily be seen on the seismic data accompanying this report (from west to east, lines 404, 916, 908, 403, 909, and 400; sheet 3).

Both dredge (Plafker and others, 1980) and seismic data show that the lower Tertiary and older strata underlying the continental shelf (strata below horizon D) outcrop on the continental slope throughout the length of the Yakutat segment. Upper Cenozoic strata (D-SF strata) are thin or not present on the slope east of about Yakutat Valley, but are truncated at the sea floor on the continental shelf as a result of uplift on the Fairweather Ground high. West of Yakutat Valley, these strata blanket the upper slope and the lower part of the sequence has prograded seaward over the older strata. The upper part of the sequence crops out with decreasing age toward land, indicating that the sediment supply was insufficient to keep pace with subsidence. The flat-lying to gently landward-dipping strata between Yakutat Valley and the structures of the Pamplona zone have been affected only by regional subsidence.

Oceanic crust at the base of the slope is overlain by sedimentary strata that are as much as 8 km thick; at least part of the sedimentary sequence appears to be of pre-Pliocene age, in agreement with von Huene and others (1979). In the vicinity of Yakobi Valley, the upper part of the abyssal section onlaps the continental margin (lines 400, 987, 909(?), sheet 3). Further to the west, these strata may be faulted against the continental margin (lines 404, 916, 906, 403; sheet 3). Nowhere can seismic reflectors from the abyssal section be traced beneath the continental slope, and reflectors terminate at the base of the slope or onlap the continental slope.

The abyssal strata are undeformed except for uplift in a narrow zone at the base of the slope in areas adjacent to the Fairweather Ground high. Two styles of deformation are present. On seismic section 908, 403, and 909 across the western part of the Fairweather Ground high (sheet 3), the amount of uplift at the base of the slope is greatest in Pliocene and older sediments (below horizon A1) and decreases upwards in Pleistocene strata (A1-SF sequence); only minor deformation is present in the upper part of the Pleistocene section. Thus, this uplift began prior to horizon A1 time, or approximately during Pliocene time. The magnitude of uplift can be estimated for lines 403 and 909 and is around 1 km and 2 km, respectively. Late Cenozoic uplift on the shelf edge and the Fairweather Ground high for these same two lines was discussed earlier and estimated as beginning during Pliocene time (D-SF time), and with magnitudes of around 1 and 2 km, respectively. Thus, the time of onset of deformation and the magnitude of deformation for strata on the shelf and at the base of the slope are about the same, suggesting the deformation at the base of the slope is due to uplift of the Fairweather Ground high. The strata of the outer shelf, slope, and base of slope between Yakobi Valley and Yakutat Valley have been warped upwards during Pliocene and Pleistocene time, with maximum uplift centered on Fairweather Ground. Additional consequences of this uplift may include the erosional truncation of strata at the sea floor along the continental shelf and over Fairweather Ground, and the steep average gradient of the continental slope, ranging from 8° to 16° with a mean of about 11° (Atwood and others, 1981).

The second style of deformation at the base of the slope is seen southwest of Fairweather Ground. Bathymetric data (fig. 3; Atwood and others, 1981) show three subparallel northwest-trending ridges that are 15 to 30 km long. A seismic line across one of these ridges (line 967, sheet 3) shows that it is a young anticline (post-horizon A1 time) that developed in the late Pleistocene. The ridges trend almost perpendicular to the N. 15, W. relative convergence vector for the Pacific and North America plates (Minster and Jordan, 1978; Chase, 1978), and the geometry suggests that

the structures developed in response to this convergence (Atwood and others, 1981).

On all of the seismic sections across the continental slope (sheet 3), faulting is present between the oceanic basalt and the outcropping continental shelf section. As previously discussed, these faults could be in part due to vertical or near vertical uplift associated with the Fairweather Ground structural high, or to recent deformation due to plate convergence. The lack of other major compressional deformation in strata of the continental shelf or at the base of the slope, and the lack of an accreted section along the continental slope suggests that if there is additional faulting, it is primarily strike-slip. Thus, major subduction of the Pacific Plate beneath the continental margin during the late Cenozoic is unlikely. No features have as yet been observed that show strike-slip offset.

An area of bathymetric relief and structural deformation is present in the area of Yushin Ridge, a large northwest-trending ridge or set of ridges present at the base of the slope southwest of Yakutat Valley (fig. 3; also line 404, sheet 3; see also Plafker and others, 1980, figs. 22, 23; and Atwood and others, 1981, sheet 2). Yushin Ridge is about 5 km wide and 27 km long, and rises more than 800 m above the adjacent sea floor. Seismic and dredge data indicate that Yushin ridge is composed of probable lower Tertiary strata, and that relatively undeformed upper Cenozoic sedimentary strata surrounds and buries the ridge (Plafker and others, 1980; Atwood and others, 1981). Thus, this feature developed prior to the late Cenozoic and most likely reflects an early or middle Tertiary erosional or tectonic event.

GEOLOGIC HISTORY AND DISCUSSION

Prior to or about early Eocene time (horizon F time), block or transform faulting along the Dangerous River zone created a structural high to the east of the zone and a basin low to the west. The subsiding basin was floored by basalt and interbedded sedimentary rocks of early Eocene age. Strata deposited into the basin overlapped the basin floor towards the Dangerous River zone. The basin axis trended generally northwest, cutting obliquely across the modern continental shelf (sheets 1 and 2). Deposition of Eocene and Oligocene strata (F-D strata) occurred in increasingly shallower water, and the lower part of the sequence is characterized by transport of shallow water sediments into the deeper parts of the basin (Plafker and others, 1980). The Dangerous River zone may mark the edge of the early Tertiary basin.

West of the Dangerous River zone, local uplift within the Paleogene basin occurred prior to about late Eocene or early Oligocene time (pre-horizon E time), as suggested by minor uplift at location 1 on the reconstructed depth section for seismic section 403 (fig. 9). At early or middle Oligocene time (horizon E time), differential subsidence within the basin formed the gentle structural high seen on seismic section 403. Along horizon E, this deformation created a local unconformity which does not appear to be affected by major erosion. In the rest of the western subbasin, Eocene and Oligocene time (F-D time) is characterized by regional subsidence along the Paleogene basin axis, and no major structures or unconformities are seen in the seismic data. Additional faulting along the Dangerous River zone may also have occurred, resulting in uplift of the area east of the zone. The seismic data leave unclear the timing of early Tertiary faulting along the Dangerous River zone, but most faulting occurred prior to the early stages of deposition of the P-D strata.

Horizon D is a time-stratigraphic break that may include part or all of Miocene time. West of the Dangerous River zone, seismic reflectors are conformable across the horizon, suggesting that no major deformation occurred during the hiatus. The time interval could be a period of regional uplift and erosion. Uplift along the Dangerous River zone during horizon D time caused truncation of P-D strata at horizon D. The terrain east of the zone may have undergone major uplift during this time, resulting in erosion and stripping of most of the lower Tertiary section.

By Pliocene time, and possibly during Miocene time,

deposition of D-SF strata began, initially with these sediments prograding across the continental shelf over the older strata. Gentle uplift began in the Fairweather Ground area during early D-SF time, with more rapid uplift during the last half of D-SF time (approximately late Pliocene and Pleistocene time). Maximum uplift of Fairweather Ground is at least 2 km and could be greater. This uplift resulted in truncation of Cretaceous through Pleistocene strata at the sea floor along the uplift zone, deformation of strata at the base of the slope, and faulting along the continental slope between Yakobl Valley and Yakutat Valley. Throughout the area, the continental shelf is characterized by regional subsidence during D-SF time, with the basin axis near and roughly parallel to the coastline. Reactivation of faults along the Dangerous River zone or faulting along the westward extension of the fault off Lituya Bay has resulted in uplift of D-SF strata offshore of the Dangerous River into a gentle anticline. Additional deformation of the upper Cenozoic strata is present along the shoreline near Lituya Bay, with marked uplift near the shoreline, and along the offshore extension of the Fairweather fault. Finally, recent deformation of strata at the base of the slope southeast of Fairweather Ground has created anticlines that trend almost perpendicular to the regional North America plate-Pacific plate relative convergence vector.

The truncation of Paleogene strata and possible Cretaceous or older rocks of Fairweather Ground at the continental slope indicates that during Cenozoic time, continental and oceanic crust of very different ages have been juxtaposed at the base of the continental slope. The thickness and extent of the Paleogene section beneath the continental shelf west of the Dangerous River zone suggest that at one time these strata were much more extensive than at present and that the seaward part of the Paleogene basin is missing. Since the current morphology of the slope is in part a response to late Cenozoic uplift, the seaward part of the basin might be the thick sediment wedge at the base of the slope. The thickness of the Paleogene shelf section at the slope is similar to the thickness of the pre-Pliocene strata at the base of the slope (see lines 906, 403; sheet 3). However, dredge data indicate the shelf strata were deposited primarily in upper bathyal to shelf environments (Plafker and others, 1980), and it seems unlikely that these strata are continuous with the thick sedimentary sequence at the base of the slope. Also, the section at the base of the slope is younger than the Cretaceous and Paleogene rocks of the shelf, as suggested by the seismic age tie to the DSDP hole 178, near Kodiak Island and discussed above, and by oceanic magnetic anomalies. The oceanic magnetic anomalies on the Pacific plate near this part of the shelf have been identified as numbers 7 through 14, or Oligocene in age, although this age must in part be inferred across a zone at the base of the slope where the anomalies are severely attenuated (Nagler and Wageman, 1973; Taylor and O'Neill, 1974; Schwab and others, 1980). Thus, the sedimentary strata overlying oceanic crust at the base of the slope are most likely Oligocene and younger, thus requiring juxtaposition of terranes of very different ages.

This juxtaposition most likely took place after Oligocene time but prior to Pliocene time, or during the inferred hiatus across horizon D. Evidence for this is: (1) juxtaposition must have taken place after creation of the Oligocene oceanic crust; (2) Eocene through Oligocene strata of the shelf are truncated at the continental slope; (3) strata above horizon D do not appear to be truncated at the slope (in particular, see lines 404, 916, and 906; sheet 3), and have not undergone major deformation during the late Cenozoic, and (4) the upper part of the abyssal section, of Pliocene and younger age, overlaps the continental slope (lines 909, 967, 400; sheet 3). The juxtaposition of the continental shelf and oceanic sections could have occurred either by subduction or by transform faulting, but the absence of an accreted section of early Tertiary age and the absence of major deformation of the Paleogene strata at the shelf edge suggest that subduction has not occurred and that the most likely mechanism is transform faulting. Transform motion is unlikely during the late Cenozoic, due to the onlap relation of Pliocene and Pleistocene oceanic strata against the

continental shelf, and to the absence of deformation of the upper Cenozoic shelf and slope strata (strata above horizon D). The shelf and slope of the Yakutat segment have, therefore, been in proximity to the adjacent oceanic section during at least Pliocene and Pleistocene time, and possibly longer, without major relative motion between the two terranes by either subduction or transform motion during this time period. Thus, the geologic history of the Yakutat segment includes removal of the seaward part of the Paleogene basin and adjacent Cretaceous basement rocks, during about latest Oligocene or early Miocene time, and northwest movement with the Pacific Plate during Pliocene and Pleistocene time, and possibly during part or all of Miocene time as well.

Additional data that support these conclusions include a linear magnetic anomaly, the slope anomaly, that lies along the continental slope between Fairweather Ground and Kayak Island and trends across the shelf between Kayak Island and the Kenai Peninsula. The source body for this anomaly lies within strata of the continental shelf (Schwab and others, 1980) and is likely to be the Eocene basalt (Plafker and others, 1980), with the anomaly resulting as an edge effect from the truncated basalt. This source body has been subducted beneath the shelf west of Kayak Island, and this relation indicates that the continental margin from Cross Sound to Kayak Island is a microplate, the Yakutat block, that has moved with the Pacific plate during at least the last 3-5 m.y. (Schwab and others, 1979, 1980; Bruns and others, 1979).

Plafker and others (1980) note that the most probable provenance for the dredged sandstone in the lower Tertiary sequence of the Yakutat segment was the plutonic and high-grade metamorphic complex of the coastal mountains of British Columbia and Alaska. Such a provenance implies that the dredged strata have been displaced northwest to the Gulf of Alaska by transform faulting along the North America plate margin. If constant movement of the Yakutat block with the Pacific plate is assumed, then a finite rotation of the segment shows that in late Oligocene or early Miocene time, the segment lay roughly between the present position of Vancouver Island and the Queen Charlotte Islands off British Columbia, in general agreement with the location favored by Plafker and others (1980). If these interpretations are correct, the geologic history of the Yakutat segment includes removal of the seaward part of the Paleogene sedimentary sequence and possible rafting of a combined Yakutat-Yakutat segment (the Yakutat block) from a position along what is now British Columbia during late Oligocene or early Miocene time, and movement of the segment northward with the Pacific plate during Miocene to present time. The Yakutat block is now colliding with the North America plate, rather than forming a part of the North America plate and overriding a subducting Pacific Plate.

PETROLEUM POTENTIAL

Three lease sales have been held in the Gulf of Alaska since 1978. Ten exploratory wells drilled in the first area sold, between Icy Bay and Kayak Island, found no commercial discoveries of oil or gas. These wells were drilled on large anticlinal structures with favorable trap potential. However, these wells encountered no adequate source or reservoir rocks, and primarily tested the Neogene sedimentary rocks. The Paleogene section was not adequately tested during this first round of exploratory drilling.

The rocks dredged from the continental slope indicate that potential petroleum source and reservoir rocks are present in the Paleogene sedimentary sequence (Plafker and Claypool, 1979; Plafker and others, 1980). These rocks are immature to marginally mature on the continental slope, but should be more mature beneath the continental shelf. The seismic data show that hydrocarbon traps may be present in the Paleogene strata (P-D strata), but are not likely in the upper Cenozoic section (strata above horizon D). This section briefly reviews the results of Plafker and Claypool (1979) and Plafker and others (1980) on source and reservoir rock characteristics and thermal maturity of rocks on the continental slope, discusses trap potential as seen in the

seismic data, and uses Lopatin's method (Waples, 1980) to estimate the thermal maturity, petroleum potential, and possible petroleum migration history for the strata beneath the continental shelf.

Source Rocks

Analysis of selected dredge samples from the continental slope (Plafker and Claypool, 1979; Plafker and others, 1980) indicates that rocks with favorable hydrocarbon source-rock characteristics are present in the Paleogene sedimentary strata (Units D, E and F of Plafker and others, 1980; F-D strata of this report). Organic matter in selected samples is dominantly herbaceous, with subordinate amounts of humic and amorphous kerogen. Total organic-carbon content of the analyzed samples ranges from 0.42 to 1.87 percent, and averages more than 1.0 percent. This average is above the generally accepted minimum value of 0.5 percent for argillaceous hydrocarbon source rocks (Tissot and Welte, 1978). The kerogen composition is of a type that can generate both liquid hydrocarbons and gas (Plafker and Claypool, 1979; Plafker and others, 1980).

Thermal Maturity

Indicators of thermal maturity show that the dredged samples from units D, E and F are immature to marginally mature but that samples from unit B are mature to overmature (Plafker and Claypool, 1979; Plafker and others, 1980). Oil generation begins at a thermal alteration index (TAI) of about 2 and a vitrinite reflectance value (Ro) of about 0.8 percent; peak oil generation ends at a TAI of around 3 and a vitrinite reflectance value of about 1.35 percent (Dow, 1977, 1979; Tissot and Welte, 1978). Vitrinite reflectance values for dredge samples from units D, E, and F are within the range of 0.3 to 0.8 percent; only a few samples are near or greater than 0.6 percent (Table 5 and fig. 22 of Plafker and others, 1980). The TAI for these samples ranges from 1.2 to 2.3, with most samples around 1.9 to 2.1 (Table 4 of Plafker and others, 1980). Thus, these rocks are immature, and only a few samples approach thermal maturity. Vitrinite reflectance values from unit B range from about 0.4 to 3 percent, and TAI values from 2.3 to 3.8 indicating that these rocks are mature to overmature (Plafker and others, 1980).

Reservoir Rocks

Plafker and others (1980) found that most of the dredged sandstone samples have poor reservoir qualities, due to the presence of abundant unstable mineral and rock fragments, common calcite cement, and siliceous, zeolite, and clay pore fillings. A few samples of sandstone with permeabilities and porosities sufficient to serve as reservoir rocks were recovered from units B, D, and E during dredging; in these rocks, the permeability and porosity appear to be largely secondary.

Hydrocarbon Traps

Except for the anticline near the Dangerous River, seismic data do not show any late Cenozoic anticlinal folds on the Yakutat shelf similar to those found on the adjacent Yakutat shelf to the west (Bruns, 1979; Bruns and Schwab, 1983). At least one large anticline is present in the early Tertiary section (line 403, sheet 3 and fig. 9, location 2), and the extent of this structure and possibly that of some smaller, similar structures, can be inferred from the blocks leased in the 1980 lease sale 55 (sheets 1 and 2). This anticline developed prior to about late Oligocene time and could serve as an effective trap. Additional structures may be present, but are not seen in the widely spaced seismic data.

The Dangerous River zone may provide possibilities for stratigraphic and structural traps. Paleogene strata lap onto the acoustic basement along the Dangerous River zone. These potential traps are updip from the basin axis, and thus could trap hydrocarbons migrating out of the basin lows. The

Dangerous River zone may have formed the margin of Paleogene basin. As noted by Plafker and others (1980), there may therefore be enhanced opportunities for the presence of coarse clastic rocks, up-dip stratigraphic traps, possible Eocene carbonate reefs, and fault seals along the zone. In addition, the overlying upper Cenozoic section could provide a seal over the Paleogene strata in some areas.

The uplift of the Fairweather Ground high is unlikely to provide significant trap possibilities, since Tertiary strata around the high generally crop out at the seafloor, and apparently no effective seal is present. Possible exceptions to this may occur on the west end of the high along the Dangerous River zone.

Lopatin Calculations On Thermal Maturity

The method of Lopatin (Waples, 1980) can be used to evaluate the thermal maturity of the strata beneath the shelf. This method takes into account the cumulative effects of both time and temperature on the maturation of organic matter. Waples (1980) described the method in detail and showed that the time-temperature index of maturity (TTI) calculated by this method generally agrees with other indicators of thermal maturity. The threshold values of Lopatin's time-temperature index (TTI), and corresponding vitrinite reflectance (Ro) and thermal alteration index (TAI) values are (see Waples, 1980, table 3):

TTI	Ro	TAI	
15	0.65	2.85	Onset of oil generation
75	1.0	2.9	Peak oil generation
180	1.3	3.2	End of oil generation

Use of Lopatin's method requires estimates of the geothermal gradient and the burial history of the section of interest. In addition, three major simplifying assumptions are made (Waples, 1980): (1) a constant temperature gradient with depth; (2) a constant temperature gradient through time; and (3) a relatively uniform rate of sedimentation for the age units used in the calculation.

The burial history for sediments beneath the Yakutat shelf can be estimated from the seismic data. The mapped horizons give sediment thicknesses throughout the shelf area. Horizon F, lying at least in part on the top of Eocene basalt, is assigned an age of 50 m.y., or about the age of the basalt as determined by Plafker and others (1980). Horizon E is considered to be on middle Oligocene strata, or about 30 m.y. age. Two end member cases are considered for horizon D. In the first case, horizon D may separate Oligocene from Pliocene strata, so that there is a hiatus from about 24 to 5 m.y. (unconformity case). The effect of this assumption is that strata remain at a constant burial depth and temperature window for the period from 24 to 5 m.y. This case corresponds to age estimates of the shelf strata as discussed earlier in this study, but does not take into account possible deposition of Miocene strata and subsequent stripping during a period of erosion. Maturity could be greater than that calculated. This case gives a conservative estimate of thermal maturity for shelf strata. In the second case deposition is considered as continuous across horizon D during Neogene and Quaternary time (continuous deposition case). The effect of this assumption is that the calculated maturity is optimized.

The geothermal gradient used for the calculation is 25° C/km, as measured in the Continental Offshore Stratigraphic Test (COST) well drilled southwest of Icy Bay in the Yakutat segment of the continental margin (Bolm and others, 1978). Well logs for Colorado Oil and Gas Yakutat wells #1 and #3 indicate bottom-hole temperatures that yield a gradient as high as 30° C/km, and in offshore wells southwest of Icy Bay, the gradient may be 20° C/km or lower. The effect of the higher and lower gradients on the calculated results will be discussed below. The Paleogene strata may initially have had a different geothermal gradient than estimated above. The measured gradients may primarily reflect the rapid deposition of the Neogene strata. This case

is not considered in the calculations.

Lopatin diagrams for the two cases are shown in figure 10. The reconstruction of burial history is for strata adjacent to the main structure leased during lease sale 55 and is at location 2 on line 403 (sheet 3 and fig. 9). The oil-generation window is shown by lines corresponding to a calculated TTI of 15 (Ro = 0.6%) and TTI of 180 (Ro = 1.30%). Several conclusions are common to both cases. (1) Only strata in the Eocene sequence have entered or gone through the oil window; the upper part of the Eocene strata, and Oligocene and younger rocks at this location are likely to be immature. (2) Oil generation within these strata may have started during late Oligocene and Miocene time, with rocks reaching a sufficient depth for peak oil generation (TTI = 75) in Miocene time. (3) Burial had to be greater than about 4 to 4.5 km for oil generation to begin. (4) Only rocks below about 4.5 km (unconformity case) or 4 km (continuous deposition case) are currently in or below the oil window. (5) The oil-generation window is about 1.5 km thick. The main difference between the two cases considered is that the oil-generation window is about 0.5 km higher in the section for the continuous deposition case than for the unconformity case. Calculations for geothermal gradients of 20° and 30° C/km move the oil generation window about 1 km deeper or 0.7 km higher, respectively, in the section than discussed above.

Lopatin calculations for other areas of the shelf have been combined to show the oil-generation window for 25° C/km in two cross sections (fig. 11; locations shown in fig. 3). Section A-A' is along line 912 and extends on strike to the west to line 404; it crosses the thickest part of the early Tertiary basin along the inferred early Tertiary basin axis. Section B-B' corresponds to seismic line 908; and runs across the continental margin. These two cross sections suggest the following conclusions. (1) The top of the oil generation window ranges from about 4 to 5.5 km in depth for the unconformity case and is about 0.5 km higher for the continuous deposition case. The average depth is about 5 km and 4.5 km, respectively. (2) The base of the oil-generation window is about 1.5 km deeper. (3) Potentially mature rocks are primarily of Eocene age. Oligocene strata may be mature only beneath the thickest parts of the Neogene and Quaternary sedimentary basin, primarily in areas beneath the axis and the western quarter of the basin. Neogene rocks are potentially mature only below the axis of the basin. (4) Potentially mature strata are most likely present only west of the Dangerous River zone. (5) Strata at the continental slope are likely to be immature. This conclusion matches observed thermal maturity indicators in the dredged rock samples.

In general, hydrocarbon generation throughout the basin may have begun in Eocene strata in about Miocene time. Migration of these hydrocarbons updip would drive the generated hydrocarbons toward the edges of the basin onshore and at the continental slope, where the potential for traps appears to be poor, and east up the basin axis towards the Dangerous River zone and the structures leased during lease sale 55. These structures formed at least in part during Oligocene and early Miocene time, or prior to or during potential hydrocarbon generation and migration, and may thus have good potential for trapping migrating hydrocarbons.

The Lopatin calculations strongly suggest that if suitable source rocks are present in the Eocene strata beneath the shelf, at least some of these strata are mature enough to have generated oil and gas. Since the dredge samples suggest source rocks are present in the exposed section, the major unknowns are whether migration has occurred and whether suitable reservoir rocks are present to trap migrating hydrocarbons.

The Lopatin calculations can be combined with volume estimates of sedimentary rocks of the Yakutat segment to estimate the volume of sediment that may be in or below the oil window. The total volume of sediment can be measured from the structural contour maps. The limits of the area considered are between the Yakobi Valley and the Pamplona zone, the shoreline (extended across the mouths of the major bays), and the lower Tertiary outcrop or subcrop at the continental shelf. In areas where contours are not defined by the seismic data, the contours are estimated by extending them along strike. With these limits and assumptions, the total

volume is about 138,500 km³, of which 75,000 km³, or 54 percent are Eocene and Oligocene strata (F-D strata) and 63,500 km³ or 46 percent are Neogene and Quaternary strata (D-SF strata). Approximately 90 percent of the total sediment volume is east of the Dangerous River zone (east of the lower Tertiary subcrop line, sheets 1 and 2). For the unconformity case and a geothermal gradient of 25° C/km, the average depth to the oil window is about 5 km, and the volume of sediment in or below the threshold for oil generation is therefore about 23 percent (fig. 12). For the continuous deposition case, averaging about 4.5 km to the oil window, the volume is about 28 percent. For geothermal gradients of 20° and 30° C/km, the corresponding volume ranges are about 13 to 18 percent and 34 to 40 percent, respectively. Since almost all the strata in the oil window are of Paleogene age, these volume ranges suggest that from 25 percent (at 20° C/km) to 70 percent (at 30° C/km) of the total Paleogene section could be within or below the oil window, depending on the assumptions of burial history and geothermal gradient.

Even though drilling results in the Gulf of Alaska have so far been disappointing, there are still reasons for optimism for the discovery of economic hydrocarbons beneath the Yakutat segment. The major positive consideration is the presence of organic-rich rocks dredged from the continental slope that could be good source rocks beneath the continental shelf. These rocks extend beneath the adjacent Yakataga segment, and are the likely source for the numerous seeps onshore. These strata are untested by drilling that has so far taken place. Thus, the major exploration strategy for the Gulf of Alaska is to search for the Oligocene and Eocene strata in favorable trap locations at drillable depths beneath the Yakutat and Yakataga segments.

REFERENCES

- Atwood, T. J., Bruns, T. R., Carlson, P. R., Molnia, B. F., and Plafker, George, 1981, Bathymetric maps of the northern Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-859, 3 sheets, scale 1:250,000.
- Bayer, K. C., Mattick, R. E., Plafker, George, and Bruns, T. R., 1978, Refraction studies between Icy Bay and Kayak Island, eastern Gulf of Alaska: U.S. Geological Survey Journal of Research, v. 6, no. 5, p. 625-636.
- Beikman, H. M., 1974 Preliminary geologic map of the southeast quadrant of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-612, scale 1:1,000,000.
- Beikman, H. M., 1975, Preliminary geologic map of southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-673, scale 1:1,000,000.
- Beikman, H. M., 1980, Geologic Map of Alaska: U.S. Geological Survey, 1 sheet, scale 1:2,500,000.
- Bolm, J. G., Chmielek, F. B., Stewart, G. H., Turner, R. F., Waetjen, H. H., and Wills, J. C., 1976, Geological and operational summary, Alaska stratigraphic test OCS 75-80, No. 3, northern Gulf of Alaska: U.S. Geological Survey Open-File Report 76-635, 40 p.
- Bruns, T. R., 1979, Late Cenozoic structure of the continental margin, northern Gulf of Alaska, in Sisson, Alexander, ed., The relationship of plate tectonics to Alaskan geology and resources: Alaska Geological Society Symposium, Anchorage Alaska, 1977, Proceedings, p. 11-130.
- Bruns, T. R., Atwood, T. J., and Childs, J. R., 1981a, Free-air gravity anomaly map, Cross Sound to Icy Bay, northern Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1306, scale 1:500,000.
- Bruns, T. R., and Bayer, K. C., 1977, Multichannel seismic reflection data acquired on the M/V Cecil H. Green in the Gulf of Alaska, June-August, 1975: U.S. Geological Survey Open-File Report 77-352.
- Bruns, T. R., Childs, J. R., and Atwood, T. J., 1981b, Free-air gravity anomaly map, Dixon Entrance to Cross Sound, northeastern Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1307, scale, 1:500,000.
- Bruns, T. R., and Plafker, George, 1982, Geology, structure, and petroleum potential of the southeastern Alaska and northern Gulf of Alaska continental margins, in Bruns, T. R., ed., Hydrocarbon resource report for proposed OCS Lease Sale 88: southeastern Alaska, northern Gulf of Alaska, Cook Inlet, and Shelikof Strait, Alaska: U.S. Geological Survey Open-File Report 82-828, 133 p.
- Bruns, T. R., and Schwab, W. C., 1983, Structure maps and seismic stratigraphy of the Yakataga segment of the continental margin, northern Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1424, 3 sheets, scale 1:250,000.
- Bruns, T. R., Schwab, W. C., von Huene, Roland, Atwood, T. J., Blakely, R. J., and Case, J. E., 1979, Geologic significance of the slope anomaly, northern Gulf of Alaska: American Geophysical Union Transactions, v. 60, no. 46, p. 951.
- Burkhard, Nardía, Bruns, T. R., McHendrie, A. G., and Ruppel, B. D., 1980a, Map of free-air gravity anomalies, eastern Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1172, scale 1:500,000.
- Burkhard, Nardía, von Huene, R. E., McHendrie, A. G., Ruppel, B. D., and Bruns, T. R., 1980b, Map of free-air gravity anomalies, western Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1173, scale 1:500,000.
- Carlson, P. R., Plafker, George, Bruns, T. R., and Levy, W. P., 1979, Seaward extension of the Fairweather fault, in Johnson, K. M., and Williams, J. R. eds., The United States Geological Survey in Alaska: Accomplishments during 1978: U.S. Geological Survey Circular 804-B, p. B135-B139.
- Chase, C. G., 1978, Plate kinematics: The Americas, East Africa, and the rest of the world: Earth and Planetary Science Letters, v. 37, p. 355-368.
- Dix, C. H., 1955, Seismic velocities from surface measurements: Geophysics, v. 20, no. 1, p. 68-86.
- Dow, W. G., 1977, Kerogen studies and geological interpretations: Journal of Geochemical Exploration, v. 2, no. 2, p. 79-99.
- , 1979, Petroleum source beds on continental slopes and rises, in Geological and geophysical investigations of continental margins: American Association of Petroleum Geologists Memoir 29, p. 423-442.
- Lahr, J. C. and Plafker, George, 1980, Holocene Pacific-North American plate interaction in southern Alaska: Implications for the Yakataga seismic gap: Geology, v. 8, no. 10, p. 483-486.
- Minster, J. B. and Jordan, T. H., 1978, Present-day plate motions: Journal of Geophysical Research, v. 83, no. B11, p. 5331-5354.
- Naugler, P. P. and Wageman, J. M., 1973, Gulf of Alaska: magnetic anomalies, fracture zones, and plate interactions: Geological Society of America Bulletin, v. 84, no. 5, p. 1575-1584.
- Perez, O. J., and Jacob, K. H., 1980, Tectonic model and seismic potential of the eastern Gulf of Alaska and Yakataga seismic gap: Journal of Geophysical Research, v. 85, no. 812, p. 7132-7150.
- Plafker, George, 1967, Geologic map of the Gulf of Alaska Tertiary province, Alaska: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-484, scale 1:500,000.
- , 1971, Possible future petroleum resources of Pacific-margin Tertiary basin, Alaska, in Future petroleum provinces of North America: American Association of Petroleum Geologists Memoir 15, p. 120-136.
- Plafker, George, and Addicott, W. O., 1976, Glaciomarine deposits of Miocene through Holocene age in the Yakataga Formation along the Gulf of Alaska margin, in Miller, T. P., ed., Recent and ancient sedimentary environments in Alaska: Alaska Geological Society Symposium, Anchorage, 1975, Proceedings, p. Q1-Q23.
- Plafker, George, Bruns, T. R., Carlson, P. R., Molnia, B. F., Scott, E. W., Kahler, Roger, and Wilson, Charles,

- 1978a, Petroleum potential, geologic hazards, and technology for exploration in the outer continental shelf of the Gulf of Alaska Tertiary province: U.S. Geological Survey Open-File Report 78-490, 33 p.
- Plafker, George, Bruns, T. R., and Page, R. A., 1975, Interim report on the petroleum resource potential and geologic hazards in the outer continental shelf of the Gulf of Alaska Tertiary province: U.S. Geological Survey Open-File Report 75-592, 74 p.
- Plafker, George, Carlson, P. R., Coonrad, W. L., Hunt, S. J., and Quintero, Paula, 1979, Geologic implications of 1978 outcrop sample data from the continental slope, in Johnson, K. M., and Williams, J. R., eds., *The United States Geological Survey in Alaska: Accomplishments during 1978*: U.S. Geological Survey Circular 804-B, p. B143-B146.
- Plafker, George, and Claypool, George, 1979, Petroleum source potential of rocks dredged from the continental slope in the eastern Gulf of Alaska: U.S. Geological Survey Open-File Report 79-295, 24 p.
- Plafker, George, Hudson, Travis, Bruns, T. R., and Rubin, Meyer, 1978b, Late Quaternary offset along the Fairweather fault and crustal plate interactions in southern Alaska: *Canadian Journal of Earth Sciences*, v. 15, no. 5, p. 805-818.
- Plafker, George, Winkler, G. R., Coonrad, W. L., and Claypool, George, 1980, Preliminary geology of the continental slope adjacent to OCS Lease sale 55, Eastern Gulf of Alaska; Petroleum Resource Implications: U.S. Geological Survey Open-File Report 80-1089, 72 p.
- Plafker, George, Winkler, G. R., Hunt, S. J., Bartsch-Winkler, Susan, Coonrad, W. L., and Quintero, Paula, 1978c, Outcrop samples from the continental slope in the eastern Gulf of Alaska: U.S. Geological Survey Circular 772-B, p. B87-B99.
- Rau, W. W., Plafker, George, and Winkler, G. R., 1977, Preliminary foraminiferal biostratigraphy and correlation of selected stratigraphic sections and wells in the Gulf of Alaska Tertiary Province: U.S. Geological Survey Open-File Report 77-747, 54 p.
- Rogers, J. P., 1977, Implications of plate tectonics for offshore Gulf of Alaska petroleum exploration: Offshore Technology Conference, Houston, Texas, 1977 Proceedings, v. 1, p. 11-18.
- Schwab, W. C., and Bruns, T. R., 1979, Preliminary residual magnetic map of the northern Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1054, scale 1:500,000.
- Schwab, W. C., Bruns, T. R., and von Huene, Roland, 1979, Magnetic lineaments in the northern Gulf of Alaska and their geologic significance (abs.): *Geological Society of America Abstracts with Programs*, v. 11, no. 7, p. 512.
- Schwab, W. C., Bruns, T. R., and von Huene, Roland, 1980, Maps showing structural interpretation of magnetic lineaments in the northern Gulf of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1245, scale 1:500,000.
- Stoneley, Robert, 1967, The structural development of the Gulf of Alaska sedimentary province in southern Alaska: *Geological Society of London Quarterly Journal*, v. 123, p. 25-37.
- Taylor, P. T., and O'Neill, N. J., 1974, Results of an aeromagnetic survey in the Gulf of Alaska: *Journal of Geophysical Research*, v. 79, no. 5, p. 719-723.
- Tissot, B. P., and Welte, D. H., 1978, *Petroleum formation and occurrence*: New York, Springer-Verlag, 538 p.
- von Huene, R. E., Molnia, B. F., Bruns, T. R., and Carlson, P. R., 1975, Seismic profiles of the offshore Gulf of Alaska Tertiary province, R/V *Thompson*, Sept.-Oct., 1974: U.S. Geological Survey Open-File Report 75-664.
- von Huene, R. E., Shor, G. G. Jr., and Wageman, J. M., 1979, Continental margins of the eastern Gulf of Alaska and boundaries of tectonic plates, in Watkins, J. S., and Montadert, L., eds., *Geological and geophysical investigations of continental margins*: American Association of Petroleum Geologists Memoir 29, p. 273-290.
- Waples, D. R., 1980, Time and temperature in petroleum formation: application of Lopatin's method to petroleum exploration: *American Association of Petroleum Geologists Bulletin*, v. 64, no. 6, p. 916-926.
- Winkler, G. R., 1976, Deep-sea fan deposition of the lower Tertiary Orca Group, eastern Prince William Sound, Alaska, in Miller, T. P., ed., *Recent and ancient sedimentary environments in Alaska*: Alaska Geological Society Symposium, Anchorage, 1975, Proceedings, p. R1-R20.



Figure 1. Index map of Alaska showing study area.




Figure 2. Simplified geologic and tectonic setting of the northern Gulf of Alaska showing structural trends and general distribution of Paleozoic, Mesozoic, and Cenozoic rocks. Geology after Beikman (1980); Pacific plate-North America plate relative-convergence vector (large arrow) from Minster and Jordan (1978). CS, Cross Sound; DF, Denali fault; DRF, Duke River fault; FF, Fairweather fault; FG, Fairweather Ground; IB, Icy Bay; KI, Kayak Island; MI, Montague Island; PZ, Pamplona zone; QCI, Queen Charlotte Islands; QC-FF, Queen Charlotte-Fairweather fault system; TF, Totschunda fault; YB, Yakutat Bay; Wrangell Volcs., Wrangell volcanoes and volcanic field.




Figure 4. Results of refraction sonobuoys and correlation of velocity layers. Velocities given in kilometers per second (km/s). Locations of sonobuoy lines shown in figure 3.




Figure 5. Time-depth curve, interval-velocity curve, error limits, and comparison with well velocity curve. Time-depth curve is used to convert two-way traveltime on seismic sections to depth; "Z" is subsurface depth and "t" is two-way travel time.




Figure 6. Correlation of seismic horizons D, E, and F of this report to geologic units A-G defined by Plafker and others (1980) from dredge data from the continental slope. Locations of correlations are shown in degrees west longitude. Vertical ruled lines indicate a hiatus; diagonal lines indicate no data. Correlation to onshore geologic formations can be determined by comparing this chart to figure 2 of Plafker (1971).




Figure 7. True-scale depth section (top) and restored section derived by flattening on horizon D for a segment of seismic section 909 north of Fairweather Ground. Location of the line segment is shown in figure 3 and on sheet 3. Reconstruction at horizon D time shows approximate configuration of lower Tertiary strata prior to major uplift of Fairweather Ground. Amount of uplift is about 2 km, and may be about 2.5 km if uplift is mainly after horizon D time.




Figure 8. True-scale depth section (top) and restored section derived by flattening on horizon U for a segment of seismic section 911 north of Fairweather Ground. Location of the line segment is shown in figure 3 and on sheet 3. Reconstruction at horizon U time shows approximate configuration of strata prior to major uplift of Fairweather Ground. Amount of uplift is about 1 km. Age of horizon U is unknown, but is probably Pliocene or younger. See text for further discussion.




Figure 9. Depth section (top) for part of seismic section 403 and sequential reconstruction of strata configurations at horizon D, U, and E times. Reconstructions show history of development of structural highs at locations 1 and 3, and a low at location 2. Location of the line segment is shown in figure 3 and on sheet 3. Large arrows on each reconstruction indicate subsidence or uplift with respect to previous reconstruction. See text for further discussion.





Figure 10. Lopatin diagrams showing the burial history and calculated time-temperature index of maturity (TTI) of 15 and 160, corresponding to beginning and end of oil generation (Waples, 1980), for strata at location 2, line 403 (fig. 9 and sheet 3). Calculation assumes a geothermal gradient of 25°C/km. Strata beneath horizon F are assumed to have little or no resource potential. A. "Unconformity case" assumes horizon D separates Oligocene from Pliocene strata, and that no major subsidence or uplift occurred during the hiatus. B. "Continuous-deposition case" assumes no hiatus at the end of Oligocene time, and continuous deposition across horizon D. The actual burial history probably lies somewhere between the two cases shown here, and could include both uplift and subsidence at horizon D time.




Figure 12. Plot of cumulative percent volume versus depth (percent of total volume at or below a particular depth). Oil and gas limits shown are a shelf-wide average based on Lopatin calculations of thermal maturity with a geothermal gradient of 26 C/km. The corresponding volume of sediment that may be mature then ranges from around 23 to 28 percent of the total volume. See text for discussion of calculations using other geothermal gradients.