








By
Bruce R. Johnson and Geoffrey S. Elliott
1984

CORRELATION OF MAP UNITS

	QUATERNARY
1n	TERTIARY (?)
	CRETACEOUS (?)
	CRETACEOUS
1a	CRETACEOUS AND JURASSIC
	TRIASSIC (?)
	MESOZOIC AND PALEOZOIC

LIST OF 90P UNITS	
Qal	Alluvial deposits, undivided
Tf	Telsic plutonic rocks
Mf	Mafic plutonic rocks
Ed	Diorite sill, extensively altered
Ko	Sitka Gneiss
Kb	Kay Group
KJf	Telsic plutonic rocks
KJm	Mafic plutonic rocks
W	Whitestripe Marble
Bg	Green Bp Gneissstone
NpPa	Unconformable, metamorphic, and plutonic rocks

SYMBOLS

 Contact, approximately located, dotted where concealed
 Boundary of study area
 Geochemical sample site
 Lead (Aa) = 30-50 ppm
 Lead (Aa) > 50 ppm
 Lead (Spec) = 50-70 ppm
 Lead (Spec) > 70 ppm

STUDIES RELATED TO WILDERNESS

The Wilderness Act (Public Law 88-577, September 3, 1964) authorized the U.S. Geological Survey and the U.S. Bureau of Mines to survey certain public lands in Alaska for potential mineral resource potential. Results must be available to the President and Congress by January 1, 1968. This report presents the results of geological survey of the Western Chikofka-Nabok Islands located in the Bering Sea off the coast of Alaska. About 65 percent of the study area has been covered by a detailed geological map under the Alaska National Interest Land Conservation Act (P.L. 86-487).

This report covers the areas of the U.S. Geological Survey investigations of the Western Chikofka-Nabok Islands located in the Bering Sea off the coast of Alaska. Samples were collected. Samples were analyzed for 31 elements by a wet digestion method followed by atomic absorption spectrophotometry (AAS). The AAS method (Cotton, 1960) was used for elements by atomic absorption spectrophotometry (Ward and others, 1966). The following elements are listed: As, Ba, Be, Bi, Br, Ca, Cd, Co, Cr, Cu, Fe, Ga, Ge, Hg, K, Li, Mn, Mo, Ni, Pb, Pt, Rb, Se, Si, Sr, Ta, Te, Ti, Tl, U, V, W, Zn, Zr. Coordinates, and a station location map are available for each sample. A map of the islands is included. Elliott, 1964. A map and discussion of the mineral resource potential of the study area is also available.

Background levels for each element vary for different lithologies in the study area. Because of this, the use of a single value for background levels such as sampling technique, analytical variance, and chemical weathering, it is impossible to select a specific analytical level above which values indicate significant enrichment. The use of a greater likelihood of bedrock mineralization, but confidence levels are low for single element high values and results which are not supported by neighboring values. This study suggests that the use of high analytical values for the element lead by two analytical techniques, as well as the locations of all 2,735 samples. Multiple symbols for a single analytical technique for a sample site represent multiple samples at that site.

REFERENCES CITED

Grimes, D. J., and Narinero, A. F., 1968, Direct correlation of aluminum content of rock with emission spectrographic field methods for the quantitative analysis of aluminum in rock materials; U.S. Geological Survey Circular 591, 10 p.

Johnson, R. W., 1962, Magmatic cane coning trace element patterns in the basal rocks of the West Chicago-Tofino Islands Volcanic Stratovolcano, Alaska; U.S. Geological Survey Information Service Report No. USGS-60-82-005, 10 p.

Johnson, R. W., and Elliott, G. R., 1966, Map showing bedrock geochronological station locations, Western Alaska; U.S. Geological Survey Bulletin 1247-B, 1 sheet, southeastern Alaska; U.S. Geological Survey Bulletin 1247-C, 1 sheet, Alaska.

Johnson, R. W., and Karl, S. W., 1962, Macromagma-tic differentiation of the basal rocks of the West Chicago-Tofino Islands Volcanic Stratovolcano, Alaska; U.S. Geological Survey Bulletin 1247-A, 1 sheet, Alaska.

Johnson, R. W., and Karl, S. W., 1963, Petrographic Field Studies Map US-1476-a, scale 1:125,000, Alaska; U.S. Geological Survey Bulletin 1247-B, 1 sheet, Alaska.


Johnson, R. W., and Karl, S. W., 1963, Mineral resource potential map of the Western Alaska Tofino Islands Volcanic Stratovolcano, Alaska; U.S. Geological Survey Bulletin 1247-C, 1 sheet, Alaska.

Johnson, R. W., and Karl, S. W., 1963, Petrographic Field Studies Map US-1476-b, scale 1:125,000, Alaska; U.S. Geological Survey Bulletin 1247-D, 1 sheet, Alaska.

Ward, R. L., Johnson, R. W., Burns, T. F., and Van Dine, J. L., 1969, Apatite, zircon, and titanite of magmatic origin in geochronal exploration; U.S. Geological Survey Bulletin 1247-E, 1 sheet, Alaska.



Index Map Showing Location of Study Area



SCALE 1:125 000

0 2 4 6

2 0 2 4 6

This map is preliminary and has not been reviewed for conformity with U. S. Geological Survey editorial standards, but the stratigraphic nomenclature has been approved geologically.