

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

RECEIVED

FEB 9 1978

U.S. DEPARTMENT OF THE INTERIOR
BUREAU OF GEOLOGICAL SURVEY

MAPS AND TABLES DESCRIBING AREAS OF METALLIFEROUS MINERAL
RESOURCE POTENTIAL OF CENTRAL ALASKA

(Text and tables to accompany Open-file Report 78-1-D)

by

G. Donald Eberlein and W. David Menzie

This report is preliminary and has
not been edited or reviewed for
conformity with Geological Survey
standards and nomenclature

Menlo Park, California
1978

Contents

	<u>Page</u>
Introduction -----	1
Acknowledgments -----	2
Philosophy and methodology -----	2
Status of geologic and mineral resource information on central Alaska -----	6
Explanation of maps and table pertaining to designated areas of mineral resource potential -----	9
Delineation and designation of areas -----	9
Summary mineral resource information -----	10
Summary and conclusions -----	38
References cited -----	43

Tables

Table 1. Reports prepared as a foundation for mineral resource assessment of central Alaska -----	3
2. Metalliferous mineral resource data for central Alaska -----	12
3. Grade and tonnage models -----	37

INTRODUCTION

This report constitutes an appraisal of the known and potential metallic mineral resources of central Alaska, utilizing the best and most up-to-date geologic and resource data available. Its main purpose is to provide information for forthcoming decisions on classification of Alaska's lands arising from the Alaska Native Claims Settlement Act of 1971. The report is one of five resource appraisals of Alaska (U.S.G.S. Open-file reports 78-1-B to F) that constitute end products of the U.S. Geological Survey's Regional Alaska Mineral Resource Assessment Program (RAMRAP). Four of the appraisals (B-E) describe mineral resources, and one (F) describes oil and gas resources. This report consists of two 1:1,000,000-scale maps (designated sheets 1 and 2 of open-file report 78-1-D) showing areas of known and speculative mineral resource potential, a summary of pertinent geologic and mineral resource data and of conclusions pertaining to resource potential (table 2), grade and tonnage models (table 3), and this explanatory text.

Central Alaska as considered in this report embraces an area of approximately 631,000 km² (156,800,000 acres; 245,000 mi²; approximately 42 percent of Alaska's total land area) that extends from the Canadian boundary (meridian 141°) on the east to the Bering Sea on the west and from the approximate southern boundary of the Brooks Range on the north to the northern flank of the Alaska Range on the south.

Specifically excluded from this analysis are the fossil fuels, geothermal energy resources, and deposits of nonmetallic minerals, excepting asbestos. However, maps and tables that provide an inventory of the known coal and thermal spring occurrences in central Alaska are included in the folio of background data outlined in table 1, and areas of interest for oil and gas in central Alaska are described in U.S. Geological Survey Open-file report 78-1-F of this series.

ACKNOWLEDGMENTS

Most of the Alaskan information upon which this analysis is based has been derived from published and unpublished works of previous investigators--mainly geologists and mining engineers representing both government and private sectors. These sources have been cited individually in the publications listed in table 1. We are especially indebted to W. P. Brosge, R. M. Chapman, H. L. Foster, J. M. Hoare and W. W. Patton, of the U.S. Geological Survey, who are presently engaged in Alaskan mineral resource investigations and who have greatly enhanced our assessment by contributing relevant up-to-date unpublished information on the distribution and geologic controls of certain previously known and newly discovered mineral deposits in central Alaska. The authors are also grateful to D. A. Singer and J. H. DeYoung, Jr., for their contributions to the development of grade/tonnage models for certain deposit types (table 3). This assessment has also been improved by several industry geologists who generously supplied pertinent data on the results of recent company exploration programs. Particularly cooperative in this regard have been C. L. Sainsbury and Don Stevens. The timely completion of this report would not have been possible without the technical assistance of Judith S. Gassaway, Beth Abramson, Anelia Lott, Mark Sander, Voncille Johnson and Barbara Thompson.

PHILOSOPHY AND METHODOLOGY

Information concerning a region's mineral endowment is an essential part of intelligent land use planning. It should be at a scale that permits the evaluation of individual tracts of land and should be in a form that is useful for analysis from a number of perspectives, including national mineral needs, economics, currently available and/or projected technologies, and competing land uses. The information should include a resource inventory of known

Table 1.--Reports prepared as a foundation for mineral resource assessment
of central Alaska

U.S. Geological Survey		
Open-file report numbers	Author(s)/Compiler(s)	Title ¹
77-168-A	G. Donald Eberlein, Judith S. Gassaway, and Helen M. Beikman	Preliminary geologic map of central Alaska
77-168-B	Edward H. Cobb	Placer deposits map of central Alaska
77-168-C	David F. Barnes	Preliminary Bouguer gravity map of central Alaska
77-168-D	G. Donald Eberlein, Robert M. Chapman, Helen L. Foster, and Judith S. Gassaway	Map and table describing known metalliferous and selected nonmetalliferous mineral deposits in central Alaska
77-168-E	John Decker and Susan Karl	Preliminary aeromagnetic map of central Alaska
77-168-F	John Decker and Susan Karl	Preliminary aeromagnetic profiles of central Alaska
77-168-G (in press)	Judith S. Gassaway	Map (with table) showing occurrences of coal in central Alaska
77-168-H (in press)	Judith S. Gassaway	Map showing distribution of known thermal springs and selected igneous rocks in central Alaska

¹All maps at scale: 1:1,000,000

deposits and numerical estimates of possible undiscovered mineral deposits grouped according to similar physical, chemical, mineralogical, and grade/tonnage characteristics.

The present analysis should be considered as a first step toward a comprehensive assessment of central Alaska's mineral resources and has used the following approach:

- (1) Relevant data on the more than one-thousand known lode and placer mineral occurrences in central Alaska were systematically reviewed and analyzed to determine the types of deposits present and, insofar as possible, their geologic controls and regional settings.
- (2) Using the information in (1), areas of mineral resource potential have been delineated (see map sheets 1 and 2) on the basis of existing deposit type(s) and/or on geologic factors and relations permissive for the occurrence of specific deposit types.
- (3) For a number of deposit types it has been possible to construct models of their expected grades and tonnages of mineral commodities or of contained metals (grade times tonnage). These models have been constructed following the procedure of Singer, Cox, and Drew (1975) and utilize data derived from similar deposit types worldwide (see table 3). These models are referenced (table 2, column 10) in cases where they have been judged to be representative of known or speculated deposit types in given areas, and they provide an indication of relative resource abundance.
- (4) In the seven cases where the data have permitted, subjective probability estimates of the number of deposits, by type, likely to exist within a delineated area have been made on the basis of

the number of known deposits, favorable geologic considerations, the nature and extent of exploration and any favorable geochemical and/or geophysical indicators. These estimates are presented in terms of high (90 percent), intermediate (50 percent), and low (10 percent) confidence levels that the indicated number of deposits, or more, may occur. For most areas, however, the information level is insufficient to permit estimation of the probable number of deposits present. In such cases qualitative statements as to resource potential and, where appropriate, applicable grade-tonnage models have been entered into table 2.

The four mineral resource assessments prepared under RAMRAP share a common philosophy, methodology and format, but details of the individual assessments differ as a function of the amount and mix of information available. For some parts of central Alaska the information level is so low that it has been possible only to generally delimit very large regions of potential for the occurrence of one or more deposit types, largely on the basis of geologic analogy and favorable setting (for example, map areas 4, 12 and 14). As geologic and mineral resource information increases and our understanding of various deposit types improves, it should become possible to better define and delimit such regions into smaller areas for which more specific and precise future assessments of contained mineral resources can be made. Thus, this assessment will become outdated as geologic and mineral resource information increases.

STATUS OF GEOLOGIC AND MINERAL RESOURCE INFORMATION
ON CENTRAL ALASKA

The adequacy of any assessment of mineral resource potential to satisfy land use planning, regional mineral exploration and other requirements is determined in large measure by the level and completeness of the available geologic information. The assessment must be made at a scale that is consonant with the information base.

Information about the geology and mineral deposits of central Alaska has been developed over a period of more than 75 years, but prior to 1959 it was mainly of a reconnaissance nature and with few exceptions was very strongly controlled by the occurrence and distribution of placer gold. Accordingly, the information level differed drastically from region to region and in many cases has proven to be inadequate as a basis for obtaining much more than the crudest perspective of overall mineral resource potential, especially in regions removed from gold-producing districts. Furthermore, even the most comprehensive of the earlier surveys did not develop certain kinds of geologic data necessary to evaluate the likelihood that some of the more recently recognized deposit types (e.g., volcanogenic sulfide deposits, stratabound and stratiform base metal deposits, island arc porphyry copper deposits, composite porphyritic biotite granite intrusives having a potential for tin-tungsten deposits) may occur in a given area. Notable exceptions were commodity-oriented investigations for tin, tungsten and quicksilver in selected areas by the U.S. Geological Survey and U.S. Bureau of Mines during World War II under the Strategic Minerals Program and reconnaissance examinations for radioactive deposits by the Geological Survey on behalf of the Division of Raw Materials, U.S. Atomic Energy Commission, during the middle and late 1940's and early 1950's.

Although great disparities still exist among different parts of central Alaska, considerable progress has been made since 1959 toward improving the level of geologic information. Mainly as a result of the Geological Survey's 1:250,000-scale regional geologic mapping program slightly more than 50 percent of the region is now covered by published, modern, ground-based geologic mapping at that scale (Bartsch-Winkler, 1977a). However, only about 3.3 percent of central Alaska is covered by geologic maps at a scale of 1:63,360 or larger (Bartsch-Winkler, 1977b), the minimum considered acceptable to satisfy most regional mineral exploration requirements in the western conterminous United States.

Considerable progress also has been made in recent years toward improving the quality and extent of geophysical and geochemical data for central Alaska. Contoured and/or profile aeromagnetic coverage, suitable for regional extrapolations at scales between 1:500,000 and 1:63,360, is now available for approximately 71 percent of the region, and reconnaissance soil, stream-sediment and bedrock geochemical data, ranging from spotty to fairly extensive, now exist for almost 40 percent of the region.

In July 1974, the Geological Survey initiated a multidisciplinary Alaskan Mineral Resource Assessment Program (AMRAP) directed toward evaluating the endowment of terranes having high economic mineral potential on a 1:250,000-scale quadrangle format. Unfortunately for this analysis the results of AMRAP investigations to date are available for all or parts of only five of the forty-eight 1:250,000 Alaska Topographic Series quadrangles that cover the region. However, from these early results it is evident that such studies are generating the kinds of data that are essential to modern regional resource analyses. As a result, future assessments can be expected to have a considerably improved geologic and mineral resource information base that

will enable them to be more definitive, precise, and conclusive.

Perhaps the most serious hindrance to the development of an understanding of the bedrock geology of central Alaska, especially as it bears on mineral resource evaluation, has been, and will continue to be, imposed by the region's natural environment. With few notable local exceptions, central Alaska physiographically is characterized by unglaciated uplands and lowlands whose evolution has been controlled mainly by periglacial erosional processes operating under a subarctic permafrost climate. The uplands have been sculptured largely by creep and solifluction, and their lower slopes are commonly mantled by windborne silt. Thus, large areas are covered by transported material and by vegetation that obscure bedrock. Unglaciated lowlands have been formed mainly by depositional processes that blanket bedrock with materials brought in from highlands and tend to erase irregularities in the lowland surface. Underlying permafrost prevents downward percolation of groundwater, and summer melting of ice at the top of permafrost keeps extensive vegetation cover relatively saturated. Thus, large areas of unglaciated lowlands are marshy flats almost wholly lacking in bedrock exposures. It is estimated that 40 percent of central Alaska ($252,000 \text{ km}^2$; $98,000 \text{ mi}^2$) is blanketed by a Quaternary surficial cover of alluvial, glaciofluvial, moraine, muck, swamp, lacustrine, flood-plain, and windblown deposits, and undeformed subaerial volcanic rocks (1.4 percent). Accordingly, most bedrock geologic investigations in these extensive areas of surficial cover are and doubtless will continue to be much less conclusive than desired for definitive mineral resource evaluation purposes, even on a regional scale.

EXPLANATION OF MAPS AND TABLE PERTAINING TO DESIGNATED AREAS OF
MINERAL RESOURCE POTENTIAL

Delineation and designation of areas

This report includes two 1:1,000,000-scale maps (designated sheets 1 and 2), covering the eastern and western parts of central Alaska, and on which 77 areas of mineral resource potential have been delineated. Each area is identified by a number which keys it to the same numbered entry in table 2. The numerical order is progressive in general from west to east and carries no implications as to the relative importance or favorability of one numbered area with respect to any other. Areas considered favorable or permissive for the occurrence of one or more of the same deposit type(s) and known or believed to have similar regional geologic characteristics from a mineral resource standpoint have been assigned the same number. Most of the major and minor commodities known or suspected to occur within each area are indicated by standard chemical symbols. Minor and suspected commodities are listed parenthetically. The order of their listing is random and is not indicative of relative abundance or potential importance.

It should be emphasized that important differences exist in the basis for delineating the various areas of resource potential. Certain areas have been identified on the basis of known deposits and on extensions of their geologic features. Other areas have been outlined because results of preliminary or reconnaissance geologic and/or mineral exploration investigations indicate that they could contain certain deposit types that have been productive in similar geologic settings elsewhere. Some areas in the first group are known to contain important resources; the resource potential of those in the second group can only be speculative; the remainder fall between these two extremes.

The maps show that some parts of central Alaska have not been included among those designated as having significant mineral resource potential. For the most part these areas have been excluded because (1) they contain no known deposits or only a few scattered small deposits in a generally unfavorable geologic setting, and/or (2) they are mantled by a thick cover of unconsolidated surficial deposits, or by young volcanic rocks. Although such areas may contain concealed deposits at depth, it is felt there is little likelihood of their being discovered and exploited using presently known or foreseeable techniques.

Summary of mineral resource information

The principal data used for defining areas of mineral resource potential are summarized in telegraphic style in table 2. Included are qualitative and, where the data permit, probabilistic estimates of potential. The table contains ten columns, which are discussed briefly in numerical order from left to right below. For more detailed information on individual deposits and mining districts, the reader is invited to consult U.S. Geological Survey Open-file Report 77-168-D.

- Column 1. Area outlined on maps.--The entry number shown in this column correlates with corresponding numbered area(s) of mineral resource potential on sheets 1 and 2 of the map.
- Column 2. Types of known deposits.--Major types of mineral deposits known to occur within the area are identified in column 2 using terminology commonly accepted among most economic geologists. In most cases there is evidence of production. Where more than one deposit type is listed, they are prefixed by lowercase letters to facilitate reference to related comments in the other columns. Principal and subordinate (in parentheses) resources are indicated by standard chemical symbols.
- Column 3. Suspected or speculative types of deposits.--Listed here in the same format as in column 2 are deposit types whose occurrence within the designated area is considered permissive because of favorable geologic considerations and/or exploration history. Also included are minor occurrences.

- Column 4. Geologic controls of mineral resources.--Geologic characteristics of known, suspected, or speculative deposit types listed in columns 2 and 3 are briefly stated and, where possible, principal geologic factors believed responsible for their localization are indicated.
- Column 5. Production and resource information.--Available data and estimates of production, estimates of identified resources and generalized statements about potential for the deposit types listed in columns 2 and 3 are summarized here.
- Column 6. Status of geologic information.--This column contains a summary of the status and scale of geologic mapping, geophysical and geochemical surveys, and mineral resource investigations. Recent industry exploration and mining activities are also noted.
- Column 7. Additional comments.--Statements in this column in general are intended to further clarify or amplify the statements concerning geologic controls in column 4. Additional information bearing on the potential for the occurrence of speculative deposit types is also presented here.
- Column 8. Summary of mineral resource potential.--This column contains generalized information relevant to the overall mineral resource potential of the area, as well as a summary of the principal geologic factors that render the area permissive for the occurrence of certain deposit types.
- Column 9. Estimated number of deposits.--Where warranted by the available information, subjective probabilistic estimates of the number of deposits, by type, likely to exist within the designated area are presented in this column. These estimates are assigned high (90 percent), intermediate (50 percent), and low (10 percent) levels of confidence that the indicated number of deposits, or more, may be present. The absence of an entry in this column indicates that data necessary to construct such estimates are lacking.
- Column 10. Grade and tonnages for deposit types.--Where occurrences in a given area are judged representative of a designated deposit type, reference is made to the appropriate grade-tonnage or contained-metal model to provide an indication of expected size (table 3). Not all models presented in table 3 have been referenced in this column because not all of the deposit types modeled are known to be represented in central Alaska.

TABLE 2. METALLIFEROUS MINERAL RESOURCE DATA FOR CENTRAL ALASKA

(See map sheets 1 and 2)

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
1.	(a)Pt, Au--placer	(b)(Cr, Hg)--placer	(a)Mainly modern stream and bench placers of Salmon River and its western tributaries that drain Red Mountain dunite intrusive (Jurassic?) and that is considered to be the ultimate source of the platinum metals. Gold reconcentrated into placers from glacial outwash (b)Small amounts of chromite and locally native mercury and cinnabar in beach and stream placers	(a)Total platinum production from Goodnews Bay district estimated to be well over 20,000 kg (650,000 troy ounces), plus about 485 kg (15,600 ounces gold). Most productive ground involved two pay-streaks mined over a distance of about 10 km (6 miles) in Salmon River valley. Drilling demonstrates Salmon River platinum-bearing ground extends at least as far south as Happy Creek but not as far as Chagvan Bay about 5.6 km (3½ miles) farther south. Platinum reportedly present in sediments of Goodnews Bay, but results of industry sampling program not available. No workable lodes containing platinum have yet been found. Although Red Mountain dunite intrusive is not considered a favorable site for the discovery of any large low-grade workable ores of platinum metals, some small high-grade concentrations may exist. Total identified resources in beaches, bays and lodes has been estimated at 210,000 kg (6.8 million troy ounces). Platinum metals potential of offshore area essentially unknown but could be sizable (b)Beach placers locally contain low grade chromite resources and restricted occurrences of gold and platinum	Reconnaissance USGS 1:250,000 geologic mapping of late 1940's and 1950's, updated 1974-1976 under AMRAP. Regional and local detailed geochemical sampling and aeromagnetic surveys by government and industry. Comprehensive USGS study of Goodnews Bay district platinum deposits	Preservation of placer ground largely controlled by extent of glacial scouring. Platinum placer mining Goodnews Bay District ceased 1975. Attempts to sample sea floor west of Red Mountain for platinum have been generally inconclusive. Bed-rock and stream sediment geochemical mercury anomaly south of Goodnews Bay	Significant past production of placer platinum > 20,000 kg; (> 650,000 oz) and byproduct gold (485 kg;15,600 oz) from bench and stream placers in Salmon River area. Significant additional resources of platinum probably exist in extensions of the Salmon River placers, and in beaches, bays and lodes		

<u>AREA OUT- LINED ON MAPS</u>	<u>MAJOR TYPES OF KNOWN DEPOSITS</u>	<u>SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)</u>	<u>GEOLOGIC CONTROL(S) OF MINERAL RESOURCES</u>	<u>PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)</u>	<u>STATUS OF GEOLOGIC IN- FORMATION</u>	<u>ADDITIONAL COMMENTS</u>	<u>SUMMARY OF MINERAL RESOURCE POTENTIAL</u>	<u>ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)</u>	<u>GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)</u>
2.	Au, (Pt)--placer	- - -	Bench and stream placers. Ultimate source of gold probably Tertiary granitic intrusive rocks and reconcentration from glacial deposits	Total production not known but estimated less than 3,100 kg (100,000 ounces) mainly from Slate and Mattanuse Creeks, which have been extensively explored and drilled. A small amount of platinum recovered from Bear Creek placers	Area covered by recent (1961) 1:250,000 scale regional geologic mapping, updated under AMRAP studies. State contoured aeromagnetic coverage (1:250,000 scale)		Estimated production of gold from stream and bench placers has been < 3,100 kg (< 100,000 oz). One stream produced a little byproduct platinum. The area has been extensively explored		
3.	Au (Pt)--placer	- - -	Ultimate source of gold in Pre-Pleistocene (?) bench and present stream gravels may have been the mineralized major deep-seated fault zone that separates Precambrian Kanektok terrane from the Ordovician to early Cretaceous Ahklun terrane to southeast. By-product platinum probably derived from Tertiary mafic intrusives	Total production not known, but may have exceeded 3,100 kg (100,000 oz) small amount of byproduct platinum also locally recovered	Recent (1961) 1:250,000 scale USGS geologic mapping updated under AMRAP studies. State contoured aeromagnetic coverage (1:250,000 scale)		Estimated production of gold from stream and bench placers has been < 3,100 kg (< 100,000 oz). Some streams have produced a little byproduct platinum		

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND YIELDAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
4.	(a)Hg(Sb)--vein and open space filling (low temperature hydrothermal) (b)Au--placer	(c)Au, (Ag, Hg, Sb, Zn, Cu, Pb, W, Sn)--mainly quartz veins associated with lower Tertiary granitic intrusives; in part porphyry gold	(a)Principal deposits and groups of deposits localized along major, deepseated, through- going fault zones. Local controls related to structurally and chemically suitable sites that offered sufficient permeability for the passage of the ore-forming solutions and adequate porosity for the deposition of cinnabar and stibnite. Most (but not all) productive deposits are in or adjacent to faulted and altered mafic dikes and sills (Tertiary) that in- trude sedimentary and volcanic rocks of the Kuskokwim (Cretaceous) and Gemuk (Carboniferous to Cretaceous) Groups. (b)Ultimate source of gold in many of the most productive pre- sent stream and bench placers believed to be irregular quartz veins in granitic plu- tons (early Tertiary) and in associated hornfels zones. Es- pecially rich placers locally formed where present streams tran- sect older bench gra- vels. (Compare Ruby district placers, area 14.) Some gold also re-concentrated from glacial and/or glaciofluvial deposits. Semi-residual placers locally developed over deeply weathered quartz monzonite in- trusives (early Tertiary) as near Flat	(a)Total production of mercury over past 75 years about 1,300 tons (38,000 flasks; 76 lbs. per flask), mainly from the Red Devil Mine about 1,100 tons (32,000 flasks). A small amount of by- product antimony also produced. A few known regional mercury geochemical anomalies. (b)Placer deposits wide distributed throughout region and have accounted for almost 13% about 70,000 kg (2,252,000 fine ounces) of Alaska's total placer gold production, plus byproduct silver, through 1960. Principal producers include Iditarod- Flat area about 41,400 kg (8,130,000 ounces) Inoko area about 17,000 kg (550,000 ounces) and Candle Creek about 3,100 kg (100,000 ounces) (c)Of 13 occur- rences, only two known mines had small production (gold, silver, lead, and zinc) and are now inac- tive. "Porphyry gold" deposits associated with quartz monzonite stocks probably have a significant low grade resource potential	Southwestern part of re- gion covered by reconnais- sance 1:250,000 scale geo- logic mapping, updated by AMRAP studies, including state contoured aeromag. (1:250,000 scale). Most of remainder covered by pre-1955 reconnaissance mapping and ranks among geologically least known areas in Alaska. Much of region characterized by poor exposures. Most known lode mercury-antimony deposits have been examined by USGS and many have been sampled by USBM. Also USBM sampling programs in Flat area for gold and mer- cury. Extensive regional geochemical sampling pro- gram by industry in early 1970's	(a>About two dozen mercury-antimony lode deposits are known in addition to the ten that have been productive and more are likely to be found. A few regional geochemical anomalies. Cinnabar also present in many gold placers, but none has produced more than a few flasks of mercury. (b)Placer concen- trates also contain ubiquitous cinnabar, stibnite, scheelite, and base metal sul- fide minerals; lo- cally cassiterite, platinum, monazite, orpiment, realgar, allanite, and blis- suth also present. Chromite found in some of the placers probably derived from small ultra- mafic intrusives	(a)Ten lode depos- its have produced mercury in this region and approx- imately two dozen additional deposits are known to occur. The region is con- sidered favorable for the occurrence of more deposits (b)Bench and stream placers have pro- duced approximately 70,000 kg (2,252,000 oz) of gold (c)Production from two small gold-bear- ing quartz vein de- posits was about 84 kg (2,700 oz) of gold and 81 kg (2,600 oz) of silver, about 4 tons of lead and a little zinc ($< .5$ ton). Addi- tional deposits are possible. "Porphyry gold" deposits, associated with quartz monzonite stocks, probably con- tain a significant resource of low-grade gold	90% 50% 10% chance that there are 30 70 110 deposits or more	(a)mercury model)

<u>AREA OUT- LINED ON MAPS</u>	<u>MAJOR TYPES OF KNOWN DEPOSITS</u>	<u>SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)</u>	<u>GEOLOGIC CONTROL(S) OF MINERAL RESOURCES</u>	<u>PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)</u>	<u>STATUS OF GEOLOGIC IN- FORMATION</u>	<u>ADDITIONAL COMMENTS</u>	<u>SUMMARY OF MINERAL RESOURCE POTENTIAL</u>	<u>ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)</u>	<u>GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)</u>
4. (cont.)			(c) Irregular quartz veins up to two inches thick cut lower Tertiary quartz monzonite stocks and contain gold, scheelite and base metal sulfide minerals, and a little cassiterite. Others contain cinnabar and stibnite. Emplacement of the plutons may have been controlled by Iditarod-Nixon fault and related structures. Widely scattered mineralized quartz veins also associated with small lower Tertiary albite rhyolite porphyry intrusions						

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
5.	(a)Fe(Ti)--- magmatic	- - - (b)Au--placer (c)Mo--vein (hydrothermal?) (d)Hg (Sb)--possible vein and open space filling lode deposits	(a)Single known south- eastward dipping tabular titaniferous magnetite bearing pyroxenite intru- sive at least 1,200 m (4,000 feet) thick under- lying area of 10 km ² (4 square miles) or more (Kamuk Mountain). (b)Stream and/or bench placers. (c)Auriferous quartz- carbonate veinlets in contact zone between serpentinite and fine grained mafic intrusive. (d)Similar to notation #1 under area 4(a)	(a)Estimated 2.4 billion tons of hypothetical re- sources containing 15-17% total iron and 10.5-12% mag- netic iron. No production. (b)No known production. (c)Single known occurrence. Channel samples contain up to 11 ppm gold. No known production	(a)Company airborne and ground geophysical surveys and diamond drilling pro- gram. Small area north of Dillingham covered by USGS contoured aeromag (1:125,000 scale). Re- gional geology poorly known. Western part covered by pre-1940 recon- naissance 1:250,000-scale geologic mapping by USGS Extensive glacial cover (including eolian).	(a)Deposit covered by 90-400 feet of glacial deposits. Most rocks are sil- icified argillite quartzite and lime- stone (Permian- Triassic?). (c)One occurrence of molybdenite reportedly found near head of Silver Horn on Lake Beverly (d)No known deposits but available geo- logic information suggest that undis- covered deposits may be present in western part of area Note: adjoins area 8, southern Alaska (USGS Open-file report 78-1-E)	(a)A single titani- ferous magnetite- bearing pyroxenite intrusive contains an estimated 2.4x10 ⁹ tons of 15-17% total iron (10.5-12% mag- netic iron) (d)Although there are no known mer- cury mines or pros- pects within this region, its western portion is similar in geology to area 4, a region of known mercury deposits and prospects	(d)mercury model	
6.	(a)Au(Pt, Hg, Sb)-- placer	(b){Au, Cu, Sb}--vein (hydrothermal) (c){Cu}--magmatic (?) (d){Cu, Zn, Ag}-- possible submarine volcanogenic sulfide deposits relate to mafic lavas (?)	(a)Modern stream, bench, and fossil placers. Source of most of gold was gold- quartz veins in and ad- jacent to Upper Cretaceous quartz monzonitic plutons and smaller Upper Creta- ceous-lower Tertiary albite rhyolite intrusives. Byproduct platinum proba- bly derived from small ultramafic (Jurassic ?) intrusives. (b)Quartz veins associ- ated with some of the Cretaceous granitic intru- sives contain gold, stib- nite and copper minerals. (c)Tuluksak River dredge yielded specimens of bornite and asbestos from ultramafic bedrock (d)Although their occur- rence has not yet been demonstrated, the belt of Jurassic (and some Permian) mafic volcanic rocks that underlies the NW part of the area may have some potential for submarine mafic volcano- genic sulfide deposits	(a)Most of the major gold pro- duction from bench placers, as at Bear Creek, where still older under- lying gravels may have potential for more gold. Minor platinum recovered from some of the placers but the amount is unknown. Many of the streams draining northward into Kuskokwim R. contain traces of gold and may have some placer poten- tial, especially beneath the blanket of glacial outwash. (b)No known pro- duction. (c) - - - (d) - - -	Reconnaissance 1:250,000 scale-geologic mapping (1959) by USGS. Little prospecting. Essentially no presently available geochemical coverage. Northern half covered by ERDA aeromag and gamma ray and profiles (1:500,000 scale)	(a)Cinnabar a common consti- tuent of many of the placers, notably those of Bear Creek	(a)Bench, stream and fossil placers have been mined. Fossil placers may have potential for addi- tional gold. (b)Quartz veins, which carry gold, copper and antimony minerals, are scat- tered throughout the region (d)The belt of Permo- Jurassic mafic vol- canic rocks, that occurs in the north- west part of the region, is permissive for the occurrence of mafic volcanogenic deposits. No occur- rences are known in this generally poorly explored region	(d)mafic volcanogenic model	

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
7.	(a)Au(Pt)--placer	(b)Au(Cu, Pb, Mo, Ag) vein (hydrothermal)	(a)Mainly modern stream placers. Source of gold probably quartz veins related to Upper Creta- ceous-lower Tertiary granitic intrusives into mafic volcanic and assoc- iated clastic rocks (Permian-Jurassic) (b)Quartz-calcite veins up to 30 cm (12 ins.) thick contain native gold, galena, molybdenite, chal- copyrite and traces of scheelite and wolframite. Arnold prospect report- edly contained high silver values	(a)Estimated total gold production about 3,000 kg (96,000 fine ounces) plus byproduct sil- ver and a little platinum, mainly from five deposits (b)Lode deposits known since about 1914, but none has been productive. Potential exists for discovery of additional vein deposits but theirs total pre- cious and base metal resources probably are small	Recent (1959, 1976) 1:250,000-scale recon- naissance geologic map- ping by USGS. Little known company activity. Regional aeromagnetic, gamma ray profiles (ERDA), and reconnaissance geo- chemical (USGS) coverage.	(a)Cinnabar pre- sent in many of the placer con- centrates. Minor scheelite at Elephant Creek. Area includes pillow basalts, diabase and gabbro with minor chert argillite and volcaniclastic rocks that may correlate with Rampart belt of volcanic ophiolites (see area 11)	(a)Stream gold pla- cers have been mined (3,000 kg; 96,000 oz). Some of the streams carry byproducts, in- cluding silver, plat- inum and tungsten (b)A number of gold and sulfide bearing quartz veins are known in the region. To date none has been product- ive. Potential exists for additional discov- eries, but resource potential considered small (c)Region contains rock types which are geologically permis- sive for the occur- rence of mafic vol- canogenic deposits. No occurrences are presently known	(c)mafic volcanogenic model	
8.	(a)Hg---veinlets and open space fillings (b)Au, Ag (Cu, Fe, Bi, W, Th, REs)-- contact metamorphic (c)Au (Bi, Sn, W, Th, U)-- placer	(a)(Sb)---possible stibnite associated with mercury minerals in veinlets, dissemina- tions and open space fillings (?)	(a)Cinnabar occurs mainly as breccia filling in dolomite and dolomitized limestone (Middle? Ordo- vician) along southeast- ward dipping faults that are probably related to Farewell fault zone (see also item #4 above) (b)Tactite zones (including skarn) in lower Paleozoic platform carbonate rocks adjacent to quartz mon- zonite stocks (Upper Cretaceous-lower Tertiary). Gold also disseminated in the monzonite (porphyry gold?). (c)Stream placer deposits, especially those that cross contacts between quartz monzonite intru- sives and carbonate rocks (see above)	(a)White Mountain deposit has pro- duced an estimated 14+ tons (400+ flasks) of mercury since 1964. Cinnabar distributed over an area 6-8 km (4-5 miles) long and up to 1.6 km (1 mi) wide (b)Lode mines in Nixon Fork area have produced an estimated 1,600 kg (50,000 ozs) of gold and a little silver, mainly between WWI and WWII. Copper con- tent ranges from 2-12% but no at- tempt made to recover it. (c)Production data on gold placers lacking but in Nixon Fork area the amount proba- bly was less than that from the lode deposits	Southern part of region is one of the geologic- ally least known areas of Alaska. Northern part included in AMRAP studies now in progress. Geochemi- cal studies in Nixon Fork mine area by state and USGS. Central areas cov- ered by 1977 USGS aeromag- netic profiles (1:250,000- scale)	(a)Stibnite not a recognized con- stituent of the White Mountain deposit, but could be present in as yet undis- covered lodes. (b)Nixon Fork lodes contained auriferous pyrite, chalcopryite, copper carbonate minerals, native bismuth, and minor amounts of bornite, scheelite and traces of radio- active minerals (parisite, allanite) (c)Placer concen- trates in Nixon Fork area also included scheelite, cassiterite, allan- ite and thorionite	(a)One deposit, White Mountain, has pro- duced 14+ tons (400+ flasks) of mercury. Other deposits may occur (b)3 skarn deposits are known. One, Nixon Fork, has produced about 1,600 kg (50,000 oz) of gold and a little silver; copper was present but not produced. Region is favorable for the occurrence of additional deposits (c)Stream placers have been mined		

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
9.	- - -	(a)(Cu, Zn, Ag)-- possible submarine volcanogenic sulfide deposits related to mafic lavas (?) (b)(Cu, Pb, Zn, Mo, Au) --- vein and porphyry deposits associated with granitic and rhy- olitic igneous rocks (?) (c)(Pt, Cr, Fe, Ti)--- magmatic deposits (?) (d)(Au)--placer (?)	Although the only known mineral resources derive from a single occurrence of chalcopyrite in con- tact zone of a diorite plu- ton and from placer gold on bars of the Anvik River and its tributaries, the following fea- tures contribute to an overall geologic setting considered suitable for the occurrence of deposits types indicated as specula- tive in the adjacent column: (1)widely distri- buted marine mafic volcanic rocks (Lower Cretaceous), in part propylitically altered, and intruded both by granitic plutons (Upper Cretaceous) and high-level rhyolitic bodies with their extrusive counter- parts (Upper Cretaceous- lower Tertiary; (2)major northeastward trending faults; (3)several mafic and at least 1 known ultramafic intrusive body	Area has no known recorded production; definitive resource data lacking	Reconnaissance geologic mapping by USGS (1:2,500- scale). Southern quarter covered by ERDA aeromag- netic and gamma ray profiles (1:500,000-scale). Little known prospecting	Area also included in the belt of upper Cretaceous- lower Tertiary rhyolitic intru- sive and extrusive rocks described under area 11 below	Although this region contains only 1 known mineral occurrence, it has been little pros- pected and the geolo- gic setting is permis- sive for the occurrence of the following deposit types: mafic volcanogen- ic deposits (associated with Cretaceous basalt and andesite), porphyry copper deposits (asso- ciated with Cretaceous- Tertiary rhyolitic vol- canics and high level intrusives), podiform chromite deposits (as- sociated with ultra- mafic rocks) and gold stream placers (asso- ciated with Quater- nary alluvial deposits)	(a)mafic vol- canogenic model (b)porphyry copper model (c)podiform chromite model	
10.	- - -	(a)(Cr, Asbestos)--- possible podiform chromite, chromite layers and cross-fiber asbestos deposits as- sociated with ultra- mafic rocks. (b)(Ni)---possible nickeliferous resid- ual deposits (?)	(a)Known occurrences of chromite and small seams of cross-fiber asbestos in typical alpine serpent- inized dunite and peridotite tectonite (and overlying layered gabbro) that con- stitute the ultramafic parts of allochthonous ophiolite complexes. (b)Potential residual deposits enriched in nickel (laterites) ultra- mafic rocks, particularly at the edge of and beneath overlapping non-marine coal-bearing mid-Creta- ceous-lower Tertiary sedimentary rocks that border the ultramafic ophiolite belt northwest of the Ray Mountains	(a)Chromium values range as high as 9% in selected USGS grab samples of dun- ite streaked with grains of chrome spinel from the Kanut-Kilotna Creeks area. Com- posite chip samples contained 2400-3000 ppm chromium, 1900- 2400 ppm nickel and below detectable (atomic absorption) amounts of platinum group metals. Since no visible sulfide minerals have been detected the anomal- ously high nickel content is consid- ered to be present in silicate miner- als. Bedrock samples from ultra- mafic rocks in the Yukl River region contain 1000-1500 ppm nickel and generally 700-1500 ppm chromium	Recent 1:250,000 geologic mapping and more detailed studies of the ophiolites by USGS. Locally good bed- rock and stream sediment sampling coverage. Con- toured state aeromagnetic maps available for most of the areas except in Nulato quadrangle. Little known industry activity	These ultramafic and layered gabbro ophiolitic complex- es have been in- terpreted as re- presenting the root zones of allochthonous sheets, the lead- ing edges of which are the basalt- diabase-chert complexes noted under area 11 below. Area 10 on map also locally contains elements described under area 12 below	(a)Known occurrences of chromite and asbestos in alpine ultramafic bodies. Podiform chro- mite deposits are possible (b)Although there are no known deposits, the geologic setting is permissive for the occurrence of laterite nickel deposits	90% 50% 10% chance that there are 6 20 66 deposits or more	(a)podiform chromite model

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
11.	- - -	(Cu, Zn, Ag)---possible submarine volcanogenic sulfide deposits	Pillow basalt, diabase and gabbro with minor chert, argillite, vol- caniclastic rocks and limestone of the Rampart belt of dismembered ophi- olite complexes (Miss- issippian to Jurassic) offer a potentially suitable geologic set- ting for the occurrence of massive and/or dissem- inated stratabound vol- canogenic sulfide deposits associated with pillow lavas similar to those at Cyprus, Oman, and else- where	No known occur- rences and re- source data essentially lack- ing. Limited geochemical data indicate the ophiolitic assem- blage northwest of Livengood carries the highest values of copper, (gen- erally 15-700 ppm; 1 sample showed 15000 ppm) and up to 0.1 ppm silver in bedrock samples	Most of the indicated areas covered by post 1966 USGS 1:250,000-scale geologic mapping except for Kaiyuh Mountains-Yukl River area. Regional ERDA aeromag and gamma ray pro- file coverage (1:500,000- scale). Reconnaissance USGS geochemical data exists for most of the areas. Little known exploration activity by industry	Small mafic and ultramafic intru- sives associated with these ophi- olites may have some potential for chromium and asbestos. (Bed- rock samples locally carry > 5000 ppm Cr).	Geologic setting is considered per- missive for the occurrence of mafic volcanogenic sulfide deposits although no such deposits, or even occurrences, are presently known. Region has not been thoroughly prospected for this deposit type	mafic volcan- ogenic model	
12.	- - -	(Pb, Zn, Ag, Cu, Au)--- possible disseminated sulfide deposits and veins associated with silicic intrusive and extrusive igneous rocks (?)	Numbered areas define a belt at least 725 km (450 miles) and up to 128 km (80 miles) wide that contains swarms of porphyritic rhyolitic to dacitic dikes, sills and other small plutons and non-marine extrusives (Upper Cretaceous-lower Tertiary) that are for the most part localized along northeast trend- ing faults. Some are as young as Miocene. Locally developed pyritiferous zones are extensively oxidized. Altered zones of silica and sericite. Area may contain undiscovered porphyry copper deposits	No known produc- tive deposits and relevant resource data almost totally lacking. However, one such altered zone about 90 m (100 yards long) in the upper Kanuti River area contains disseminated galena, sphalerite and py- rite. Grab samples contain 700-20,000 ppm lead, trace to 3,000 ppm zinc, 3-30 ppm silver and 70-500 ppm copper. Similar altered rhyolite bodies on the Indian River about 120 km (75 miles) to the southwest contain small, but anomal- ous, amounts of lead copper, sil- ver, gold and molybdenum	Northeastern half covered by recent USGS 1:250,000 scale regional geologic mapping. Other data in- clude ERDA aeromag and gamma ray 1:500,000 pro- files, and some USGS bed- rock and stream sediment geochemical data. South- western part generally poorly known. No known systematic prospecting for the indicated specu- lative targets	The geologic setting is also considered po- tentially suitable for the occurrence of uranium	This region has not been through- ly prospected for base metals. The rock types present (small felsic in- trusive and extru- sive rocks) and their structural setting, together with extensive altered areas and associated geo- chemical anomalies, contribute to a geologic setting which is consid- ered permissive for the occur- rence of porphyry copper deposits and uranium de- posits associated with granitic and rhyolitic rocks	porphyry copper model	

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
13.	(a)Au, W(Sb)--- placer	- - - (b){W, Sn}---placer (c){Sb, Au}---vein	(a)Gold in bench and modern stream placers of Ungalik River area pro- bably derived from small mineralized intrusives (Upper-Cretaceous) into marine Cretaceous (Albian?) sedimentary rocks. (b)Beach placers (c)Scores of claims staked pre-1909 on Christ- mas Mountain Upper Creta- ceous quartz monzonite- granodiorite intrusive body	(a)Production data lacking, but prob- ably well over 370 kg (12,000 oz) of gold. Mining (including dredging) has spanned over 60 years at least through 1975. A few pounds of scheelite recovered 1910. Placer gold may also be con- centrated at base of submerged alluv- ium along former valley of Ungalik River and as an offshore residual deposit formed when sea level rose to its present position. (b)Beach deposits along northwest shore Norton Bay contain traces of cassiterite (tin) and scheelite (tungsten). (c)Stibnite float and colors of gold from streams drain- ing Christmas Mountain, but no record of any development work or production	Recent reconnaissance unpublished 1:250,000-scale geologic mapping and coastal studies by USGS; USBR reconnaissance sampling of beach and river deposits, Horton Bay	(a)Stibnite present in heavy concen- trates. Offshore and beach placers may be present at mouth of Ungalik River (c)Stibnite re- ported in quartz veins on ridge between Shaktolik and Ungalik Rivers	(a)Stream and bench gold placers have been mined as re- cently as 1975. Total production well over 370 kg (12,000 oz). A few kilograms of byproduct scheelite (tungsten) have been recovered. Buried Pleistocene stream channels beneath Norton Bay have some potential for placer gold (c)Stibnite-gold veins associated with Upper Creta- ceous granitic intrusive bodies occur in the region		

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE J)
14.	(a)Au(Sn, Pt, Bi)-- placer (b)Ag, Pb(Zn, Cu, Au, Mo)--polymetal veins	(c)(U, Th)--possible disseminations in igne- ous rocks(?) (d)(Sn, W, Be)--possi- ble vein, disseminations and/or tactite deposits (e)(Cu, Zn, Pb, Ag)-- possible stratabound (volcanogenic?) massive and/or disseminated sulfide deposits	Areas outlined general- ly define the Ruby ge- anticline which is under- lain by pelitic and cal- careous schist, quartzite, and carbonate rocks (Pale- ozoic and Precambrian?) and widely intruded by Cretaceous quartz monzo- nite, biotite granite and by granodiorite, and smaller granitic plutons of Late Cretaceous-early Tertiary age. (a)About 75% of the more than 70 placer gold mines, prospects and occurrences are concentrated in the Poorman-Long-Ruby area where pay streaks are for most part, deeply buried, discontinuous, and bear little relation to present stream channels. Ultimate source of metals considered to be small quartz veins in highly decomposed metamor- phic bedrocks and related to Upper Cretaceous-early Tertiary granitic intru- sives. (Compare Ophir dis- trict, area 4.) (b)Mainly argentiferous galena-bearing quartz veins in quartz-mica schist. (c)Quartz monzonite and granite of the Melozitna pluton contain above- average amounts of uranium and thorium. Radiometric anomalies over parts of the Sithylenkat Lake and Melozitna plutons suggest possible disseminated ura- nium and thorium in intru- sives and wall rocks. (d)Many of the Cretaceous granitic plutons of the Ruby geanticline and their associated contact zones contain pegmatites and quartz-tourmaline veins, as well as geochemically anom- alous amounts of tin, boron, beryllium, lithium and, lo- cally, lead and rare earths. These and other geologic considerations make the in- dicated areas permissive for the occurrence of	(a)Poorman-Ruby- Long areas has pro- duced well over 12,000 kg (400,000 fine ounces) of gold (about 2% of Alaska's total placer gold out- put), plus byprod- uct silver and a little platinum. A few tens of tons of cassiterite also recovered, but most was not shipped to smelter. Some of the ground was ex- ceedingly rich. Geologic nature of the placers suggest some undiscovered pay streaks may still exist. Pro- duction from the Morelock and Tozi- moran Creeks area pro- bably did not exceed 150 kg (5,000 oz) of gold, plus byprod- uct silver. Cas- siterite a common constituent of those placers and some was recovered, but it is uncertain that any was ever marketed. (b)Of seven known occurrences only one has produced. Perseverance mine and adjacent Valley claims yielded 204 tons (225 short tons) of ore that averaged 73% lead and 3.6 kg per ton (104 oz silver per short ton). Undis- covered vein depos- its probably exist, but their resource potential for the indicated metals is likely to be low. (c)(d)(e)No produc- tion. Definitive resource data lack- ing	Recent 1:250,000-scale geologic mapping by USGS. Government airborne mag- netic and radiometric re- gional coverage fairly complete except southwest part of area. Some USGS bedrock and stream sedi- ment geochemistry. Recent claim staking in Melozitna Hot Springs area (radio- metric anomaly). Other- wise little known system- atic prospecting except for lode and placer gold	(a)Native bismuth, platinum, scheelite and allanite pres- ent in some of the placer concen- trates. Locally includes elements of area 12 NOTE: There are differences in dis- tribution and boundaries of area 14 between Central and Northern Alaska (USGS Open-file Rept. 78-18) re- gions owing to dif- ferences in geo- logic interpreta- tions of the avail- able data in the respective regions	(a)Fossil gold placer deposits have been mined in the region. Production has been well over 12,400 kg (400,000 oz). By- products recovered include silver, platinum and tin. There is a potential for additional de- posits in the region. (b)One deposit and six occurrences of argentiferous galena- bearing quartz veins are known in the re- gion. The one depos- it was low tonnage (204 tons) but very high grade (73% lead and 3.6 kg silver per ton). Some potential exists for additional deposits. (c)The rock types (granitic intrusives) and known local radio- metric and geochemical anomalies signify a geologic setting per- missive for the occur- rence of disseminated uranium and thorium minerals. (d)The rock types present (granitic in- trusive rocks), to- gether with anomalous- ly high values of tin, beryllium and lead in geochemical samples, suggest a geologic setting permissive for the occurrence of tin and tungsten in vein, greisen and/or tactite deposits. (e)Although this re- gion contains no known occurrences of acid- intermediate volcano- genic sulfide deposits, it is not known to have been thoroughly pros- pected for base metal deposits, especially of that type. The rocks present are coextensive with and	(d)tungsten tactite model (e)felsic and intermediate volcanogenic sulfide model	

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
14. (cont.)			tin-tungsten vein or greisen deposits and metalliferous tactite (skarn) deposits, especially where the plutons have intruded calcareous schist. (e) The pelitic schists and associated metamorphic rocks or the Ruby geanticline are lithogenetically similar to, and are considered to be co-extensive with, the low- to medium-grade metamorphic rocks of the southern Brooks Range that contain the recently discovered Arctic and Picnic Creeks stratabound (volcanogenic?) massive and/or disseminated sulfide deposits. This region of central Alaska is therefore considered to be geologically permissive for the occurrence of such as yet undiscovered deposits				lithologically similar to pelitic schists in the southern Brooks Range that contain the Arctic and Picnic Creek deposits. Thus, the region is considered permissive for the occurrence of felsic and intermediate volcanogenic sulfide deposits		
15.	(a) Au--placer	(b) Cu, Pb, An, Ag, (Mo, Au, W)--veins and disseminations	(a) Mainly modern stream gravels that drain contact zones between Lower Cretaceous andesitic rocks and Upper Cretaceous quartz monzonite and granodiorite plutons, as well as mineralized zones in the volcanics. Some reconcentration from glaciofluvial deposits. (b) Tetrahedrite-galena-sphalerite veins in altered andesitic volcanic rocks (Lower Cretaceous) near the Indian Mountain granodiorite-quartz monzonite pluton (Upper Cretaceous); disseminated sulfide minerals in fault controlled, altered, silicified, fine-grained felsic intrusive rocks (Upper Cretaceous or lower Tertiary) and andesitic host rocks (see area 12 above)	(a) Total known production about 260 kg (8,500 oz) gold plus byproduct silver, an amount much too small in view of the extensive mining on Indian River and Bear and Utopia Creeks. Dredging on Bear Creek as recently as 1975. (b) Anomalous amounts of copper (up to 1,500 ppm), lead (up to 3,000 ppm), zinc (up to 3,000 ppm), silver (up to 700 ppm), and gold (up to 6 ppm) in stream sediment and/or selected grab samples. No known development work. Potential resource for indicated metals in known and undiscovered lodes is probably small	Reconnaissance 1:250,000 geologic mapping (1966, 1977) and reconnaissance geochemical sampling of selected areas by USGS. Regional ERDA aeromagnetic and gamma-ray profiles (1:500,000 scale)	Some creeks contain abundant barite boulders with fine-grained tetrahedrite, galena and sphalerite. Cassiterite and platinum group metals reported in Bear Creek placer concentrates. Area contains elements of area 12 above	Stream gold placers have been extensively mined locally. Production has been well over 260 kg (8,500 oz) of gold plus byproduct silver		

<u>AREA OUT- LINED ON MAPS</u>	<u>MAJOR TYPES OF KNOWN DEPOSITS</u>	<u>SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)</u>	<u>GEOLOGIC CONTROL(S) OF MINERAL RESOURCES</u>	<u>PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)</u>	<u>STATUS OF GEOLOGIC IN- FORMATION</u>	<u>ADDITIONAL COMMENTS</u>	<u>SUMMARY OF MINERAL RESOURCE POTENTIAL</u>	<u>ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)</u>	<u>GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)</u>
16.	(a)Au, Sn (Cr, W, RE's)--placer	(b)(Pb, Ag, Au, Cu)-- mineralized shear zones and veins	(a)Placer deposits re- stricted to several pay streaks in lower Quaternary stream gravels and gener- ally related to the drain- age basin that existed be- fore 3 to 52 m (10 to 170 ft) of silt and muck cov- ered the area. Definite lode sources for the gold and cassiterite have not been found anywhere in area. Placer gravels may have been derived from un- exposed quartz veins imme- diately north of the well defined tin belt. The veins may have been brec- ciated and mineralized with cassiterite and tourmaline during Tertiary intrusive activity. Cassiterite and gold need not have come from same source. Chro- mite, picotite, ilmenite and magnetite were probably derived from several ser- pentinized ultramafic bod- ies in area. (b)Mineralized shear zones in hornfels related to Tertiary granitic intru- sives	(a)Gold production 1899-1961 about 14,000 kg (451,000 oz, about 2.2% Alaska's placer output), plus sev- eral hundred tons byproduct cassiter- ite from belt about 19 km (12 mi) long and 1.6 km (1 mi) wide. 238 tons (262 short tons) cassiterite report- edly produced 1911- 1919. Wayland (USGS, 1961) esti- mated as of 1941 tailings contained about 201 tons (222 short tons) of cas- siterite (indicated reserves) and about 820 tons (900 short tons) of proven and indicated cassiter- ite reserves in buried placers in areas with system- atic drilling and near mined pay streaks. Addition- al inferred cassit- erite in the basins of Patterson and Baker Creeks and west of Woodchopper Creek amounted to about 1,000 tons 1,100 short tons). Reserves of gold in tailings were esti- mated to be about 130 kg (4,300 oz). Lode re- sources entirely speculative. Re- source estimates probably valid at least through 1958 (b)None of the lode deposits has been productive. Their resource potential appears to be small	Area covered by recent (1975) 1:250,000-scale regional geologic mapping. USGS contoured aeromag at 1:250,000 scale. Local detailed aeromag by indus- try. (a)Detailed studies and sampling programs in Tofty tin belt by USGS and USBM. (b)USBM diamond drilling and sampling program (1954) on Hot Springs Dome mineralized zones	(a)Rare earth min- erals (monazite, aeschnite, and xenotime) rather widespread among the placers but not concentrated into minable de- posits. Small amounts of schee- lite, cinnabar and native copper lo- cally present in concentrates	(a)Stream gold plac- ers have yielded about 14,000 kg (451,000 oz) of gold plus byproduct silver and several hundred tons of cassiterite (tin). Remaining cassiterite resources (all categories) probably do not exceed a few thousand tons of cassiterite. (b)Mineralized shear zones and quartz veins in contact zones of Tertiary granitic intrusives contain sulfide min- erals (Pb, Ag, Au, Cu). No production is known		

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
17.	(a)Au(Ag, Sn, W, Cr, Pb, Bi, Hg, RE's)-- placer (b)Sb(Ag, Pb, Cu, Mn)--vein	- - -	(a)Bench and present stream placers. Source of gold probably in mineralized quartz veins in schistose lower Paleozoic rocks and contact zones related to Upper Cretaceous-lower Ter- tiary plutons. Locally common chromite and some of the copper present in heavy concentrates probably derived from mafic and ultramafic rocks of the Rampart ophiolitic belt. (b)Small vein deposits bearing silver-lead, lead- gold-copper, and antimony may be related to Upper Cretaceous-lower Tertiary intrusive activity. Two manganese occurrences con- sist of irregular masses and thin seams of psilo- melane of probably hydro- thermal origin in lower Paleozoic chert and/or quartzite	(a)Production data incomplete but small- and large- scale placer mining operations probably have accounted for up to 3,100 kg (100,000 oz) of gold from the 1880's to present. Much of the gold quite pure (>900 fine). Potential for future mining not known but for most part little virgin ground re- mains. Some streams (e.g., Minook Creek) may warrant reevalua- tion for large- scale operations at elevated gold prices. (b)Seven known oc- currences. Only recorded lode pro- duction has been a few hundred tons of antimony ore in the early 1950's from a deposit on Sawtooth Mountain. Potential for future lode min- ing unknown but probably small. Area may contain plutons that have characteristics of tin granites	Recent (post 1970) 1:250,000-scale geologic mapping and regional geo- chemical sampling by USGS. Contoured aeromag at 1 mi spacing (USGS). ERDA aeromag and gamma-ray pro- files (1:500,000 scale)	(a)Noneconomic amounts of chro- mium, copper, lead, silver, tungsten, bismuth, mercury, and rare earth minerals present in concentrates of many of the plac- ers. (b)Geochemically anomalous amounts of tin, lead, sil- ver, copper, zinc, arsenic and rare earths present in bedrock and stream sediment samples in vicinity of gra- nitic intrusives	(a)Bench and stream gold placers have been mined. Produc- tion has been approx- imately 3,100 kg (100,000 oz) of gold. Potential for future production is unknown, but probably little virgin ground remains. (b)One of seven known polymetal vein depos- its has produced a few hundred tons of antimony ore		
18.	- - -	(Ni, Cr, Pt, Asbestos)-- Podiform chromite	Discontinuous belt about 113 km (70 mi) long and up to 3 km (1 3/4 mi) wide of tectonically emplaced alpine-type serpentinite contains occurrences of podiform chromite, minor cross-fiber veinlets of asbestos, and geochemically anomalous amounts of nickel with respect to similar rocks in other parts of the world	Grab rock samples contain up to 0.5% nickel and low, but detectable, amounts of platinum, palla- dium and rhodium. One selected sample of chromite con- tained 12.8% chro- mium. General ab- sence of sulfide minerals suggest the observed anoma- lous nickel content may be associated with silicate min- erals. Additional covered bodies of	Regional 1:250,000-scale geologic mapping by USGS (1971); USGS contoured aeromag (1:250,000 scale). USGS regional bedrock and stream sediment sampling. Much of belt staked by industry about 1970	Serpentinized ul- tramafic rocks may represent part of dismembered ophi- olitic complex	The occurrence of chromite and minor cross-fiber asbestos in serpentinite sug- gests that the re- gions have a poten- tial for the occur- rence of podiform chromite and asbes- tos deposits	90% 50% 10% 4 14 45 chance that there are deposits or more	podiform chromite model

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
18. (cont.)				tectonized serpen- tinite may occur on strike with or par- allel to the major ultramafic trend. Total resource po- tential not known					
19.	(a)Au--placer (b)Sb, Hg (Au, Ag)-- vein and breccia filling	(c){Sn, W}--possible vein, disseminated and/or tactite deposits (d){Au}--possible low- grade disseminated deposits	(a)Stream and bench gold placers representing modern and ancient stream-channel deposits developed on mature Quaternary erosion surface and largely buried by later sediments includ- ing silt and muck (frozen). Source of gold believed to be metalliferous quartz- calcite veins that consti- tute small lodes associated with numerous small quartz monzonite and monzonite (Tertiary?) intrusives. (b)Mainly small quartz- carbonate veinlets contain- ing gold, silver and stib- nite spatially related to monzonite stocks and dikes or probable Tertiary age. Altered granite at head of Olive Creek contains low- grade deposits of cinnabar. (c)The known local occur- rence of cassiterite, scheelite, topaz, tourma- line and rare earth miner- als in selected placer concentrates, as well as geochemically anomalous amounts of boron in stream sediment samples, within an area known to contain Ter- tiary granitic intrusives (including biotite granite) suggests the possibility that unrecognized or undis- covered tin granites with associated tin-tungsten vein deposits may be pres- ent. (d)The geologic setting and suite of geochemically anomalous metals (espe- cially arsenic, mercury and antimony) known to be pres- ent in certain of the drainage areas suggest the region may be permissive for the occurrence of low- grade disseminated gold deposits (Carlin-type)	(a)Livengood placer district has produced well over 12,000 kg (380,000 fine ounces) of gold (about 2% of Alaska's total gold placer produc- tion). One of the most recently de- veloped placer dis- tricts in state. Available data sug- gest district con- tains sizable gold resources that mer- it additional eco- nomic appraisal. Information from 640 drill holes in gravels of Liven- good Creek area alone indicates about 13 million m ³ (17 million yd ³) of material with average recoverable gold content of 0.8 g/m ³ [\$3.20/yd ³ (gold at \$5.14/g; \$160/oz)]. Area may contain as yet undiscovered plac- ers representing former stream chan- nels buried beneath frozen muck. (b)Of the eight known lode occur- rences, only two have had even token production. A small amount of stibnite was shipped from a vein on Livengood Creek and a little cinnabar was recov- ered from the de- posit at the head of Olive Creek. Stream sediment and spring sediment sam- ples locally contain anomalous amounts of	See area 18 above (a)Considerable company exploratory drilling of placers, especially in Livengood Creek area. (b)USGS bedrock, stream sediment and spring sedi- ment sampling in vicinity of Livengood (1967)	(a)Heavy minerals identified in con- centrates include chromite and chrome spinel (derived from ultramafics-- see area 18), cin- nabar, stibnite and less abundant scheelite, cassit- erite, and uranium- thorium-rare earth minerals	(a)Stream and bench placers have produced well over 12,000 kg (380,000 oz) of gold. Considerable poten- tial for additional production exists. Based on 640 drill holes, indicated re- sources in the Liven- good Creek area alone are equal to 13 x 10 ⁶ m ³ (17 x 10 ⁶ yd ³) at 0.8 g/m ³ [\$3.20/yd ³ ; gold at \$160/oz]. (b)Eight sulfide- bearing quartz- carbonate veins are known to occur within the region. Only two have produced even small amounts of ore. Some potential for additional deposits exists. (c)The presence of Tertiary granites, together with anoma- lous amounts of boron in stream sediment samples and the pres- ence of cassiterite, scheelite, topaz, tourmaline and rare earth minerals in placer concentrates, suggests a geologic setting permissive for the occurrence of tin and tungsten in veins, greisens and/or tactite deposits. (d)The region has an unknown potential for low-grade disseminated gold deposits (Carlin type)	(c) tungsten tactite mode	

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
-------------------------------------	----------------------------------	--	---	---	-------------------------------------	---------------------	--	--	--

19. (cont.)

gold, arsenic, mer-
cury, antimony,
molybdenum, silver
and boron, espe-
cially along fault
zones cut by Ter-
tiary intrusive
rocks. Although the
known lode deposits
have heretofore
proven uneconomic
because they are too
small and low grade
to mine, they sug-
gest the possibility
of additional con-
cealed deposits in
this poorly exposed
region of seemingly
favorable geology.
Some company explo-
ration for lode de-
posits.
(c)No production.
Definitive resource
data lacking.
(d)Definitive re-
source data lacking

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
20	(a)Au(Sn, W, Sb)-- placer (b)Au, (Ag)--vein (c)Sb--vein (d)W--contact meta- morphitic and vein	(e)(W, Sn)--possible vein, greisen, and/or contact metamorphic (tactite) deposits	(a)Principally stream placers formed in creek valleys in Pliocene and Pleistocene time during a complex series of events that included an early period of gold-bearing gravel deposition, fol- lowed by erosion and re- concentration of much of the gold from the earlier gravels into later placers that were offset from the location of the earlier channels. Some of the initial accumulations re- main as fragmentary bench deposits. Deposits buried beneath cover of later Quaternary loess and fro- zen silt and muck. Source of the gold and associated heavy metals believed to be local, mainly from min- eralized quartz veins associated with Upper Cre- taceous(?) lower Tertiary granitic intrusives. (b)Fissure veins 2 in. to 3 ft thick that cut schist and are spatially related to Upper Cretaceous-lower Tertiary granitic intru- sive rocks, including quartz diorite, biotite granite, and related por- phyries. (c)Stibnite and sulfanti- monide minerals generally present in the same quartz vein systems and geologic setting as the lode gold- silver deposits. However, at a few places, stibnite was deposited to the prac- tical exclusion of other sulfide minerals and gold. The deposits occur princi- pally in the Pedro Dome, Ester Dome, and Sourdough Creek areas in a belt about 21 km (13 mi) long northeast of Fairbanks. (d)Scheelite in tactite zones developed at contacts between pegmatitic granitic intrusive rocks and lenses and beds of marble and cal- careous schist. Scheelite	(a)1902 through 1963, Fairbanks area was largest placer producer in Alaska, yielding about 240,000 kg (7,650,000 fine ounces) of gold roughly (37% of total Alaskan placer production) plus byproduct sil- ver. Main produc- ing creeks were Cleary, Ester, Fairbanks, Dome, and Goldstream and their tributaries. Placer operations by USSR & M, once involving as many as six dredges, were closed down in 1964. Most of the potential ground has been exten- sively drilled by former operators. Some of the margin- al placer gold re- sources doubtless remain, especially in tailings and areas that could not be worked dur- ing former dredging operations, but quantitative re- source data are unavailable. (b)About 7,500 kg (240,000 oz) of gold and 1,200 kg (39,000 oz) of sil- ver produced to 1960 from 65 mines. Most of the silver occurs in lead sul- fantimonide and sulfide minerals. In some cases, gold content of closely spaced quartz vein- lets was sufficient to permit mining over zones 8-12 ft wide. Lode poten- tial unknown but probably moderate	Regional USGS 1:250,000- geologic mapping of Fair- banks (1966) and Livengood (1977) quadrangles, and 1:24,000-scale coverage of D-1, 2, 3 Fairbanks sub- quadrangles. Detailed USGS and University of Alaska studies of surfi- cial and bedrock geology, Fairbanks area. Contoured 1:250,000-scale aeromag- netic coverage by USGS and State. Area considered to be rather thoroughly pros- pected for gold, antimony, and tungsten deposits, but outcrops are poor owing to extensive cover of surfi- cial deposits. (b)(c)(d)Fair to good geo- logic information level exists as result of de- tailed examinations and exploration programs by government, industry, and University of Alaska	(a)Heavy minerals in placer concen- trates mainly the same as in lodes and include stib- nite, scheelite, wolframite, chrom- ite, and a wide variety of sulfide minerals. Locally, native bismuth, galena, native cop- per, cinnabar, and uranium, thorium and rare earth min- erals also present. No known lode source for cassi- terite which occurs in minor amounts in stream concentrates throughout area. Scheelite fairly common in several creeks but no known lode tungsten re- sources. (b)(c)(d)Current mine activity almost nil	(a)Stream and buried bench placers have produced large (240,000 kg; 7,650,000 oz) amounts of gold. The area has been extensively drilled by previous opera- tors and only marg- inal lands are thought to remain unworked (b)Production from 65 mines has been about 7,500 kg (240,000 oz) of gold and 1,200 kg (39,000 oz) of silver from gold bearing quartz veins. Potential for additional production unknown but considered to be moderate to small (c)Approximately 2,700 tons of anti- mony have been pro- duced from sulfide bearing quartz veins. Deposits generally contain small ton- nages of ore (d)Tungsten has been produced from tactite deposit and as a by- product from gold- quartz veins. The largest known deposit (Stepovich) produced 36 metric tons of W03. Some potential exists for additional tactite deposits but they are likely to be small and scattered (e)The presence of cassiterite in placer concentrates and gran- itic intrusions (in- cluding porphyritic varieties) suggests the area is geologi- cally permissive for the occurrence of tin granite with associated tin and tungsten vein deposits	90% 50% 10% chance that there are deposits or more	(d)tungsten tactite model

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
-------------------------------------	----------------------------------	--	---	---	-------------------------------------	---------------------	--	--	--

20 (cont.)

also present in auriferous quartz veins in schist and in granitic and pegmatic dikes.

(e) The presence of minor but ubiquitous amounts of cassiterite in placer concentrates of streams draining poorly exposed bedrock areas underlain by Late Cretaceous-early Tertiary biotite granite plutons with associated porphyries and local pegmatites that have intruded calcareous schist and marble suggests the area could conceivably contain as yet undiscovered tin granites with associated tin and tungsten vein or contact metamorphic deposits

to small.

(c) Total production of antimony ore has been about 2,700 metric (3,000 short) tons, mainly prior to World War II and from 18 of more than 50 localities known to contain antimony minerals. Most of the production was from five mines, two of which were operated mainly for antimony. The remaining production was a byproduct of gold mining. In 1942 USGS estimated about 27 tons (30 short tons) of stockpiled high-grade ore containing more than 30% antimony, about 2,700 tons (3,000 short tons) of low-grade dump material and about 45 tons (50 short tons) of proven high-grade material at various mines and prospects. Inferred resources not estimated, but doubtless present. Probably several hundred tons of ore shipped since 1942.

(d) Scheelite lode deposits known at about two dozen localities, all in Pedro Dome area. Largest known deposit (Stepovich) produced about 36 tons (4,000 units, 20 lbs to a short ton) of WO_3 through 1944, mainly from tactite zones developed at intersections of pegmatite dikes and marble. Small tonnages of scheelite concentrate also produced

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
20. (cont.)				as byproduct from gold-quartz veins. Known resources are small. Area may contain undiscover- ed deposits, but they are likely to be small and sparsely distrib- uted					
21.	(a)Au, Sn, W (Pb, Bi, Th, U, RE's)-- placer	(b)Au, Ag, Pb, Cu-- vein and vein networks (c)(Sn, W, Cu, Pb, Mo, Th, RE's)--vein; dis- seminations in ig- neous rocks; por- phyry(?)	(a)Stream and bench plac- ers 1.2-4.6 m (4-15 ft) thick buried beneath 0-3 m (0-10 ft) of muck (mostly unfrozen). Gold believed locally derived from min- eralized quartz veins (some of which contain visible gold) in schist and Upper Cretaceous- lower Tertiary granitic plutons that constitute the bedrocks of the drainage basins. (b)Quartz veins and vein networks in schist, spa- tially related to small Upper Cretaceous-lower Tertiary intrusives, con- tain gold, argentiferous galena, and malachite. (c)Bedrock samples from small Upper Cretaceous- lower Tertiary intru- sives, including both equigranular and porphy- ritic varieties of bio- tite granite, contain small amounts of schee- elite, galena, molybde- nite, stibnite, monazite, and allanite, with acces- sory fluorite and topaz. Associated quartz veins contain scheelite, wol- framite, and cassiterite. The primary and second- ary (placer) occurrence of these minerals in the indicated geologic set- tings suggests that the area may contain unrecog- nized and/or undiscovered porphyry copper deposits and tin granites with associated tin-tungsten veins [see (c), additional com- ments column].	(a)Gold placers have produced about 26,000 kg (835,000 fine ounces) or more of gold since their discovery in 1893. Many of the placers contain subeconomic amounts of cassiterite which was recovered with gold and wol- framite at Deadwood Creek. (b)No known pro- duction from three occurrences. Known resources thus are small but potential exists for the dis- covery of additional deposits. (c)Three known oc- currences. No pro- duction or known development work. (d)Definitive resource data lacking	Area poorly known geolog- ically. Little or no geochemical control. Re- cent USGS (1974) contoured aeromagnetic coverage (1:250,000 scale). Uni- versity of Alaska thesis work in Mount Prindle area. (d)Reconnaissance mineral sampling program by USBM/ BLM/USGS (1977)	(a)Heavy concen- trates also locally include wolframite, allanite, bismuth- enite, galena, cinnabar, xenotime, topaz, and tourma- line. Some of these are slightly uraniferous. (b)... (c)At Mount Prin- dle, lower Tertiary porphyritic, fine- to coarse-grained biotite granite pluton is cut by aplite, quartz porphyry, and tour- maline-bearing peg- matite dikes, and contains vein fluorite, as well as anomalously high tin and boron	(a)Stream and bench placers have produced about 26,000 kg (835,000 oz) of gold (b)Though none has produced, two pros- pects and one occur- rence of gold bearing quartz veins are known in the region. There is some potential for the discovery of add- itional deposits of small tonnage (c)The rock (porphyri- tic granites) and the minerals present in bedrock samples (scheelite, galena, molybdenite, stibnite, monazite, allanite, fluorite and topaz) suggest a setting which is geologically permissive for the occurrence of porphyry copper deposits and possibly tin-bearing granites. (d)Values of uranium (up to 400 ppm) in stream sediments from the contact area of the Mount Prindle pluton suggests that some of the granitic intru- sions of the region have some potential for uranium	(c)porphyry copper model	

AREA OUT- LINE ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDING MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
------------------------------------	----------------------------------	---	---	---	-------------------------------------	---------------------	--	--	--

21. (cont.)			(d) Anomalous uranium detected in stream sediment (up to 400 ppm) and spring sediment (up to 570 ppm) samples from contact zone of Mt. Prindle granitic pluton (early Tertiary) [see (c), additional comments column] suggests that this and similar intrusives have some potential for disseminated uranium						
22.	(a) Au(Sn, W, Bi)-- placer	(b) (Cr, Ni)--possible magmatic deposits (c) (Sn, W, Bi)--possible vein, greisen, and/or tactite deposits	(a) Buried stream placers. Gold and subeconomic amounts of scheelite, cassiterite and bismuth, probably derived from quartz-carbonate veinlets in Paleozoic and/or Precambrian(?) schist. (b) Area contains several known small ultramafic intrusive bodies with a virtually unknown but probably small potential for chromium, cross-fiber asbestos, and nickel. (c) Presence of small but ubiquitous amounts of cassiterite, locally abundant scheelite, and bismuth, as well as accessory tourmaline in some stream concentrates, suggests the possibility of as yet undiscovered tin-tungsten vein, greisen and/or tactite deposits especially associated with Upper Cretaceous-lower Tertiary granitic intrusives known to occur in the area	(a) Total gold production from ten known deposits not known but probably did not exceed 310 kg (10,000 oz). Stream concentrates from Salcha River drainage area contain locally abundant native bismuth and scheelite, but there is no record of either having been recovered. Cassiterite also present in concentrates. (b) Definitive resource data lacking. (c) Early AMRAP stream-sediment sample results show anomalous tin values	Area currently being studied under USGS AMRAP program. State and USGS contoured aeromagnetic coverage (1:250,000 scale). Area believed to have been thoroughly prospected but mainly for gold placers. Bedrock exposures generally poor below 2,000-3,000 ft altitude due to heavy vegetation and cover of surficial deposits	(b) General absence of sulfide minerals suggests that any nickel present is likely to be silicate phases	(a) Ten known placer deposits have produced less than 310 kg (10,000 oz) of gold. The region has been extensively prospected but mainly for gold (b) The presence of small bodies of ultramafic rock suggests the area is permissive for the occurrence of podiform chromite and cross-fiber asbestos deposits (c) The rock types present (granitic intrusions into locally calcareous schist), together with the presence of cassiterite and tourmaline in stream concentrates suggest the area is permissive for the occurrence of tin and tungsten in veins or greisen and/or tactite zones		(c) tungsten tactite model

[illegible]

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
24. (cont.)			significant resources is considered small. (d)Gold-bearing quartz- carbonate veins asso- ciated with mafic and ultramafic igneous rocks.	up to 5,000 ppm nickel and up to 7,000 ppm arsenic. Since no sulfide mineral phases re- cognized, nickel probably present in silicate mineral. Silicate-carbonate rock also locally contains traces of gold (0.9 ppm) and silver (0.5 ppm). (c)Bedrock samples from Eagle Bluff contain 7 to >10,000 ppm copper, <10 to 1,500 ppm lead, <200 to 1,500 ppm zinc, <5 to 2,000 ppm cobalt, <0.5 to 1.5 ppm silver and <10 to 500 ppm tin. (d)Two known occur- rences. Samples across one zone 19 m (63 ft) wide yielded up to 11 ppm gold. Undis- covered occurrences probably exist, but they are likely to be small and of negligible poten- tial					
25.	(a)Au(Pt, RE's)-- placer	(b)(U, Th)--possible peneconcordant depos- its(?)	(a)Modern stream placers and less productive bench placers localized within a belt of Upper Creta- ceous to Pliocene (?) non- marine sandstone, mud- stone, and quartz-rich conglomerate that strad- dles the Tintina fault zone and from which most of the placer gold is believed to have been reconcentrated. Irregu- lar distribution of the gold in modern placers probably related to ini- tial erratic distribu- tion in the older con- glomerates. Original source of gold considered at least in part to be quartz veins in meta- morphitic and granitic intrusive rocks that	(a)Total production not known but prob- ably amounts to at least 930 kg (30,000 fine ounces) of gold, plus byprod- uct silver. Gold also alloyed with as much as 0.42% platinum metals, claimed as seign- iorage by U.S. Treasury. Gold placer remaining resources not known. Although gold is present in the non- marine Late Creta- ceous to Pliocene conglomerates, it is sparsely and erratically distri- buted. (No known gold placer	Entire area covered by recent (1969, 1972) USGS 1:250,000 geologic map- ping. Eastern part near Eagle covered by 1:63,360- scale USGS geologic map- ping. Part in Eagle quadrangle covered by state contoured aeromag (1:250,000 scale); that in Charley River quad- rangle covered by 1:1,000,000 scale con- toured aeromag and ERDA 1:500,000 aeromag and gamma ray profiles. USGS geochemical sampling in Eagle quadrangle	Placer concentrates locally include native platinum, native silver, cin- nabar, monazite, cassiterite, and chromite. Bedrocks in Tintina fault zone contain local- ly high but sporadic concentrates of copper, lead, zinc, and tin. (b)Selected sam- ples of black shale from Calico Bluff formation (Upper Mississip- pian-Lower Pennsyl- vanian) contain 0.001-0.009% uran- ium, up to 0.68% V2O5 and 1.9% P2O5	(a)Stream and bench deposits have proba- bly produced at least 930 kg (30,000 oz) of gold plus byproduct silver (b)The lithology and organic content of the Upper Cretaceous-Ter- tiary conglomerates together with the presence of small amounts of uranifer- ous, thorium and rare earth minerals in some of the pla- cer deposits suggest a geologic setting permissive for the occurrence of sand- stone uranium deposits		

AREA OUT- LINED ON MAPS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)	STATUS OF GEOLOGIC IN- FORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)
25. (cont.)			underlie region to south. However, not all lithic constituents in the conglomerate are represented in that igneous and metamorphic bedrock terrane. Platinum locally present in heavy concentrates probably derived from ultramafic intrusives	occurrences in northeastern area.) (b)Geologic relationships, including the presence of small amounts of uraniferous thorium and rare earth minerals in some of the placer concentrates suggest that the areas of continental, coal-bearing clastic rocks may be permissive for the occurrence of peneconcordant uranium deposits					
26.	(a)Fe--sedimentary	---	(a)Low-grade banded hematitic red beds of sedimentary origin occur in Tindir Group (late Precambrian). Hematite occurs in beds a fraction of an inch to 1 in. thick, as a replacement mineral of volcanic fragments, as a cement, and as a matrix constituent	Section considered to have best potential about 550 m (1800 ft) thick. (a)USBM sampling of 40 and 60 m (133 and 200 ft) intervals judged to have highest iron content showed 20.10% and 21.85 soluble iron, respectively. One 2 cm (3/4 in.) bed arrayed 33.4% soluble iron, the maximum tenor found. Hypothetical and speculative resources could amount to several billion tons or more	Recent (1969) 1:250,000-scale geologic mapping by USGS. USGS aeromag coverage at 1:1,000,000 scale. ERDA aeromag and gamma-ray profiles (1:500,000 scale). USBM sampling program in Tatonduk River area		Low-grade hematitic beds of the Tindir Group occur in this region. The portion of the stratigraphic section with the greatest known resource potential is 550 m (1800 ft) thick. Sampling of two portions of the section judged to have the highest iron content showed approximately 20% soluble iron. Speculative and hypothetical resources could amount to several billion tons		
27.	(a)Au(Sn, W, Hg, Cr, Ag, RE's)--placer	(b)Au, Ag, Pb, Cu, Zn, Sb--vein (c)(Cu, Pb, Zn)--possible stratabound volcanogenic(?) massive and/or disseminated sulfide deposits. (d)(Sn, W)--possible vein, greisen, and/or tactite deposits	(a)Stream and bench placers in zone of discontinuous permafrost. Much of ground permanently frozen. Loess, derived from proglacial flood plains, mantles much of area. Source of gold considered to be mineralized quartz veins in Paleozoic metamorphic rocks near contacts with felsic intrusives, mainly of Late Cretaceous-early Tertiary age.	(a)Approximately 13,000 kg (417,000 oz) (about 2% total Alaska placer gold production) of gold produced from 1887 through 1961. Production figures not available since 1961, but large dredge operated at Chicken Creek through 1967. No reliable	Area of generally poor exposures covered by recent (1976) 1:250,000-scale USGS geologic mapping. Most of area covered by 1965 or later 1:63,360 USGS geologic mapping. Regional USGS geochemical sampling control generally good except southeastern part. State contoured aeromag coverage (1:250,000 scale). Area has been target of considerable	(a)Stream concentrates locally include noneconomic amounts of scheelite, cassiterite, cinnabar, chromite, native silver, monazite and allanite. Lode source of the cinnabar not known. Area also contains small serpentized ultramafic intrusive bodies considered to have a	(a)Stream and bench placers have produced approximately 13,000 kg (417,000 oz) of gold. Small-scale (<3) kg/yr; 1000 oz/yr operations continue to produce gold in 1977. Area has been actively prospected for almost 100 years. Potential for undiscovered stream	(c)felsic and intermediate volcanogenic sulfide model. (d)tungsten tactite model	

<u>AREA OUT- LINED ON MAPS</u>	<u>MAJOR TYPES OF KNOWN DEPOSITS</u>	<u>SUSPECTED OR SPECULA- TIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES</u>	<u>GEOLOGIC CONTROL(S) OF MINERAL RESOURCES</u>	<u>PRODUCTION AND RE- SOURCE INFORMATION (ALL UNITS METRIC UNLESS OTHERWISE INDICATED)</u>	<u>STATUS OF GEOLOGIC IN- FORMATION</u>	<u>ADDITIONAL COMMENTS</u>	<u>SUMMARY OF MINERAL RESOURCE POTENTIAL</u>	<u>ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS</u>	<u>GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 3)</u>
27. (cont.)		<p>(b)Polymetal quartz-carbonate veins in meta-morphic rocks (Paleozoic and/or Precambrian?) near contacts with granitic intrusive rocks (mainly Upper Cretaceous-lower Tertiary).</p> <p>(c)Although no such deposits are presently known, the area contains geologic elements (including metamorphosed rocks of probable felsic to intermediate volcanic origin) considered permissive for the occurrence of stratabound (volcanogenic?) base metal sulfide deposits.</p> <p>(d)The presence of ubiquitous scheelite and cassiterite in several of the gold placers suggests that the area has some potential for the occurrences of as yet undiscovered tin-tungsten vein, disseminated, or tactite deposits, especially in the vicinity of composite intrusives of Upper Cretaceous-lower Tertiary porphyritic biotite granite</p>	<p>(b)Polymetal quartz-carbonate veins in meta-morphic rocks (Paleozoic and/or Precambrian?) near contacts with granitic intrusive rocks (mainly Upper Cretaceous-lower Tertiary).</p> <p>(c)Although no such deposits are presently known, the area contains geologic elements (including metamorphosed rocks of probable felsic to intermediate volcanic origin) considered permissive for the occurrence of stratabound (volcanogenic?) base metal sulfide deposits.</p> <p>(d)The presence of ubiquitous scheelite and cassiterite in several of the gold placers suggests that the area has some potential for the occurrences of as yet undiscovered tin-tungsten vein, disseminated, or tactite deposits, especially in the vicinity of composite intrusives of Upper Cretaceous-lower Tertiary porphyritic biotite granite</p>	<p>estimate of amount of gold mined annually since then, but it probably has not exceeded 31 kg (1,000 oz), even in good years. At least 20 known active placer operations, 1977. Potential for undiscovered modern stream placers probably small because district has been actively prospected for almost 100 years. Remaining gold resources mainly are in ground that could not be profitably worked earlier at lower gold prices (especially dredged area) and in tailing piles. Principal virgin placer ground lies in high level terraces, which could contain rich pay streaks (e.g., Napoleon Creek).</p> <p>(b)Of six known occurrences, only one is known to have recorded production from a rich gold-quartz vein which extended to depth of about 2 m (6 ft). Resource potential of such vein deposits not known but believed to be small</p>	recent prospecting and exploration activity by industry	low potential for cross-fiber asbestos, chromite, and possibly nickel	<p>placers is probably low. Potential does exist for undiscovered placers in high-level terraces.</p> <p>(b)Six occurrences of sulfide-bearing quartz veins are known. One has produced a small quantity of gold.</p> <p>(c)The presence of metamorphosed felsic and intermediate volcanic rocks suggests a geologic setting permissive for the occurrence of felsic and intermediate volcano-genic sulfide deposits.</p> <p>(d)The presence of scheelite and cassiterite in placers, together with the occurrence of composite intrusives of granitic composition, suggest a geologic setting permissive for the occurrence of tin and tungsten veins, greisen and/or tactite zones</p>		

[illegible]

TABLE 3. GRADE AND TONNAGE MODELS

[Related data occur on line from column to column: all data in metric units;
NS, not significant; *, significant at 5-percent level; **, significant at 1 percent level]

Deposit Type	Tonnage and grade variables (units in parenthesis)	Number of deposits used in developing model	Correlation coefficient of listed variable with variable on line with it in column 2	90 percent of deposits have at least	50 percent of deposits have at least	10 percent of deposits have at least
Porphyry Copper	Tonnage of ore (millions of tons)	41		20	190	430
	Average copper grade (percent)	41	with tonnage of ore = -0.07 NS	0.1	0.3	0.55
	Average molybdenum grade (percent Mo)	41		0.0	0.003	0.031
Island Arc Porphyry Copper	Tonnage of ore (millions of tons)	41		20	190	430
	Average copper grade (percent)	41	with tonnage of ore = -0.07 NS	0.1	0.3	0.55
	Average molybdenum grade (percent Mo)	41		0.0	0.008	0.031
	Average gold grade—locally significant but not determined					
Porphyry Molybdenum	Tonnage of ore (millions of tons)	31		1.5	24	340
	Average molybdenum grade (percent Mo)	31	with tonnage of ore = -0.05 NS	0.065	0.13	0.26
Podiform Chromite	Tonnage of Cr_2O_3 (tons)	268		15	200	2,700
Copper Skarn	Tonnage of ore (millions of tons)	38		0.08	1.4	24
	Average copper grade (percent)	38	with tonnage of ore = -0.44**	0.86	1.7	3.5
	Average gold grade—locally significant, but not determined					
Mafic Volcanogenic	Tonnage of ore (millions of tons)	37		0.24	2.3	22.0
	Average copper grade (percent)	37	with tonnage of ore = -0.13 NS	1.1	2.2	4.1
	Average zinc grade excluding deposits without reported grades (percent)	19	with tonnage of ore = 0.03 NS	0.3	1.3	5.5
	Average gold grade—locally significant but not determined					
Felsic and Intermediate Volcanogenic Massive Sulfide	Tonnage of ore (millions of tons)	89		0.19	1.9	18.0
	Average copper grade (percent)	89	with tonnage of ore = -0.41**	0.54	1.70	5.40
	Average zinc grade excluding deposits without reported grades (percent)	41	with tonnage of ore = 0.25 NS	1.40	3.80	10.00
	Average lead grade excluding deposits without reported grades (percent)	14	with tonnage of ore = -0.02 NS	0.20	0.95	4.80
	Tonnage contained gold excluding deposits without reported gold (tons)	38	with tonnage of ore = 0.78**	0.27	2.90	32.00
	Tonnage contained silver excluding deposits without reported silver (tons)	46	with tonnage of ore = 0.82**	5.00	30.00	1300.00
Nickel Sulfide	Tonnage of ore (millions of tons)	48		0.23	1.20	5.90
	Average nickel grade (percent)	48	with tonnage of ore = -0.03 NS	0.32	0.61	1.20
	Average copper grade (percent)	48	with tonnage of ore = 0.03 NS with nickel grade = 0.04 NS	0.13	0.47	1.20
Mercury	Tonnage of contained mercury (tons)	165		0.09	3.10	120.00
Vein Gold	Tonnage of contained gold (tons)	43		0.29	3.30	33.00
Skarn/Tactite Tungsten	Tonnage of ore (millions of tons)	31		0.024	0.63	17
	Average tungsten grade (percent W)	31	with tonnage of ore = -0.34 NS	0.24	0.51	1.10