
UNITED STATES DEPARTMENT OF THE INTERIOR GEOLOGICAL SURVEY

PRELIMINARY GEOMAGNETIC DATA COLLEGE OBSERVATORY FAIRBANKS, ALASKA

MARCH 1987

OPEN FILE REPORT 87-0300C



THIS REPORT WAS PREPARED UNDER THE DIRECTION OF JOHN B, TOWNSHEND, CHIEF OF THE COLLEGE OBSERVATORY, WITH THE ASSISTANCE OF THE OBSERVATORY STAFF MEMBERS: R.V. O'CONNELL AND L.Y. TORRENCE AND IN COOPERATION WITH THE GEOPHYSICAL INSTITUTE OF THE UNIVERSITY OF ALASKA, THE COLLEGE OBSERVATORY IS A PART OF THE BRANCH OF GLOBAL SEISMOLOGY AND GEOMAGNETISM OF THE U.S. GEOLOGICAL SURVEY.

Explanation of Data and Reports

Magnetic Activity Report

Principal Magnetic Storms

Preliminary Calibration Data and Monthly Mean Absolute Values

Magnetogram Hourly Scalings - Five Quietest Days

Sample Format for Normal and Storm Magnetograms

Normal Magnetograms

Storm Magnetograms (When Normal is too disturbed to read)

COLLEGE OBSERVATORY PRELIMINARY GEOMAGNETIC DATA

EXPLANATION OF DATA AND REPORTS

INTRODUCTION

The preliminary geomagnetic data included here is made available to scientific personnel and organizations as part of a cooperative effort and on a data exchange basis because of the early need by some users. To avoid delay, all of the data is copied from original forms processed at the observatory; therefore, it should be regarded as preliminary. Inquiries about this report or about the College Observatory should be addressed to:

Chief, College Observatory U.S. Geological Survey 800 Yukon Drive Fairbanks, Alaska 99775-5160

Requests for copies of the magnetograms except for the current month should be addressed to:

> World Data Center A NOAA D63m 325 Broadway Boulder, Colorado 80303

OBSERVATORY LOCATION

The College Observatory, operated by the U.S. Geological Survey, is located at the University of Alaska, Fairbanks, Alaska. It is near the Auroral Zone and the northern limit of the world's greatest earthquake belt, the Circum-Pacific Seismic Belt. Although the observatory's basic operation is in geomagnetism and seismology, it cooperates with other scientists and organizations in areas where the facility and personnel can be of service.

The observatory is one of three operated by the USGS in Alaska. The others are located at Barrow and Sitka.

GEOMAGNETIC DATA

Normal and storm magnetograms and appropriate calibration data are processed at the observatory and are available for analysis or copying. Also available are mean hourly scalings for the five quietest days for the month and K-Indices.

Magnetic Activity

The K-Index: The K-Index is a logarithmic measurement of the range of the most disturbed component (D or H) of the geomagnetic field for eight intervals 0000-0300, 0300-0600...2100-2400 UT. It is a measure of the difference between the highest and lowest deviation from a smooth curve to be expected for a component on a magnetically quiet day, within a three hour interval.

The Equivalent Daily Amplitude, AK: The K-Index is converted into an equivalent range, ak, which is near the center of the limiting gamma ranges for a given K. The average of the eight values is called equivalent daily amplitude AK. The unit 10y has been chosen so as not to give the illusion of a accuracy not justified.

The schedule for converting gamma range to K, and K to ak is as follows:

Centres	Range	K - Index	ak
<u>~</u>	25	0	~0
25<	50	1	3
50<	100	2	7
100<	200	3	15
200<	350	4	27
350<	600	5	48
600<	1000	6	80
1000<	1650	7	140
1650<	2500	8	240
2500+		9	400 (10y)

Principal Magnetic Storms

Gradual and sudden commencement magnetic disturbances with at least one K-Index of 5 or greater, which are believed to be part of a world-wide disturbance, are classified as principal magnetic storms. The time of the storm beginning and ending; direction and amplitude of sudden commencements; period of maximum activity; and storm range are reported. Monthly reports of these data are forwarded to the World Data Center A in Boulder, Colorado.

Magnetogram Hourly Scalings

Magnetogram bourly scalings are averaged for successive periods of one hour for the D, H, and Z elements. The Value in the column headed "Ol" is the average for the hour beginning 0000 and ending 0100. Note that the values on the scaling sheet are in tenths of mm with the decimal point omitted. The user of these scalings should keep in mind that the tabular values are hourly means and if one is interested in the detailed morphology of the magnetic field, refer directly to the magnetograms.

Magnetograms

The normal magnetograms in this report are reproduced at about one-third the size of the originals. Preliminary base-line values and scale values adopted for use with the original magnetograms are included. For days when the magnetic field is too disturbed for the Normal magnetogram to be readable, Storm magnetograms are reproduced.

Absolutes, Base-lines and Scale Values

To determine the absolute value of the magnetic field from the hourly means or from point scalings the following equations should be used:

D=B_D+d·S_D; H=B_H+h·S_H; Z=B_Z+z·S_Z
where D, H and Z are absolute values;
E_D. B_H and B_Z are base-line values;
S_D. S_H and S_Z are scale values;
and d, h and z are scalings in millimaters.

NUAA FORM 76-133 NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION

OBSERVATORY

MAGNETIC ACTIVITY
(Greenwich civil time, counted from midnight to midnight)

College Alaska

MONTH AND YEAR

March 1987

					K-10	OICE	5				TIME SCALE ON
DATE		93-06	6 0- 9 0	09-12	12-18	5 ¥ 10	18-21	21-24	\$UM	AK	MAGNETOGRAMS 20 mm/hr
1 2 3 4 5	1 0 0 2 1	2 0 0 1 3	1 2 2 0	2 1 4 4 6	2 0 1 4 6	3 1 2 3 5	2 1 1 3	2 0 1 1 3	15 04 11 18 27	07 02 06 12 32	SUDDEN COMMENCEMENTS d h , m
6 7 8 9	2 3 2 1 0	1 2 2 3 2	2 6 3 2 6	4 5 5 1 4	2 6 4 0 6	0 1 5 0 5	i 2 1 1	2 1 2 1 0	14 26 24 09 24	08 30 20 04 31	
11 12 13 14 15	2 1 2 2 2	1 2 1 3 3	1 2 3 1 2	3 5 4 4 3	4 2 3 3 3	3 1 2 1	3 2 1 1 2	1 2 1 1	18 17 17 16 17	11 11 10 10 09	
16 17 18 19 20	2 2 1 2	1 4 0 2 0	3 2 1 0	6 1 0 5 1	6 3 1 4 1	1 3 1	2 1 3 1 0	1 0 2 1 0	22 15 12 17 04	25 09 06 13 02	POSSIBLE SOLAR-FLARE EFFECTS BASED ON INSPECTION OF GRAMS ALONE (WITHOUT
21 22 23 24 25	0 3 2 0 0	0 1 1 0 0	2 3 1 2 0	5 5 3 0	5 2 3 0	7 2 1 1 3	3 1 1	2 2 0 1	24 21 12 05 05	33 15 06 02 03	REFERENCE TO DATA FROM OTHER SOURCES) BEGIN END d h m d h m
26 27 28 29 30 31	2 1 3 0 0	2 4 3 1 0 1	3 4 4 2 1	5 7 5 4 1 4	4 5 3 0 3 4	4 6 2 0 2 0	1 4 1 1 1	1 2 0 0 0	22 33 21 08 08 08	17 45 16 05 04 08	

K SOALE USED:	D	н	Z	
LOWER LIMIT FOR K = 9	675.7	322.2		(mm) ·
CURRENT SCALE VALUE	3.70.	7.80		(γ/mm)
LOWER LIMIT FOR K # 9	2500	2510		(to pearest 107)

SCALINGS AND COMPUTATIONS HAVE BEEN CHECKED

APPROVED John B. Townshend, Chief, College Observatory

OBSERVER IN CHARGE

PRINCIPAL MAGNETIC STORMS

WOC-A FOR GOLAR-TERRESTRIAL PHYSICS ENVIRONMENTAL DAYA SERVICE, NOAA BGULDER, COLORADO SOBGE U.S.A.

Data from Individual Observatories:


COLLEGE OBSERVATORY, COLLEGE, ALASKA
March 1987

	,	,				March		19 <u>87</u>							
Obs. Geomag.			Commencement			amplit	udea '	Max. 3 hr - index K				Ranges		UT End	
2 Infter IAGA code lat.	day	hr min (UT)	type	D(')	H(Y)	2(Y)	day	(3 hr -	- pariod)	K	(י)מ	μ (γ)	Z(Y)	day br	
CO	64.06 N	No I	rincipal	Magnet:	c Storms	Noted	During	This Mo	nth				,		
		,													
	ļ														
												,			
	l]										•	
					J						:				
								j		1	į.				
				.											
										_					
										•					r
	1						l		} .			j			l .


	PER	MORNAL MAGRETON	TRAPE	CALIDRA	MY 6.7		
COMPONENT		100 ·	BCALE 1		BASILLINE		
	FROM	J					
	0000117,3/1/87	2400 UT, 3/31/87	1.0/mm	3.78/mm	27° 0.3' E		
α							
					,		
	0000UT, 3/1/87	2569 UT, 3/21/87	7.8	8/mm	1262		
ı	0000 UT, 3/22/87	1400 UT, 3/31/87	7.8	mm	12632		
H	Jesus Diff. Clark	, 3/3//					
z	0000 UT, 3/1/87	2359 UT, 3/21/87	7.7	8/mm	55174		
	0000 UT, 3/22/87	2400 UT, 3/31/87	17.	1/mm	<i>5</i> 5170		
	, ,						
		STORM MAGRETOG	RAPE				
COMPONENT	YER			CALTER			
Cital cartai	PROM	70	SCATE		HASELINE		
	0000 UT, 3/1/87	2400UT, 3/31/87	7.9 mm	39.5 /mm			
מ							
			1 4~	78/mm			
	(same)	(same)	43.	7°/mm			
B							
	(same)	(bame)	41	8.7 8/mm			
Z	(***)	_ ()		/ / / / / / / / / / / / / / / / / / / /			
L							
		RAPID RUN MAGNE	TOGRAPH				
	PS	OED)		CALIBR	ATTON		
COMPONENT	FROM	TO		SCALE	VALUE		
p							
-			· ·				
· · ••					1111		
Ħ			+				
			 				
							
Z							
		MORTHLY MEAN ABSOLUT	VATTROA				
			- TAMES	_			
	D	R O C O Y	Z				
	" ACA' E	12862			5316 4		
17	UTED PRIM FIVE QUIETEST DAY:				2016		


Compare Company of Company and the state of the Company of the company


FORMAT FOR NORMAL & STORM MAGNETOGRAMS (SAMPLE ONLY)





SEE PRELIMINARY CALIBRATION DATA FOR SCALE VALUES & BASELINE VALUES
























MAGNETOGRAMS

STORM



MAGNETOGRAMS

STORM