

Jurassic Paleobiogeography of Alaska

By RALPH W. IMLAY and ROBERT L. DETTERMAN

GEOLOGICAL SURVEY PROFESSIONAL PAPER 801

Changes in the paleogeography of Alaska during Jurassic time are revealed by the distribution, succession, and differentiation of molluscan faunas and by the characteristics of the associated sedimentary rocks



UNITED STATES DEPARTMENT OF THE INTERIOR

ROGERS C. B. MORTON, *Secretary*

GEOLOGICAL SURVEY

V. E. McKelvey, *Director*

Library of Congress catalog-card No. 73-600141

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C. 20402 — Price 65 cents domestic postpaid or 45 cents GPO Bookstore
Stock Number 2401-00347

CONTENTS

	Page		Page
Abstract	1	Fossil occurrences by stages	21
Introduction	1	Hettangian	21
Acknowledgments	2	Sinemurian	22
Distribution of Jurassic rocks	2	Pliensbachian	22
Comparisons of stratigraphic and lithologic features--	8	Toarcian	23
Southern Alaska	8	Bajocian	23
Northern Alaska	9	Bathonian	24
Occurrences of Jurassic unconformities	14	Callovian	24
Southern Alaska	14	Early Oxfordian	25
Northern Alaska	16	Late Oxfordian to early Kimmeridgian	25
Extent of Jurassic seas	16	Late Kimmeridgian to early middle Tithonian	26
Succession and differentiation of ammonites	20	Late middle to late Tithonian	27
		References	28
		Index	33

ILLUSTRATIONS

		Page
FIGURES	1-9. Maps showing distribution of Jurassic fossils in Alaska:	
	1. All stages of the Jurassic	2
	2. Hettangian and Sinemurian	3
	3. Pliensbachian and Toarcian	3
	4. Bajocian	4
	5. Bathonian and Callovian	4
	6. Early Oxfordian	5
	7. Late Oxfordian to early Kimmeridgian	5
	8. Late Kimmeridgian to early middle Tithonian	6
	9. Late middle to late Tithonian	6
	10. Index map of Jurassic areas in Alaska	7
	11. Correlation charts of Jurassic rocks in Alaska	10
12-15.	Chart showing—	
	12. Occurrences of unconformities in the Jurassic of Alaska	15
	13. Correlation of Hettangian to Toarcian ammonite faunas in Alaska	17
	14. Correlation of Bajocian to Callovian ammonite faunas in Alaska	18
	15. Correlation of Oxfordian to Tithonian molluscan faunas in Alaska	19

JURASSIC PALEOBIOGEOGRAPHY OF ALASKA

By RALPH W. IMLAY and ROBERT L. DETTERMAN

ABSTRACT

Jurassic marginal seas occupied considerable areas in southern and northern Alaska and in the western part of the Kuskokwim region of southwestern Alaska. They appear to have been absent during late Callovian time, much restricted during Hettangian, Bathonian, early Oxfordian and late Tithonian time, and most extensive during Sinemurian, Bajocian, and late Oxfordian to middle Tithonian time. A large area in central Alaska was probably never covered. A southwestern prolongation of that area from the Talkeetna Mountains westward to the western end of the Alaska Peninsula was the site of granitic intrusions during late Early Jurassic time and of extensive erosion during Middle and Late Jurassic time. Variations in the rate of uplift of the area of these granitic intrusive rocks may explain why marine transgressions and regressions were at different times in southern than in northern Alaska during the Bajocian and Bathonian. Connection of the northern and southern marginal seas occurred through Yukon Territory and easternmost Alaska.

The Jurassic ammonite succession in Alaska is similar to that in central and northern Europe and northern Asia. In Lower Jurassic beds, it is essentially identical. In Bajocian and in Oxfordian to lower Kimmeridgian beds, the ammonite succession in Alaska differs from that in the other areas mainly by the presence of some genera found only in areas bordering the Pacific Ocean and by the absence of a few genera common in central and northern Europe. In contrast, the Bathonian rocks of Alaska contain ammonites, such as *Arcticoceras*, *Arctocephalites*, and *Cranocephalites*, that are widespread in the Arctic region but are unknown in central Europe. Comparisons with the Tithonian of Europe are not possible because ammonites of that age, other than *Lytoceras* and *Phylloceras*, are not yet known from Alaska. The Alaskan Jurassic ammonites of late Pliensbachian Age and of Bathonian to early Kimmeridgian Age belong mostly to the Boreal realm and have very little in common with Tethyan realm ammonites such as those found in areas bordering the Mediterranean Sea.

INTRODUCTION

This report deals in a broad way with changes in geography, stratigraphy, and ammonite assemblages in Alaska during Jurassic time. It is based on data of which at least half has been obtained during the last 25 years; it presents maps depicting the approximate positions and distribution of megafossil

occurrences of the various stages; and it describes those fossil occurrences or refers to publications in which they have been described. In addition, the report summarizes existing knowledge concerning gross stratigraphic and lithologic features, position and duration of unconformities, extent and changes in marginal Jurassic seas, succession and differentiation of ammonite faunas, comparisons with ammonite successions elsewhere, and possible relationship of ammonite differentiation to major events in the Gulf of Mexico and in the Atlantic Ocean.

The paleogeographic maps presented herein for the stages of the Jurassic are intended primarily to show the main areas of marine deposition and the provenance of the sediments; at the same time it is recognized that the areas probably are not in their true palinspastic position. Considerable evidence has been collected in recent years for large-scale transcurrent fault movements of regional proportions. Many of the Jurassic localities in southern Alaska certainly were involved in major dislocations bordering the rim of the north Pacific Ocean (Brew and others, 1966; Grantz, 1966; Richter and Jones, 1970; and St. Amand, 1957). Some dislocation of Jurassic strata occurred also in northern Alaska (Brosgé and TAILLEUR, 1970; TAILLEUR and Brosgé, 1970), but the dislocations were not as severe as in southern Alaska. The authors hope that locality and stratigraphic data published herein will stimulate field geologists to obtain more data on individual basins so that more accurate paleogeographic maps can be drawn.

Most of the geographic features, areas, and towns mentioned herein are shown on a physiographic map of Alaska prepared by Wahrhaftig (1965, pl. 1). Many of the creeks mentioned under locality descriptions dealing with northern Alaska are shown on an index map prepared by Imlay (1955, p. 79). Other streams mentioned under locality descriptions are shown on maps published by the authors listed in the locality descriptions at the end of this report.

ACKNOWLEDGMENTS

For the data published herein, the writers are greatly indebted to many geologists of the major oil companies and the U.S. Geological Survey and to a few individuals such as prospectors and mountain climbers. Especial thanks for their aid in reading the manuscript, in checking or rewriting the fossil locality data, or for contributing valuable collections are given to consulting geologist M. D. Mangus, to George Gryc, Arthur Grantz, J. M. Hoare, D. L. Jones, A. S. Keller, E. M. MacKevett, G. W. Moore, W. W. Patton, Jr., H. N. Reiser, Donald Richter, and E. G. Sable of the Geological Survey, and N. J. Silberling of Stanford University. The major oil companies whose geologists in recent years have furnished valuable fossil collections and locality data include the BP Alaska Inc., the Standard Oil Company of California, and the Atlantic Richfield Oil Company.

DISTRIBUTION OF JURASSIC ROCKS

Marine Jurassic fossils have been found in Alaska mostly in its northern, southern, and southwestern parts (fig. 1). The overall distribution of the fossils by stages is shown in figures 2-9. Most of the localities indicated therein represent single fossil collections, but some represent two or more collections from a sequence of beds in a small area, or many collections from an area of appreciable size, such as

the Iniskin Peninsula or the Talkeetna Mountains (fig. 10; also see Wahrhaftig, 1965, pl. 1).

Marine Jurassic fossils have not been found in a large area in central Alaska, nor in a southwestern prolongation (fig. 1) that trends southwest near and parallel to the Alaska Range. Most of the boundaries of this area are approximations that probably will be altered by future geologic mapping. The southern boundary, however, from the Talkeetna Mountains westward and southwestward to the Alaska Peninsula is well defined by intermittent occurrences of granitic rocks that were intruded during Early Jurassic time and that shed much coarse debris southward during Middle and Late Jurassic time (Grantz and others, 1963; Detterman and others, 1965; Detterman and Hartsock, 1966, p. 64, 70, 71; Burk, 1965, p. 73, 75-77).

The apparent absence of any marine Jurassic throughout the southern part of central Alaska north of Cook Inlet may be ascribed to erosion during Middle and Late Jurassic time. The lack of Jurassic fossils throughout the rest of central Alaska may have a similar explanation, but may be due to nondeposition. Their absence cannot be explained by post-Jurassic erosion, because at many places, marine Triassic rocks are overlain directly by lowermost Cretaceous marine rocks. Overall, the evidence favors the presence of a landmass in central Alaska during Jurassic time.




FIGURE 1.—Distribution of Jurassic fossils of all stages in Alaska.




FIGURE 2.—Distribution of Hettangian and Sinemurian fossils and inferred seas in Alaska. Land areas ruled. X, Hettangian and Sinemurian; H, Hettangian only; S, Sinemurian only. Numbers keyed to descriptions of Hettangian and Sinemurian fossil occurrences.




FIGURE 3.—Distribution of Pliensbachian and Toarcian fossils and inferred seas in Alaska. Land areas ruled. X, Pliensbachian and Toarcian; P, Pliensbachian only; T, Toarcian only. Numbers keyed to descriptions of Pliensbachian and Toarcian fossil occurrences.

JURASSIC PALEOBIOGEOGRAPHY OF ALASKA




FIGURE 4.—Distribution of Bajocian fossils and inferred seas in Alaska. Land areas ruled. Numbers keyed to description of Bajocian fossil occurrences.




FIGURE 5.—Distribution of Bathonian and Callovian fossils and inferred seas in Alaska. Land areas ruled. X, Bathonian and Callovian; B, Bathonian only; C, Callovian only. Numbers keyed to descriptions of Bathonian and Callovian fossil occurrences.




FIGURE 6.—Distribution of early Oxfordian fossils and inferred seas in Alaska. Land areas ruled. Numbers keyed to descriptions of early Oxfordian fossil occurrences.




FIGURE 7.—Distribution of late Oxfordian to early Kimmeridgian fossils and seas in Alaska. Land areas ruled. Numbers keyed to descriptions of late Oxfordian to early Kimmeridgian fossil occurrences.

JURASSIC PALEOBIOGEOGRAPHY OF ALASKA




FIGURE 8.—Distribution of late Kimmeridgian to early middle Tithonian fossils and inferred seas in Alaska. Land areas ruled. Numbers keyed to descriptions of late Kimmeridgian to early middle Tithonian fossil occurrences.




FIGURE 9.—Distribution of late middle to late Tithonian fossils and inferred seas in Alaska. Land areas ruled. Numbers keyed to descriptions of late middle to late Tithonian fossil occurrences.




FIGURE 10.—Index map of Jurassic areas in Alaska. Numbers keyed to figure 11.

COMPARISONS OF STRATIGRAPHIC AND LITHOLOGIC FEATURES

Jurassic rocks in Alaska have been studied in greatest detail in the southern and northern parts of the State. Consequently, critical stratigraphic and fossil control is available for more areas within those parts than elsewhere, as is reflected in the correlation charts (fig. 11). Igneous rocks that are radiometrically dated (Detterman and others, 1965; Grantz and others, 1963, 1966; Reed and Lanphere, 1969; Reiser and others, 1965) are present in some of the same areas and confirm ages established by paleontology.

Stratigraphic analysis of Jurassic rocks in some areas of Alaska indicate that strata now in close proximity were originally deposited under different conditions and from different source areas. These rocks are generally near known faults of demonstrably large displacement, and the original site of deposition can be inferred with some degree of accuracy. The movement of these rocks to their present localities may result from plate tectonics initiated by continental drift, as suggested by many authors (Carey, 1958; Hamilton, 1968; TAILLEUR and Brosgé, 1970; Richter and Jones, 1970; Rickwood, 1970). This report is not primarily concerned with plate tectonics, but some palinspatic interpretations are pertinent to the discussion of the stratigraphy.

SOUTHERN ALASKA

In southern Alaska, Jurassic strata are exposed in a nearly continuous arcuate belt from the extreme southeastern part near Ketchikan to near the tip of the Alaska Peninsula (figs. 10 and 11A). Lower Jurassic rocks, however, are known only from the central part of this belt between Juneau on the east and Wide Bay on the west (Burk, 1965; Detterman and Hartsock, 1966; Detterman and Reed, 1964; Grantz, 1960a,b, 1961a,b; Imlay, 1952; Juhle, 1955; Kirschner and Minard, 1949; MacKevett, 1963, 1965a,b, 1969, 1970a,b; Martin, 1905, 1926; Martin and Katz, 1912; Moffitt, 1922a,b, 1927; Paige and Knopf, 1907; Smith, 1917, 1939; Stanton and Martin, 1905).

During Early Jurassic time, deposition occurred within a nearly continuous belt, at least part of which bordered a volcanic island-arc system. Volcanic and volcanoclastic rocks form most of the record from the Katmai area in the south (Mather, 1925; Keller and Reiser, 1959) to the Talkeetna Mountains in the north (Grantz, 1960a,b, 1961a,b), but the main eruptive centers were in the Cook Inlet area (Detterman and Hartsock, 1966). At the same

time, argillite, limestone, chert, spiculite, and shale were deposited both as beds within the volcanic rocks, and at a considerable distance from the island-arc system in areas now represented by the Kuskokwim region (Cady and others, 1955; Hoare, 1961; Hoare and Coonrad, 1961a,b) and the Wrangell Mountains (MacKevett, 1963, 1965a,b, 1969, 1970a,b). Near the end of the subperiod, the rocks of southern Alaska were highly deformed and intruded by granitic batholiths in the Talkeetna Mountains (Grantz and others, 1963) and in the Cook Inlet area (Detterman and others, 1965; Reed and Lanphere, 1969). These granitic rocks are dated at 155 to 175 m.y. The oldest date probably represents the beginning of plutonism and may correspond to an unconformity dated paleontologically as late Toarcian to early Bajocian. Small mafic intrusive bodies are associated with the Lower Jurassic volcanic rocks in the Cook Inlet area and predate the granitic plutons.

Middle Jurassic rocks in southern Alaska are mainly graywacke-type sandstone, polymictic conglomerate, siltstone, and shale suggestive of rapid deposition in a subsiding epi-epieugeosynclinal trough. Volcanic rocks are a major part of the sequence only in the Kuskokwim region (Cady and others, 1955; Hoare, 1961). In general, the characteristics of the rocks in southern Alaska indicate a considerable change in depositional environment between Early and Middle Jurassic time. This change probably reflects orogenic activity that resulted in plutonism near the boundary of the subperiods. Similar orogenic activity continued throughout Middle Jurassic time, as indicated by the shifting positions of the seas and the uplift and erosion of the bordering uplands to the west and northwest. The many polymictic conglomerates in the Middle Jurassic sequence that contain clasts of both the granitic plutons and the Lower Jurassic volcanic rocks, which were mainly of marine origin, are evidence of these events. However, the presence of laumontite in the rocks of the Cook Inlet area suggests derivation from active volcanoes in the Kuskokwim region and indicates, therefore, that the relative geographic position of the two regions has not greatly changed since Middle Jurassic time.

During most of Late Jurassic time, marine deposition in southern Alaska took place in two or more basins that fragmented from an earlier larger Jurassic basin after a period of general emergence at about the transition from Callovian to Oxfordian time. One basin occupied the site of the Alaska Peninsula-Cook Inlet region and continued as far

east as the Wrangell Mountains; it was filled mainly with arkosic sediments derived from nearby granitic batholiths to the west and north (Atwood, 1911; Burk, 1965; Capps, 1923, 1934; Detterman and Hartsock, 1966; Grantz and others, 1966; Keller and Reiser, 1959; Kellum, 1945; Kirschner and Minard, 1949; Kirschner and Lyon, 1973; MacKevett, 1963, 1965a,b, 1969, 1970a,b, 1971; Martin, 1905, 1921, 1925, 1926; Martin and Katz, 1912; Mather, 1925; Miller and others, 1959; Moffit, 1922a,b, 1927, 1938a, 1943, 1954; Paige, 1906; Smith, 1925; Smith and Baker, 1924; Spurr, 1900; Stanton and Martin, 1905; Stone, 1905). Some silt and clay were deposited as beds within the arkosic sediments throughout most of the basin and locally formed a considerable part of the section. Deposition was continuous to the end of the Jurassic only at the extreme southern end of the Alaska Peninsula; elsewhere, the uppermost Jurassic beds were removed or deposition did not occur.

Another basin, or basins, existed in easternmost Alaska and probably extended into adjoining British Columbia, as shown by the presence of thousands of feet of Upper Jurassic graywacke, mudstone, conglomerate, argillite, and slate now exposed in southeastern Alaska (Brew and others, 1966; Lathram and others, 1960; Loney, 1964; Martin, 1926) and in the eastern Alaska Range (Richter, 1971). The characteristics of these rocks contrast so much with those of the arkosic rocks deposited at the same time in the Cook Inlet region that the provenance of deposition and the lithology of the source areas must have been quite different. The differences could be explained by extensive crustal movements, as postulated by Richter and Jones (1970). These crustal movements have apparently foreshortened the distance between the basins by many tens of miles along major faults in southeastern Alaska.

NORTHERN ALASKA

Jurassic rocks in northern Alaska have been studied in detail only in the eastern part (figs. 10 and 11B). Characteristics of the strata indicate that the provenance of deposition and lithology of the source areas were considerably different than in southern Alaska during most of the Jurassic (Brosge and Tailleux, 1970; Detterman, 1973; Gryc and Lathram, 1954; Gryc and Mangus, 1954; Keller and others, 1961; Leffingwell, 1919; Reed, 1968; Reiser and Tailleux, 1969; Reiser and others, 1970; Sable, 1965; Smith and Mertie, 1930). However, marine connections with southern Alaska probably existed via Yukon Territory and east-central Alaska (Frebold and others, 1967; Jeletzky, 1962). The

stratigraphic sequence in the western part of northern Alaska is complicated by thrust faulting, which has juxtaposed two dissimilar Jurassic sections; one is typical of northern Alaska sedimentary rocks, and the other is more closely related to Jurassic volcanic rocks south of the Brooks Range (Brosge and Tailleux, 1970; Detterman, 1972; Gates and Gryc, 1963; Jones and Grantz, 1964; Lathram, 1965; Patton and Grantz, 1962; Tailleux, 1969).

Lower Jurassic rocks in northern Alaska are mainly fissile clay shale and claystone that commonly contain limestone concretions. Siltstone and silty shale are abundant locally. Chert, tuff, graywacke, and oil shale are present at a few localities in the western part (Brosge and Tailleux, 1970; Detterman, 1973; Imlay, 1955, 1967; Mangus and others, 1954; Tailleux, 1964; Tailleux and Kent, 1954; Tailleux and others, 1966). A glauconitic sandstone is present in the subsurface along the north coast (Bergquist, 1966; Collins, 1958, 1961; Detterman, 1973; Gates and others, 1968; Imlay, 1955; Loeblich, 1954; Payne, 1955; Payne and others, 1951; Robinson, 1959), and a massive sandstone of earliest Jurassic or latest Triassic age is present along the north flank of the Brooks Range in northeastern Alaska (Detterman, 1973; Sable, 1965). The sandstones would indicate that sediment sources were both north and south. The Lower Jurassic section is not complete at any locality and is only sparsely fossiliferous; consequently, regional unconformities are impossible to determine. There are probably many local unconformities, and paleontologic evidence suggests that parts of the Hettangian, Sinemurian, and Pliensbachian are missing.

Siltstone, silty shale, and pyritic clay shale are the main rock types present in the Middle Jurassic of northern Alaska. Orange-weathering clay ironstone concretions and beds are also common. Graywacke and tuff are present locally in the central part (Chapman and others, 1964; Jones and Grantz, 1964; Patton and Keller, 1954; Patton and Tailleux, 1954, 1964). The only good exposures for this part of the Jurassic are in northeastern Alaska (Brosge and Tailleux, 1970; Detterman, 1973; Gryc, 1951; Gryc and Mangus, 1954; Keller and others, 1961; Leffingwell, 1919; Reiser and Tailleux, 1969; Reiser and others, 1970; Sable, 1965; Whittington and Sable, 1954), where a major middle Bajocian to early Bathonian unconformity can be detected both lithologically and paleontologically (Imlay, 1955, 1970). Elsewhere in northern Alaska, the rocks are too poorly exposed to determine the presence of unconformities.

JURASSIC PALEOBIOGEOGRAPHY OF ALASKA





FIGURE 11.—A, Correlation of Jurassic rocks in southern Alaska. Vertical lines indicate strata missing; right diagonal lines indicate gradational or indefinite contact; jagged lines indicate gradational or indefinite contact. Numbered columns keyed



indicate beds not exposed; left diagonal lines indicate lack of fossil data; wavy lines indicate unconformity or disconformity to figure 10. Compiled October 1971. (Figure 11 continued on following pages.)





FIGURE 11.—Continued. B, Correlation of Jurassic rocks in northern and east-central Alaska. Vertical lines indicate strata unconformity or disconformity; jagged lines indicate gradational or



missing; right diagonal lines indicate beds not exposed; left diagonal lines indicate lack of fossil data; wavy lines indicate indefinite contact. Numbered columns keyed to figure 10. Compiled October 1971.

Upper Jurassic rocks in northern Alaska consist mainly of siltstone and shale similar to those of the Middle Jurassic. The principal exceptions are a thick sequence of sandstone, siltstone, and mudstone turbidites exposed in the far western part of northern Alaska near Cape Lisburne (Campbell, 1967; Chapman and Sable, 1960; Detterman, 1972; Knowlton, 1914; Sable and Mangus, 1954; Tailleur, 1969) and a thin sequence of tuffaceous graywacke, conglomerate, and chert exposed in the central part of northern Alaska (Brosgé and Tailleur, 1970; Chapman and others, 1964; Patton, 1956; Patton and Tailleur, 1964). These rocks that are exceptions may have been faulted into their present positions from somewhere south of the modern Brooks Range (Brosgé and Tailleur, 1970; Tailleur, 1969).

Most of northern Alaska was apparently emergent during late Callovian and early Oxfordian time and again during the late Tithonian. These periods of emergence are marked by major unconformities and by a change in the faunal assemblages. No ammonites have been found in rocks of Tithonian Age, but the pelecypod *Buchia* is present in many areas (Brosgé and Tailleur, 1970; Detterman, 1973; Im-lay, 1959; Patton, 1956).

A sequence of graywacke, siltstone, shale, chert, and conglomerate in the central part of northern Alaska was formerly mapped as the Tiglukpuk Formation of Late Jurassic age (Patton, 1956). The type locality on Tiglukpuk Creek is now known to be a faulted section consisting of Middle Jurassic and Lower Cretaceous rocks (Jones and Grantz, 1964); consequently, the formation name is herein abandoned.

Jurassic strata are present in east-central Alaska (figs. 10 and 11B) but have not been studied in detail. The rocks are poorly exposed, and the stratigraphic succession has not been completely determined. Two facies are present, one of shale, siltstone, and argillite in the Kandik and Nation Rivers area (Brabb, 1969, 1970; Brabb and Churkin, 1969), and the other of sandstone, quartzite, and shale in the Porcupine, Coleen, and Sheenjek Rivers area (Brosgé and Reiser, 1964, 1969; Maddren, 1912; Maddren and Harrington, 1914; Mertie, 1929, 1930). Sandstone and quartzite of the Porcupine River area contain a Hettangian fauna and probably represent the initial sediments in a southward-expanding northern Alaska-Yukon Territory sea (Jeletzky, 1962). The sea continued to expand and is believed to have connected northern and southern Alaska through Middle and early Late Jurassic time (figs. 2-9). Volcanic rocks dated at 155 to 165 m.y. (Reiser

and others, 1965) are present in the Christian area northwest of Porcupine and may indicate tectonic activity associated with uplift and subsequent sub-aerial erosion during the later part of Late Jurassic time.

OCCURRENCES OF JURASSIC UNCONFORMITIES

SOUTHERN ALASKA

The presence of unconformities in southern Alaska has been fairly well established (fig. 12). The lowest known unconformity is at the top of Lower Jurassic volcanic rocks north of Cook Inlet (Detterman and Hartsock, 1966, p. 69, 71; Grantz and others, 1963, p. B58); it represents a time of pronounced folding, and, on the basis of ammonites from the Talkeetna Mountains, must have formed during the latest Toarcian and the earliest Bajocian time (*Leioceras opalinum* zone). In the Wrangell Mountains, an unconformity possibly formed at the same time occurs between the Nizina Mountain and Lubbe Creek Formations (MacKevett, 1969, p. A41, A44).

The next higher well-recognized unconformity in southern Alaska formed during the later part of late Bajocian time. It is most clearly shown on the Iniskin Peninsula (Detterman and Hartsock, 1966, p. 35, 40) where the Bowser Formation of Bathonian Age cuts across 420 feet of the Twist Creek Siltstone of early late Bajocian Age onto beds of late middle Bajocian Age. The same unconformity is probably represented also on the Alaskan Peninsula in the lower part of a hiatus of middle Bajocian to late Bathonian Age. The fact that the youngest Bajocian beds exposed at Wide Bay are as old as early middle Bajocian (*Otoites sauzei* zone) is probably a result of differential erosion such as occurred on the Iniskin Peninsula. Such a possibility is supported by the presence on Alinchak Bay of *Inoceramus ambiguus* Eichwald (Mesozoic loc. 12390), which in Alaska is not known in beds older than the late middle Bajocian (*Stephanoceras humphriesianum* zone).

A third unconformity above the base of the Jurassic in the Cook Inlet region formed late in the Bathonian. It is marked between Chinitna Bay and Tuxedni Bay by northward truncation of the uppermost part of the Bowser Formation. It is marked in the Talkeetna Mountains by erosion that locally removed all beds equivalent to the Bowser Formation. It may be represented in the Wrangell Mountains in the basal part of a hiatus of probable late Bathonian to early Oxfordian Age. It is probably represented in the Alaska Peninsula in the upper

part of a hiatus between beds of early middle Bajocian and early Callovian Age. On Wide Bay and Puale Bay, lower Bathonian sediments were either never deposited or were eroded away during late Bathonian time. If the latter possibility is true, then some Bathonian beds may still be present in the subsurface of the Alaska Peninsula.

A fourth unconformity in southern Alaska formed during the late Callovian and in some areas persisted into the early Oxfordian. It represents this entire time span throughout most of the Alaska Peninsula except near Mount Peulik (Mesozoic loc. 10798) west of Puale Bay. It represents only the late Callovian in the Cook Inlet region where middle Callovian beds in the upper part of the Chinitna Formation are overlain unconformably by the Nak-

nek Formation that basally contains the early Oxfordian ammonite *Cardioceras* (Detterman and Hartsock, 1966, p. 47-49). Farther east in the Wrangell Mountains, the unconformity represents the upper part of a hiatus that apparently extends from late Bathonian to early Oxfordian time.

Another unconformity of latest Jurassic age may extend across most of southern Alaska east of Wide Bay as indicated by an absence of such late Tithonian mollusks as *Buchia piochii* (Gabb) or *B. fischeriana* (d'Orbigny) and by the presence locally of beds of Valanginian to Hauterivian Age resting on beds of late Kimmeridgian to middle Tithonian Age. For example, in the Talkeetna Mountains, the uppermost Jurassic beds containing *B. rugosa*




FIGURE 12.—Occurrences of unconformities in the Jurassic of Alaska. Blank spaces indicate strata of known ages; vertical lines indicate strata missing; right diagonal lines indicate beds not exposed; left diagonal lines indicate lack of fossil data.

(Fischer) and *B. mosquensis* (von Buch) are overlain by conglomerates and sandstones containing Buchias of early to middle Valanginian Age (Imlay and Reeside, 1954, p. 233; Grantz and others, 1966, p. C46). Evidently the unconformity below the Valanginian beds could have formed during the late Tithonian, or the Berriasian, or the entire time interval.

NORTHERN ALASKA

In northern Alaska, only the area east of the Lupine River has been studied in sufficient detail for the presence and positions of unconformities to be fairly well documented. In that area, one unconformity has been dated as late middle Bajocian to possibly early Bathonian because the youngest known Bajocian beds contain *Arkelloceras*, and the oldest Bathonian beds contain *Cranocephalites*. A slightly higher unconformity may exist at the Bathonian-Callovian boundary (see columns 15 and 16, fig. 11B), although the westward pinchout of the Bathonian beds between the Canning and Sagavanirktok Rivers could be depositional against a landmass in the area of the Killik and Itkillik Rivers. The next higher unconformity has been dated as late Callovian because the underlying beds contain *Cadoceras* and *Pseudocadoceras*?, and the overlying beds contain *Cardioceras*.

The absence of any ammonites of late Callovian Age deserves special mention because beds of that age are also absent in Canada (Frebold, 1961, table 1 opposite p. 26; 1964a, p. 4; Frebold and Tipper, 1967, p. 18) and have been found in North America only in east-central California (Imlay, 1961b, p. D6, D27) and in the states of Oaxaca and Guerrero in southern Mexico (Burckhardt, 1927; 1930, p. 26, 32-36, 43; Imlay, 1961b, p. D12). Evidently marine waters covered very little of North America during late Callovian time.

The existence of other unconformities in northern Alaska has not as yet been proven. The fact that the latest Tithonian has been identified at only two places (Mesozoic locs. 26390, M2531) suggests however, that marine deposition was not widespread at that time. Also, nondeposition of Upper Jurassic beds throughout a large area in north-central Alaska is indicated by the fact that to date only Early Cretaceous species of *Buchia* have been collected in the foothills of the Brooks Range between the Killik and Itkillik Rivers (Jones and Grantz, 1964), and those species were obtained near beds of Late Triassic age. Similarly, the reported Late Jurassic age of certain beds in the subsurface of north-central Alaska is questioned because their age determina-

tion was based on species of Foraminifera that were originally obtained from the beds just mentioned (Tappan, 1955, p. 25).

The sequence of events in northern Alaska during the Jurassic both resembled and differed from that in southern Alaska. Resemblances include moderately widespread seas from Sinemurian to early middle Bajocian time and from Oxfordian to middle Tithonian time. Resemblances also include formation of unconformities in late Callovian and late Tithonian time. Differences from southern Alaska include the absence of an unconformity in northern Alaska at the end of Toarcian time and the formation of an unconformity at a different time during Bajocian and Bathonian time. For example, in northern Alaska no ammonites have been found that represent the late middle Bajocian (*Stephanoceras humphriesianum* zone), the late Bajocian, and the early Bathonian. In southern Alaska, by contrast, that time is well represented, except for the latest late Bajocian. If failure to find ammonites of certain ages in northern Alaska is not due to inadequate collecting, then the advances and retreats of the seas in general occurred at different times in northern than in southern Alaska and reflect local diastrophic movements. These differences are probably related to times of intrusion and uplift of granitic rocks of the Aleutian Range batholith in southern Alaska (Detterman and Hartsock, 1966, p. 63, 64, 69-71).

EXTENT OF JURASSIC SEAS

In the Early Jurassic, the fossil record indicates that marine waters spread most widely across southern Alaska during Sinemurian time and most widely across northern Alaska during Pliensbachian and Toarcian time (figs. 2, 3, and 13). The Sinemurian in northern Alaska may be coextensive, however, with the geographic range of *Otapiria tailleuri* Imlay (1967, p. B3-B7, pl. 1, figs. 1-23, pl. 2, fig. 32), which species occurs in a distinctive, rather thin organic shale and has been found over a distance of at least 200 miles north of the Brooks Range between the Etivluk and Itkillik Rivers. Evidence for such an extension of the Sinemurian consists (1) of a large well-preserved float specimen of *Crucilobicer* (Mesozoic loc. 29774 from the same place as localities M 2451 and 29280 in Imlay, 1967, p. B7); (2) four small ammonite molds identical with the inner whorls of *Crucilobicer* (Mesozoic loc. 29775 from the same place as locality 24060 in Imlay, 1967, p. B7); (3) similar small ammonite molds (Mesozoic locs. 29281 and 29282) that previously were compared with *Tmetoceras* and other genera

(Imlay, 1967, p. B6, B7); and (4) the association of the ammonites with *Otapiria tailleuri* Imlay at USGS Mesozoic localities 29282 and 29775. The inferences from this data are that the Sinemurian sea was probably nearly as extensive in northern Alaska as the Pliensbachian and Toarcian seas, and in Alaska as a whole the Sinemurian sea was nearly as extensive as the Bajocian sea.

During Bajocian time, marine waters transgressed widely in southern, southwestern, north-central and northeastern Alaska and were connected eastward with seas in Canada. Near Wide Bay on the Alaska Peninsula, deposition continued at least until the early middle Bajocian (*Parabigotites crassicosatus* beds), in the Cook Inlet region until the early late Bajocian (*Megasphaeroceras rotundum* beds), and in northeastern Alaska until the early middle Bajocian (*Arkelloceras* beds) (fig. 14). The duration of the sea in the Kuskokwim region of southwestern

Alaska is not known because the fossil evidence consists almost entirely of *Inoceramus ambiguus* Eichwald, which in the Cook Inlet regions range from late middle Bajocian (*Teloceras itinsae* beds) through the Bathonian and occurs rarely in beds of early Callovian Age. Southeastern Alaska was probably also covered by a sea during at least middle Bajocian time, as indicated by occurrences of fossils of that age in nearby northwestern British Columbia (Frebold and Tipper, 1970, fig. 6 on p. 10). Whether the Bajocian sea ever extended across northwestern Alaska west of the Okpikruak River must await more fieldwork.

During Bathonian time, marginal seas definitely occurred in northeastern Alaska east of the Sagavanirktok River and in southern Alaska between the Iniskin Peninsula and the Wrangell Mountains (fig. 5). These clearly were westward extensions of seas of that age in central and northern Yukon Terri-

Stages	Western Nevada (Muller and Ferguson, 1939; Hallam, 1965)	Southern and eastern Mexico (Erben, 1956, 1957)	East-central Oregon (Imlay, 1968)	Southern Alaska (Imlay, 1968)	Northern Alaska (Imlay, 1955)
Toarcian	<i>Grammoceras</i> and <i>Pseudolioceras</i>	Lower part of coal- and plant-bearing beds	<i>Catullocheras</i> and <i>Dumortieria</i>		<i>Pseudolioceras</i>
	<i>Catacoeloceras</i> and <i>Dactyloceras</i>		<i>Haugia</i> and <i>Polyplectes</i>	<i>Grammoceras</i>	
	?		?	<i>Haugia</i> and <i>Phymatoceras</i>	<i>Dactyloceras</i> cf. <i>D. commune</i> <i>Dactyloceras</i> cf. <i>D. semicelatum</i> and <i>Nodicoeloceras</i>
	<i>Nodicoeloceras</i>			<i>Dactyloceras</i> cf. <i>D. commune</i>	
Pliensbachian	<i>Arietoceras</i>	<i>Arietoceras</i>	<i>Arietoceras</i> , <i>Reynesoceras</i> , <i>Paltarpites</i> , <i>Prodactyloceras</i> , and <i>Leptaleoceras</i>	<i>Pleuroceras</i> , <i>Amaltheus</i> , and <i>Paltarpites</i>	<i>Amaltheus</i>
		?	?		?
	?	?		?	
	<i>Uptonia</i> ?	<i>Uptonia</i>		<i>Uptonia</i> , <i>Apodoceras</i> , and <i>Acanthopleuroceras</i>	
Sinemurian	<i>Eoderoceras</i> and <i>Cruciloboceras</i>	<i>Microderoceras</i> and <i>Echioceras</i>	<i>Cruciloboceras</i>	<i>Cruciloboceras</i>	<i>Cruciloboceras</i>
	<i>Oxynoticeras</i>	<i>Pleurechioceras</i>			
	?	?			
		<i>Vermiceras</i>			
		<i>Oxynoticeras</i>			
		<i>Euagastoceras</i>			
Hettangian	?	?		<i>Microderoceras</i>	
	<i>Arnioceras</i> , <i>Arietites</i> , <i>Coroniceras</i> , and <i>Megarietites</i>	<i>Arnioceras</i>	<i>Arnioceras</i>	<i>Arnioceras</i>	<i>Arietites</i>
		?	?	?	
		<i>Coroniceras</i>		<i>Coroniceras</i>	
Hettangian	<i>Schlotheimia</i> and <i>Alsatites</i>	Not exposed	<i>Schlotheimia</i> and <i>Alsatites</i>		
	<i>Waehneroceras</i> and <i>Psiloceras</i>		<i>Waehneroceras</i> and <i>Psiloceras</i>	<i>Waehneroceras</i>	<i>Psiloceras</i>

FIGURE 13.—Correlation of Hettangian to Toarcian ammonite faunas in Alaska.

tory, Canada (Frebold and others, 1967, p. 11, 12, 23). The fossil record (fig. 14) indicates that the sea in southern Alaska existed during the entire Bathonian, and the sea in northeastern Alaska existed only during the middle to late Bathonian. Marine waters probably did not extend as far southwest as Puale Bay and Wide Bay in the Alaska Peninsula, because beds of early Callovian Age rest directly on beds of middle Bajocian Age in both places. Whether or not the Bathonian sea ever extended westward across north-central and north-western Alaska must await detailed mapping and fossil collecting comparable with that now nearing completion in northeastern Alaska. Failure to find any Bathonian fossils in southeastern Alaska coin-

cides with an absence of such fossils in western British Columbia (Frebold and Tipper, 1967, p. 4; 1970, p. 12, 16, 17). The Bathonian may be represented in the Kuskokwim region by some occurrences of *Inoceramus ambiguus* Eichwald.

The seas of Callovian time in Alaska were continuations and extensions of the Bathonian seas (fig. 5). They extended at least as far southwest as Wide Bay on the Alaska Peninsula, which is 200 miles beyond the southernmost Bathonian occurrence in the Cook Inlet region. Failure to find any megafossils of Callovian Age in north-central and northwestern Alaska may have some significance paleogeographically but could be due to inadequate collecting. Failure to find any Callovian megafos-

Stages	Southern Mexico Guerrero and Oaxaco (Burckhardt, 1927, 1930; Arkell, 1956; Imlay, 1952; Erben, 1956, 1957)	Eastern Oregon and California (Imlay, 1961b, 1964a, in part)	Southern Alaska (Imlay, 1953b, 1962a, b, 1964b; Westermann, 1964, 1969)	Northern Alaska (Imlay, 1955)
Callovian	<i>Peltoceras</i> spp.	<i>Peltoceras</i> (<i>Metapeltoceras</i>)? (eastern California only)	No fossil evidence	No fossil evidence
	?	?	?	?
	<i>Erymnoceras mixtecorum</i>	<i>Lilloettia</i> cf. <i>L. stantoni</i> and <i>Pseudocadoceras growingki</i>	<i>Cadoceras</i> (<i>Stenocadoceras</i>) <i>stenoloboide</i>	<i>Cadoceras</i> , and <i>Pseudocadoceras</i> <i>growingki</i>
	<i>Reineckella neogaea</i>	<i>Lilloettia buckmani</i> , <i>Xenocephalites vicarius</i> , and <i>Keplerites</i>	<i>Lilloettia buckmani</i> , <i>Cadoceras catostoma</i> , and <i>C. (Paracadoceras) tonniense</i>	?
Bathonian	<i>Epiptenoceras paracontrarium</i>	?	Disconformity locally	Not known
	?	Probably represented by <i>Keplerites</i> , <i>Cobbanites</i> , <i>Parareineckella</i> , <i>Xenocephalites</i> , and <i>Choffatia</i> in upper part of Snowshoe Formation	?	?
	No fossil evidence		<i>Arctocephalites</i> cf. <i>A. elegans</i>	<i>Arctocephalites elegans</i>
	?		<i>Cranocephalites</i>	<i>Cranocephalites</i>
Bajocian	<i>Zigzagoceras</i>	?	<i>Cranocephalites costidensis</i>	?
			?	
	upper	No fossil evidence	Angular unconformity on Iniskin Peninsula	No fossil evidence
	?	?	?	
		<i>Spiroceras</i> , <i>Megasphaeroceras</i> , <i>Leptosphinctes</i> , and <i>Normannites</i>	<i>Megasphaeroceras rotundum</i> , <i>Normannites</i> , <i>Leptosphinctes</i> , and <i>Sphaeroceras</i>	
	" <i>Stephanoceras</i> " and <i>Normannites</i>	<i>Chondroceras allani</i> , <i>Normannites</i> <i>crickmayi</i> , and <i>Teloceras itinsae</i>	<i>Chondroceras allani</i> , <i>Normannites</i> <i>crickmayi</i> , and <i>Teloceras itinsae</i>	
	?	<i>Dorsetensia</i>	?	
		<i>Parabigotites crassicostratus</i>	<i>Papilliceras</i> , <i>Witchellia</i> , <i>Otoites</i> , <i>Emileia</i> , <i>Dorsetensia</i> , and <i>Bradfordia</i>	
		<i>Papilliceras stantoni</i> and <i>Witchellia</i>	<i>Normannites</i>	<i>Arkelloceras</i>
	No fossil evidence		<i>Parabigotites crassicostratus</i> , <i>Dorsetensia adnata</i> , and <i>Stephanoceras juhlei</i>	?
middle		?	<i>Witchellia sutneroides</i>	
		<i>Sonninia</i> (<i>Euhoplaceras</i>), <i>Fontannesia</i> , <i>Dicidoceras</i> , <i>Hebetoxyites</i> , <i>Praestrigites</i> , and <i>Strigoceras</i>	<i>Sonninia</i> (<i>Euhoplaceras</i>) and <i>S. (Alaskoceras)</i> <i>Dicidoceras? paucinodosum</i>	
		<i>Tmetoceras scissum</i> , <i>Praestrigites</i> , and <i>Eudometoceras</i>	<i>Erycitoides howelli</i> , <i>Erycites</i> , and <i>Tmetoceras</i>	
		<i>Tmetoceras scissum</i>	<i>Tmetoceras scissum</i>	
lower		?	?	
	Upper part of coal- and plant-bearing beds	No fossil evidence	No fossil evidence	<i>Pseudoloceras</i> <i>maclintocki</i>

FIGURE 14.—Correlation of Bajocian to Callovian ammonite faunas in Alaska.

sils in the Wrangell Mountains is probably related to a disconformity of pre-late Oxfordian Age at the base of the Root Glacier Formation (MacKevett, 1969, p. A46). The fossil record (fig. 14) indicates that the sea withdrew entirely from Alaska at the end of the middle Callovian.

Early in the Oxfordian, marine waters covered the same areas in Alaska as during the Callovian (compare figs. 5 and 6), but later in the epoch and during the early Kimmeridgian they spread widely across both northern and southern Alaska (fig. 7). Evidently in southern Alaska the sea spread over an uneven terrain south of mountains that shed much conglomerate and sandstone into the sea (Detter-

man and Hartsock, 1966, p. 49, 71). The uneven character of the terrain is shown by the fact that the basal beds of the Oxfordian sequence contain the early Oxfordian ammonite *Cardioceras* in four areas, whereas in the Wrangell Mountains and at most places on the Alaska Peninsula the basal beds contain the younger ammonite *Amoeboceras* and its associate *Buchia concentrica* (Sowerby) (figs. 11A and 15). Failure to find any Oxfordian or early Kimmeridgian fossils in the Kuskokwim region or in southeastern Alaska suggests that rocks of that age are missing in those areas, but more verification is needed before paleogeographical conclusions can be drawn.

Stages		Eastern and northern Mexico (Imlay, 1943, 1952; Erben, 1957)		California, Oregon, and western Idaho (Imlay, 1953a; 1961b; Imlay and Jones, 1970)		Southern Alaska (Imlay, 1953b, 1961a)		Northern Alaska (Imlay, 1955)			
Tithonian	upper	Substeuoceras and Proniceras		Microcanthoceras, Parodontoceras, and Hildglochiceras	Substeuoceras and Spiticeras		Buchia aff. B. okensis ?		Buchia piochii and B. fischeri	Buchia cf. B. fischeriana	
		?			Parodontoceras		?				
		Kossmatia, Durangites, and Corongoceras			?		Buchia piochii				
	?		Kossmatia								
	middle				?						?
lower	Pseudolissoceras and Subplanites		Virgatosphinctes and Aulacosphinctoides	Not identified		B. rugosa and B. mosquensis		B. rugosa and B. mosquensis			
	Mazapillites										
Kimmeridgian	upper	Hybonoticeras cf. H. beckeri		Amoeboceras (Amoelites) and Idoceras		Buchia concentrica		Amoeboceras (Prionodoceras) spiniferum and Dichotomosphinctes		Buchia concentrica	
		Idoceras durangense and Glochiceras fialar									
	lower	Idoceras balderum									
Sutneria cf. S. platynota											
Oxfordian	upper	Discosphinctes caribbeanus and Ochetoceras canaliculatum		Discosphinctes virgulatus and Dichosphinctes mulbachi		Buchia concentrica		Amoeboceras (Prionodoceras) spiniferum and Dichotomosphinctes		Amoeboceras (Prionodoceras)	
		Dichotomosphinctes durangensis									
	lower	Not identified		Cardioceras martini (Idaho only)		Cardioceras distans		Cardioceras (Scarburgiceras) martini			
		Creniceras and Parapaltoceras locally								Not identified	

FIGURE 15.—Correlation of Oxfordian to Tithonian molluscan faunas in Alaska.

From late Kimmeridgian until middle Tithonian time (fig. 8) marine waters spread even more widely over Alaska. They covered large parts of the Kuskokwim region and of southeastern Alaska, and, to judge by the fossil record, covered more of Alaska than at any other time during the Jurassic. The absence of Jurassic sediments younger than early Kimmeridgian in the area between Iniskin Bay and Tuxedni Bay on Cook Inlet is probably due to Late or post-Jurassic uplift and erosion (Detterman and Hartsock, 1966, p. 71) rather than to nondeposition.

During late Tithonian time, the marginal seas in Alaska became much reduced, compared with those of the immediately preceding time (compare figs. 8 and 9). One sea covered most of the present site of the Alaska Peninsula southwest of Wide Bay. Another sea existed in an area now occupied by the eastern part of the Alaska Range and apparently extended eastward into the Yukon Territory (Frebold and Tipper, 1970, p. 15; Frebold and others, 1967, p. 10). Failure to find any marine fossils of that age elsewhere in southern Alaska could be explained by erosion or by nondeposition, or both. In southeastern Alaska the sequence exposed on Admiralty Island (Loney, 1964, p. 59, 92, 97) apparently represents continuous deposition from Kimmeridgian until Valanginian time. In northern Alaska, the geographic position of a single specimen of *Buchia* cf. *B. fischeri* (d'Orbigny) indicates the presence of a fairly widespread sea that presumably extended eastward into the northern Yukon (Jeletzky, 1958, p. 5, 1960, p. 4; 1966, p. 25, 30; 1967, p. VI; Frebold and others, 1967, p. 10).

In summation, Jurassic marginal seas occupied considerable areas in northern, southern, and southwestern Alaska, but probably never covered a huge area in central Alaska or a southwestern prolongation of that area. From the presently known fossil occurrences (figs. 2-9), it is inferred that marine waters were absent during late Callovian time, much restricted during Hettangian, Bathonian, early Oxfordian and late Tithonian time, somewhat less restricted during Pliensbachian, Toarcian, and Callovian time, fairly widespread during Sinemurian, Bajocian, and late Oxfordian to early Kimmeridgian time, and most widespread in late Kimmeridgian to middle Tithonian time.

SUCCESSION AND DIFFERENTIATION OF AMMONITES

Most of the Early Jurassic ammonites of Alaska are identical with and occur in essentially the same succession as ammonites of that age in central and northern Europe and in Canada (fig. 13). In fact,

the Alaskan ammonites of late Pliensbachian Age have more in common with the ammonites of those areas than with late Pliensbachian ammonites of eastern Oregon, which have Mediterranean affinities (Imlay, 1968, p. C21). Such close resemblances eastward across Canada show that the seaways between Alaska and northwest Europe were open during Early Jurassic time. They suggest that continental drift either did not occur during that time or had no influence on the distribution of ammonites between those areas.

The succession of ammonites in Alaska during Bajocian time (fig. 14), as previously discussed (Imlay, 1965, p. 1030), was similar to that in Europe and in the Tethyan region except for the presence in Alaska of a few genera that have been found only in areas near the Pacific Ocean and the absence in Alaska of a few genera that are fairly common in Europe. These differences were slight during the early Bajocian but became more pronounced by late Bajocian. In southern Alaska, the succession ended early in late Bajocian time. In northern Alaska the succession ended in early middle Bajocian time.

The succession of ammonites in Alaska during Bathonian time (fig. 14) was essentially identical with that in northern Canada (Frebold, 1964b), in East Greenland (Callomon, 1959), and in other lands bordering the Arctic Ocean. It was dominated by the genera *Cranocephalites*, *Artocephalites*, and *Arcticoceras*. It had almost nothing in common generically with the Bathonian ammonite assemblages of central Europe and the Tethyan region (Imlay, 1965, p. 1024, 1030, 1031). In southern Alaska, however, the Bathonian ammonite succession includes genera, such as *Parareineckeia* and *Xenocephalites*, that are known only from the Western Hemisphere. Evidently the Bathonian ammonites of the Boreal region were almost completely separated from Bathonian ammonites farther south by barriers of some kind, except along the Pacific Coast south of Alaska. As a consequence, the ammonite faunas of the world became differentiated into distinct Boreal and Tethyan realms and a somewhat less distinct Pacific realm.

A Callovian ammonite succession has been established in Alaska only in the southern part between the Talkeetna Mountains and Wide Bay on the Alaska Peninsula (fig. 14). This succession is essentially the same as along the Pacific Coast in British Columbia (Frebold and Tipper, 1967, p. 5), Oregon (Imlay, 1964a, p. D7-D10), and California (Imlay, 1961b, p. D5). It is characterized by such taxa as

Cadoceras, *C. (Paracadoceras)*, *C. (Stenocadoceras)*, *Pseudocadoceras*, *Kepplerites*, *Lilloettia*, and *Xenoccephalites*; the first five are of Boreal origin, and the last two originated in seas of the Western Hemisphere. Except for *Lilloettia* and *Xenoccephalites*, the succession has very little in common generically with ammonites of that age in Mexico and even less in common with ammonites of the Mediterranean region, as discussed elsewhere (Imlay, 1953b, p. 56; 1965, p. 1023-1033). It bears no resemblances on the specific levels to ammonites in the western interior of the continent and in northern Canada (Imlay, 1953a; Frebold, 1961, p. 17-22; 1964a, p. 4, 1964b. The succession differs even more with ammonites of the same age in Greenland and northern Eurasia (see comparisons in Frebold, 1961). Overall, the Callovian ammonites of southern Alaska are dominantly of Boreal origin but are in part of local origin and in part of Pacific origin.

The succession in Alaska of ammonites of Oxfordian to early Kimmeridgian Age (Imlay, 1952, p. 977-978) (fig. 15) is nearly the same as that along the Pacific Coast from British Columbia to California (Frebold and Tipper, 1967, p. 5; Imlay, 1961b, p. D5, 1964a, D7-10). It is likewise identical with, or closely similar to, the succession in the western interior of the continent and in northern Canada (Frebold and others, 1959, 1967, p. 10; Frebold, 1964b, p. 481). It can be correlated with ammonite successions in East Greenland and northern Eurasia by means of *Cardioceras* and *Amoeboceras* but otherwise differs by having far fewer genera (see comparisons in Frebold, 1961, table 1). The succession has nothing in common with ammonites of that age in Mexico or in the Mediterranean region, except for the presence of *Dichotomosphinctes*.

An ammonite succession has not been established in the uppermost Jurassic beds of Alaska. A few specimens of *Lytoceras* and *Phylloceras* occur in beds of middle Kimmeridgian to middle Tithonian Age which are identified by the presence of *Buchia rugosa* (Fischer) and *B. mosquensis* (von Buch). The very latest Jurassic, characterized by *Buchia piochii* (Gabb), has not furnished any ammonites. This scarcity of ammonites in Alaska in beds younger than early Kimmeridgian is in marked contrast to a fair abundance of ammonites in equivalent beds in East Greenland and northern Eurasia. As a consequence of this scarcity, most correlations above the lower Oxfordian are made on the basis of species of *Buchia* whose succession during the Late Jurassic appears to be the same throughout the Arctic.

The succession of Jurassic ammonites in Alaska, in comparison with successions in other parts of the world, shows that differentiation into Boreal and Tethyan realms started in late Pliensbachian time, increased gradually during the Bajocian, was completed by the Bathonian, and persisted into the Early Cretaceous. Evidently faunal differentiation during Bathonian and later Jurassic time coincides very well with the deposition of Jurassic salt and some associated red beds in the Gulf of Mexico region (Imlay, 1943, p. 1438, 1508, 1509; Bishop, 1967, p. 249; Viniegra-O, 1971, p. 478-484; fig. 10 on p. 492) which apparently coincides in time with early stages of continental drift. Faunal differentiation probably resulted from a combination of events (Imlay, 1965) which may include the separation of the continents.

In summation, the ammonite succession in Alaska from Hettangian to Toarcian time is essentially the same as that in central and northern Europe. It is also similar to that in the Mediterranean region except for certain ammonites of late Pliensbachian Age. The Bajocian ammonite succession in Alaska is likewise similar to that in the above-named regions except for a few genera known only from areas bordering the Pacific Ocean. In contrast, the Bathonian succession in Alaska is dominated by the Boreal genera *Cranoccephalites*, *Arctoccephalites*, and *Arcticoceras*, which are unknown in central Europe or farther south. The Callovian ammonite genera of Alaska are likewise dominantly Boreal in origin, but, as in the Bajocian and Bathonian ammonites, include some genera that are known only from the Western Hemisphere. During the rest of Jurassic time, ammonites became uncommon in Alaska but include the Boreal genera *Cardioceras* of early Oxfordian Age and *Amoeboceras* of late Oxfordian to early Kimmeridgian Age.

FOSSIL OCCURRENCES BY STAGES

Locality in figure 2	Description
Hettangian	
3----	USGS Mesozoic locs. 29737 and 29738. From Spike Mountain near the Porcupine River about 30 miles north of Old Rampart, Coleen quadrangle, lat 63°35.75' N., long 141°40' W. <i>Psiloceras</i> .
8----	USGS Mesozoic locs. 29890 and 29891. About 15 miles northeast of McCarthy, northern part of sec. 26, T. 4 S., R. 16 E., near NW. cor. McCarthy B-4 quadrangle, southern Wrangell Mountains, eastern Alaska. <i>Wachneroceras</i> and <i>Psiloceras</i> .

Locality in figure 2	Description
Hettangian—Continued	
11----USGS Mesozoic loc. 29874. East bank of Allen River at middle rapids, Tikchik Lakes area, lat 60°7' N., long 158°38' W., Taylor Mountains A-8 quadrangle. Kuskokwim region, southwest Alaska. <i>Weyla</i> and other pelecypods identical with species in basal Jurassic near Seldovia on the Kenai Peninsula (loc. 13). These could be of Hettangian or Sinemurian Age.	
12----USGS Mesozoic loc. 24357. Hagermeister Island in Bristol Bay area, lat 58°43'50" N., long 160°58'10" W. Kuskokwim region, southwest Alaska. <i>Weyla</i> and other pelecypods identical with species near Seldovia (loc. 13). These could be Hettangian or Sinemurian Age.	
13----USGS Mesozoic locs. 21242 and 22664. From 2.8 to 2.95 miles S. 67° W. of Pt. Naskowhak on south shore of Kachemak Bay, Seldovia B-5 quadrangle, Kenai Peninsula. <i>Waehneroceras</i> .	
14----USGS Mesozoic locs. 3110, 12394, 19803, 25694 and 29268 on peninsula between Puale Bay and Alinchak Bay about 1 mile N. 65° W. of Cape Kekurnoi, Alaska Peninsula. <i>Waehneroceras</i> .	
15----Richfield Oil Co. Wide Bay test well 1, core 5 at depth of 2,235 to 2,236 feet, on north side of Wide Bay, Alaska Peninsula. <i>Waehneroceras</i> .	

Sinemurian

- 1----Avak test well 1 at depth of 1,836 feet on Point Barrow, northern Alaska (See Imlay, 1955, p. 82, 87). *Arietites*.
- 2----USGS Mesozoic loc. 29774. On Blankenship Creek in Lisburne Ridge area west of Etivluk River, lat 68°37.5' N., long 156°42.5' W., northwest Alaska. *Cruciloboceras*.
- 4----USGS Mesozoic loc. 16229. Head of small eastern tributary of Partin Creek that heads against Little Shotgun Creek north of Eldridge Glacier on west side of Chulitna Valley, eastern Alaska. *Arnioceras*.
- 5----USGS Mesozoic loc. 26722. Talkeetna Mountains A-2 quadrangle, coordinates 1.20, 17.23, lat 62°59'58" N., long 147°57'48" W. Mesozoic loc. 28661 in Talkeetna Mountains A-3 quadrangle, coordinates 15.35, 15.63, lat 62°13'29" N., long 148°01'27" W. Talkeetna Mountains, Cook Inlet region. *Cruciloboceras*.
- 6----USGS Mesozoic loc. 27586. Carbon Creek, 2.3 miles S. 7° E. of its mouth, which enters Matanuska River from south near mouth of Chickaloon River, near south boundary of sec. 12, T. 19 N., R. 5 E., Anchorage D-4 quadrangle, Cook Inlet region. *Arnioceras*.
- 7----USGS Mesozoic loc. 30137. Near center of north half of SW¼ sec. 8, T. 4 S., R. 11 E., in south-central part of McCarthy (C-7) quadrangle. Mesozoic locs. 30138-30140 in west half of sec. 17, T. 4 S., R. 12 E., near southeast corner of McCarthy (C-7) quadrangle. Wrangell Mountains, eastern Alaska. All localities are in upper member of McCarthy Formation. *Arnioceras* (loc. 30140), *Megarietites*?, and *Arnioceras*? (loc. 30139), and arietitid ammonites (locs. 30137 and 30138).

Locality in figure 2	Description
Sinemurian—Continued	
8----Many localities in upper member of McCarthy Formation, in McCarthy C-4, C-5, and B-4, quadrangles, Wrangell Mountains, eastern Alaska (MacKevett, 1970a, b, 1971). Ammonites include <i>Arietites</i> (loc. 29870), <i>Arnioceras</i> (locs. 28535, 28536, 28538, 28688), <i>Cruciloboceras</i> (loc. 28537, 28540, 28671-73), and <i>Microderoceras</i> (loc. 14472).	
9----USGS Mesozoic loc. 29773. Float, sec. 37, T. 23 S., R. 35 E., Yakutat D-4 quadrangle, southeast Alaska. <i>Cruciloboceras</i> .	
10----Property of State Museum at Juneau. Ammonite found near center of face of Norris Glacier about 14 miles northeast of Juneau. Juneau B-1 quadrangle, southeast Alaska. <i>Xiphoceras</i> ?	
11----Described under Hettangian fossil occurrences.	
12----Do.	
13----USGS Mesozoic loc. 2981. Two miles west of Pt. Naskowhak on west side of Seldovia Bay. Seldovia B-5 quad., Kenai Peninsula. <i>Coroniceras</i> .	
14----USGS Mesozoic locs. 3111, 12396, and 21237 on east side of Puale Bay about 8,400 feet northwest of Cape Kekurnoi, Alaska Peninsula. <i>Coroniceras</i> .	

Locality in figure 3	Description
Pliensbachian	
1----South Barrow test well 3 at depths of 2,069 to 2,198 feet, Point Barrow, northern Alaska (Imlay, 1955, p. 82, 87; Howarth, 1958, p. XXVI). <i>Amaltheus</i> spp.	
2----Simpson test well 1 at depth of 5,680 feet, southeast of Point Barrow, northern Alaska. (Imlay, 1955, p. 82, 87). <i>Amaltheus</i> sp.	
4----USGS Mesozoic loc. M2441. Ipewik River, De Long Mountains, lat 68°37' N., long 164°11' W. Mesozoic loc. 29164 on east bank Ipewik River, lat 68°40' N., long 164° 13' W., northern Alaska. <i>Amaltheus</i> sp.	
7----USGS Mesozoic loc. 29165. Three miles east of Shrader Lake, lat 69°22' N., long 144°48' W., near head of Sadlerochit River, northern Alaska. <i>Amaltheus</i> .	
8----USGS Mesozoic loc. 30074. West side of Aichilik River about 2 miles west of VABM ATTE, lat 69° 33' N., long 143°05' W., from 800-1,000 feet above base of Kingak Shale. <i>Amaltheus</i> cf. <i>A. stokesi</i> (J. Sowerby).	
9----USGS Mesozoic loc. 29340. Float from bluff on the Middle Salmon Trout River near Old Rampart, Kandik area, east central Alaska. <i>Amaltheus</i> .	
10----Many USGS Mesozoic localities in the Talkeetna Mountains A-3 quadrangle and the Anchorage D-1 and D-2 quadrangles, Talkeetna Mountains, Cook Inlet region. <i>Paltarpites</i> , <i>Harpoceras</i> , <i>Arietoceras</i> , <i>Leptaleoceras</i> , <i>Fontanelliceras</i> , <i>Amaltheus</i> , <i>Protogrammoceras</i> , <i>Fanninoceras</i> , and <i>Acanthopleuroceras</i> .	
11----USGS Mesozoic loc. 28675, SW¼SW¼ sec. 23, T. 4 S., R. 16 E., McCarthy C-4 quadrangle. Less than 100 feet below Lubbe Creek Formation. <i>Uptonia</i> . Mesozoic loc. 28531 on east side of McCarthy Creek, probably a little northwest of center of sec. 25, T. 3 S., R. 14 E., McCarthy C-5, quadrangle, southern Wrangell Mountains, Lubbe Creek Formation. <i>Arietoceras</i> .	

Locality in figure 3	Description
Toarcian	
1-----	South Barrow test well 3 at depths of 1,772 to 2,063 feet at Point Barrow, northern Alaska (Imlay, 1955, p. 82, 87-89). <i>Catacoeloceras</i> at 2,063 feet. <i>Dactylioceras</i> at 1,772 to 2,018 feet.
3-----	USGS Mesozoic locs. 29159-29161, 29163. North bank of Thetis Creek, lat 68°41'50" N., long 164°45'05" W., northwestern Alaska. <i>Harpoceras</i> cf. <i>H. exaratum</i> (Young and Bird) and <i>Dactylioceras</i> . Mesozoic loc. 29776 near Thetis Creek, lat 68°40.8' N., long 164°45.5' W., <i>Harpoceras</i> .
5-----	USGS Mesozoic loc. 23772 and 24035 (Imlay, 1955, p. 80, 89). <i>Pseudolioceras</i> .
6-----	USGS Mesozoic loc. 22081 (Imlay, 1955, p. 80). <i>Harpoceras</i> cf. <i>H. exaratum</i> (Young and Bird) (not <i>Pseudolioceras</i>).
10-----	Many USGS Mesozoic localities in the Talkeetna Mountains A-1, A-2, and A-3 quadrangles, the Anchorage D-2 and D-5 quadrangles, and the Valdez D-8 quadrangle, Talkeetna Mountains, Cook Inlet region. <i>Dactylioceras</i> , <i>Harpoceras</i> , <i>Grammoceras</i> , <i>Haugia</i> , <i>Phymatoceras</i> , and <i>Pseudolioceras</i> .
11-----	USGS Mesozoic locs. 28546-28548, 28678, and 28679 in southern part of McCarthy C-5 quadrangle, southern Wrangell Mountains. Upper part of Lubbe Creek Formation. <i>Weyla dufrenoyi</i> (d'Orbigny), identified by S. W. Müller.
12-----	USGS Mesozoic loc. 19804. East shore of Puale Bay about 9,500 feet northwest of Cape Kekurnoi, Alaska Peninsula. <i>Haugia</i> .

Locality in figure 4	Description
Bajocian	
1-----	South Barrow test well 2 at depth of 2,391 feet near Point Barrow, northern Alaska (Imlay; 1955, p. 82, 89). <i>Tmetoceras</i> .
2-----	Topogarak test well 1 at depth of 8,113 feet about 58 miles south-southeast of Point Barrow, lat 70°37'30" N., long 155°53'36" W., northern Alaska (Imlay, 1955, p. 82, 89). <i>Tmetoceras</i> .
3-----	USGS Mesozoic loc. 24035. Canning River, lat 69°34' N., long 146°23' W., northern Alaska (Keller and others, 1961, p. 193). <i>Pseudolioceras maclintocki</i> (Haughton) and <i>Oxytoma jacksoni</i> (Pompeckj).
4-----	USGS Mesozoic locs. 29152-29156. West end of Ignek Valley, lat 69°41' N., long 145°36' W. to lat 69°33' N., long 145°43' W., northern Alaska. <i>Pseudolioceras maclintocki</i> (Haughton), <i>Erycitoides</i> cf. <i>E. howelli</i> (White), <i>Canavarella</i> cf. <i>C. belophora</i> Buckman, and <i>Inoceramus</i> cf. <i>I. lucifer</i> Eichwald.
5-----	USGS Mesozoic loc. 29147 at junction of Sadlerochit and Kekiktuk Rivers, lat 69°33' N., long 144°43' W.; Mesozoic loc. 29150 on Fire Creek in central part of Ignek Valley, lat 69°32' N., long 145°09' W.; Mesozoic loc. 29880 on Fire Creek, lat 69°32'22" N., long 145°12'30" W.; Mesozoic locs. 29884 to 29886 on Kaviak Creek, 0.7 miles SW. of Sadlerochit River, lat 69°29' N., long 145°03' W.; Mesozoic locs. 10307 and 10308 (Imlay, 1955, p. 81). Northeastern Alaska. <i>Pseudolioceras maclintocki</i> (Haugh-

Locality in figure 4	Description
Bajocian—Continued	
	ton), <i>Erycitoides</i> , <i>Canavarella</i> (loc. 29884 only), and <i>Inoceramus lucifer</i> Eichwald.
6-----	USGS Mesozoic loc. 29157 on Canning River near Shublik Island, lat 69°24' N., long 146°10' W., northern Alaska, Mesozoic locs. 21023 and 22595 (Imlay, 1955, p. 80). <i>Pseudolioceras whiteavesi</i> (White) (loc. 21023 only) and <i>P. maclintocki</i> (Haughton); Mesozoic locs. 29138-29142, Canning River near Shublik Island, lat 69°24' N., long 146°10' W., northeastern Alaska. <i>Arkelloceras</i> ; Mesozoic locs. 21024, 22597, and 24033 on west bank of Canning River opposite mouths of Eagle and Cache Creeks, lat 69°27' N., long 146°13'30" W. (Imlay, 1955, p. 80, 91; Keller and others, 1961, p. 193). <i>Arkelloceras</i> .
7-----	USGS Mesozoic loc. 30135 on ridge west of Okerokvik River, 7.6 miles S. 80° W. of VABM ATTE, lat 69°29'45" N., long 143°26' W. <i>Pseudolioceras maclintocki</i> (Haughton).
8-----	USGS Mesozoic loc. 21552. Cutbank on Fortress Creek, lat 68°31' N., long 153°03' W. (Patton and Tailleir, 1964, p. 444). <i>Pseudolioceras</i> ? and <i>Inoceramus lucifer</i> Eichwald.
9-----	USGS Mesozoic loc. 22591. Cutbank on Tiglukpuk Creek, lat 68°20' N., long 151°50' W. (Patton and Tailleir, 1964, p. 444). <i>Arkelloceras</i> ? sp. juv. and <i>Inoceramus</i> cf. <i>I. lucifer</i> Eichwald.
10-----	USGS Mesozoic loc. M1717 (Imlay, 1967, p. B8), about 3 miles east-southeast of Nation. <i>Otapiria</i> sp. undet. and <i>Pentacrinus subangularis alaska</i> Springer. This variety of <i>Pentacrinus</i> has been found in the northeast Alaska near the Sadlerochit River associated with <i>Inoceramus</i> cf. <i>I. lucifer</i> Eichwald (Mesozoic loc. 29885) and directly above beds containing <i>Canavarella</i> , <i>Pseudolioceras</i> cf. <i>P. maclintocki</i> (Haughton), and <i>Inoceramus</i> cf. <i>I. lucifer</i> Eichwald (Mesozoic loc. 29884) of early Bajocian Age. As the full stratigraphic range of the crinoid subspecies has not yet been established, its presence alone is not proof of a Bajocian Age.
11-----	USGS Mesozoic loc. 13430. Northwest side of Innoko River, about 30 miles airline upstream from confluence of Innoko and Iditarod Rivers, Innoko District, Kuskokwim region, Alaska, <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald.
12-----	USGS Mesozoic loc. 27716. East bank of Yukon River about 7 miles N. 25° W. from village of Ohogament, coordinates 0.65, 11.35, lat 61°39' N., long 161°55.5' W., Russian Mission quadrangle, Kuskokwim region. <i>Stephanoceras</i> ? sp. Probably Middle Jurassic.
13-----	USGS Mesozoic loc. 21481 on south side of Tuluksak River, 2.7 miles S. 55° W. of Nyac, and 1.5 miles N. 50° W. of Nyac hydroelectric power plant; Mesozoic loc. 21029 on north side of Tuluksak River about 1 mile north of loc. 21481, Bethel quadrangle, Kuskokwim region, southwest Alaska (Hoare and Coonrad, 1959a). <i>Inoceramus ambiguus</i> Eichwald.
14-----	USGS Mesozoic loc. 27091. Crest of low ridge 12.7 miles S. 59° W. from junction of Buckstock and

Locality in figure 4	Description
Bajocian—Continued	
	Aniak Rivers and 6.8 miles N. 55° E. of Mt. Hamilton, coordinates 20.5, 4.4. Russian Mission quadrangle, Kuskokwim region, southwest Alaska. <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald.
15----	USGS Mesozoic loc. 21031. About 3 miles west of Salmon River, about 23.5 miles S. 36° E. of Mt. Hamilton, Kuskokwim region, southwest Alaska (Hoare and Coonrad, 1959b). <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald.
16----	USGS Mesozoic loc. 20717. About 4½ miles S. 6° E. of confluence of Cripple Creek with Salmon River, about 23.5 miles E. 36° E. of Nyac, Bethel quadrangle, Kuskokwim region, southwest Alaska (Hoare and Coonrad, 1959b). <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald.
17----	USGS Mesozoic loc. 19729. Cutbank southeast of Holokuk River, 27 miles S. 20° E. of Napaiment and 3.6 miles upstream (southwest) from mouth of Boss Creek, Kuskokwim region, southwest Alaska. <i>Inoceramus</i> sp.
18----	USGS Mesozoic loc. 19395b. East of Holitna River, 4½ miles S. 42° E. of Nagamut, Kuskokwim region, southwest Alaska. <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald.
19----	USGS Mesozoic loc. 29889. South-central part of Hagemeister Island, Bristol Bay, lat 58°38'25" N., long 160°58'50" W. <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald.
20----	USGS Mesozoic loc. 28682, in part. Near center SE¼NW¼ sec. 20, T. 3 S., R. 16 E., McCarthy C-5 quadrangle, southern Wrangell Mountains, eastern Alaska (MacKevett, 1963; 1971). <i>Normannites</i> cf. <i>N. variabilis</i> Imlay and <i>Teloceras</i> cf. <i>T. blagdeni</i> (Sowerby) obtained with <i>Cranocephalites</i> , <i>Parareineckeia</i> , and <i>Cobbanites</i> as float derived from a fairly thin sequence.
21----	Many USGS Mesozoic localities in Talkeetna Mountains A-1 and A-2 quadrangles and Anchorage D-2, D-3, and D-4 quadrangles, Talkeetna Mountains, Cook Inlet region (Imlay, 1962a, p. A5, A6, 1964b, p. B2, B22, B27).
22 and	
23----	Many USGS Mesozoic localities from area between Iniskin Bay and Tuxedni Bay, north side of Cook Inlet (Imlay, 1961a, p. A-5, 1964b, p. B22-B27).
24----	USGS Mesozoic loc. 29115. Iliamna Lake region, 2.1 miles N. 30° E. of mouth of Amakdedori Sreek, lat 59°18'25" N., long 154°05'40" W., coordinates 1.25, 4.95, Iliamna B-3 quadrangle, Cook, Inlet region. <i>Inoceramus ambiguus</i> Eichwald.
25----	USGS Mesozoic loc. 12390. Half a mile south of head of Alinchak Bay, Alaska Peninsula. <i>Inoceramus lucifer</i> Eichwald (5 specimens) and <i>I. ambiguus</i> Eichwald (1 specimen found loose at bottom of cliff).
26----	Many localities near Wide Bay, Alaska Peninsula (Imlay, 1964b, p. B18; Westermann, 1964, 1969).

Locality in figure 5	Description
Bathonian	
1----	USGS Mesozoic locs. 28817, 29143, 29144, 29435, 29855, 29875 to 29877. Central part of Ignek Valley, lat 69°33'30"-34' N., long 145°20' W., northeastern Alaska. <i>Arcticoceras</i> , <i>Arctocephalites</i> , <i>Cranocephalites</i> , and <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald.
2----	USGS Mesozoic loc. 22083. North bank of Sadlerochit River, between Gravel and Fire Creeks, lat 69°31' N., long 145°02' W., northeastern Alaska. <i>Arctocephalites</i> ?
3----	USGS Mesozoic loc. 21023 (in part), Canning River at mouth of Eagle and Cache Creeks, lat 69°25' N., long 146°08' W., <i>Arcticoceras</i> ; Mesozoic loc. 22596, west side of Canning River, lat 69°23'30" N., long 146°07'30" W., <i>Arcticoceras</i> ; Mesozoic loc. 29146, Canning River near Shublik Island, lat 69°24' N., long 146°10' W. <i>Arctocephalites</i> .
4----	USGS Mesozoic loc. 29145, west end of Ignek Valley, lat 69°33' N., long 145°50' W. <i>Arctocephalites</i> .
5----	USGS Mesozoic loc. 30075. Aichilik River, lat 69°33' N., long 143°05' N., northeastern Alaska. <i>Arcticoceras ishmae</i> (Keyserling) and <i>Choffatia</i> ? sp.
7----	Many USGS Mesozoic localities in the Nizina Mountain Formation in southeastern part of McCarthy C-5 quadrangle, southern Wrangell Mountains (MacKevett, 1969, p. A42-A45). <i>Parareineckeia</i> , <i>Cobbanites</i> , <i>Cranocephalites</i> , and <i>Inoceramus</i> cf. <i>I. ambiguus</i> Eichwald. (Most collections are from float.)
8----	Seven USGS Mesozoic localities in the Talkeetna Mountains A-2 quadrangle and one in the Anchorage D-4 quadrangle (Imlay, 1962a, p. C3-C5, C16), Talkeetna Mountains, Cook Inlet region, <i>Cranocephalites</i> , <i>Parareineckeia</i> , <i>Cobbanites</i> , <i>Oecotraustes</i> , and <i>Inoceramus ambiguus</i> Eichwald.
9----	Peninsula between Tuxedni Bay and Chinitna Bay, west side of Cook Inlet. <i>Parareineckeia</i> and <i>Cranocephalites</i> in lower two-thirds of Bowser Formation (Imlay, 1962a, p. C16, 17, 20; Detterman and Hartsock, 1966, p. 38, 39) are overlain by beds containing <i>Arctocephalites</i> cf. <i>A. elegans</i> Spath (Mesozoic loc. 22699).
10----	Iniskin Peninsula, west side of Cook Inlet. <i>Parareineckeia</i> and <i>Cranocephalites</i> in the lower two-thirds of the Bowser Formation are overlain by beds containing <i>Kepplerites</i> ? (Mesozoic loc. 22553) (Imlay, 1962a, p. 16, 17, 20; Detterman and Hartsock, 1966, p. 38, 39).
Callovian	
5----	USGS Mesozoic loc. 30136. Float from beds directly overlying the beds at Mesozoic loc. 30075. <i>Cadoceras</i> sp.
6----	USGS Mesozoic loc. 22745, West Fork Ivishak River, lat 69°0' N., long 148°04'30" W. (Keller and others 1961, p. 193; Imlay, 1955, p. 80, 90). <i>Pseudocadoceras grewinkii</i> (Pompeckj)?, or possibly immature <i>Arcticoceras</i> .

Locality in figure 5	Description
Callovian—Continued	
8----	Many USGS Mesozoic localities in the Talkeetna Mountains A-1 and A-2 quadrangles and the Anchorage D-3 quadrangle, Talkeetna Mountains, Cook Inlet region. Ammonites include most of the same genera and species described by Imlay (1953b, p. 50).
9, 10----	Many localities (see Imlay, 1953b, p. 65-69).
11----	USGS Mesozoic loc. 29271, lat 57°57.3' N., long 155°4.4' W., Kashvik Bay, Karluk quadrangle Alaska Peninsula. <i>Cadoceras</i> (<i>Stenocadoceras</i>) <i>multicostatum</i> Imlay.
12----	Many USGS Mesozoic localities in the Puale Bay area, Alaska Peninsula (Imlay, 1953b, p. 63 and table 6). Common ammonites include <i>Cadoceras tenuicostatum</i> Imlay, <i>C. doroschini</i> (Eichwald), and many species of <i>C.</i> (<i>Stenocadoceras</i>) and <i>Pseudocadoceras</i> . The abundance of <i>C.</i> (<i>Stenocadoceras</i>) and the absence of <i>C.</i> (<i>Paracadoceras</i>) show that the beds are equivalent to the upper part of the Chinitna Formation north of Cook Inlet.
13----	A few USGS Mesozoic localities between Jute Bay and Portage Bay, Alaska Peninsula (Imlay, 1953b, p. 63 and table 6). <i>Cadoceras</i> (<i>Stenocadoceras</i>) <i>multicostatum</i> Imlay and <i>Pseudocadoceras grewinkii</i> (Pompeckj).
14----	Many localities near Wide Bay, Alaska Peninsula (Imlay, 1953b, p. 64, 70, 71, and table 6). Same ammonites present as in the Chinitna Formation north of Cook Inlet.
15----	USGS Mesozoic loc. 29342. Axial region of Chignik Bay anticline near Conglomerate Creek. Approximate lat 65°26'13" N., long 158°36'48" W., Chignik Bay, Alaska Peninsula. <i>Xenoccephalites vicarius</i> Imlay.

Locality in figure 6	Description
Early Oxfordian	
1----	USGS Mesozoic loc. 29856. Ignek Mesa in Ignek Valley, 6.8 miles S. 75° E. of Katakturuk Canyon, lat 69°33'30" N., long 145°20' W., northeast Alaska. <i>Cardioceras</i> (<i>Scarburgiceras</i>).
2----	USGS Mesozoic loc. 29137. Fire Creek in Ignek Valley, lat 69°32' N., long 145°09' W., northeast Alaska. <i>Cardioceras</i> (<i>Scarburgiceras</i>).
3----	USGS Mesozoic locs. 24165, 24166, 24168, 24837, and 24841. Near headwaters of Little Nelchina River in central and east-central part of Talkeetna Mountains A-2 quadrangle, Talkeetna Mountains, Cook Inlet region. <i>Cardioceras</i> (<i>Scarburgiceras</i>) <i>martini</i> Reeside and <i>C. distans</i> (Whitfield).
4, 5----	Many localities between Iniskin Bay and the Hickerson Lake area west of Cook Inlet (Detterman and Hartsock, 1966, p. 50, 51). <i>Cardioceras</i> (<i>Scarburgiceras</i>) <i>martini</i> Reeside, <i>C. distans</i> Whitfield, and <i>C.</i> (<i>Scoticardioceras</i>) <i>alaskense</i> Reeside.

Locality in figure 6	Description
Early Oxfordian—Continued	
6----	USGS Mesozoic loc. 10798. Five miles southeast of Mount Peulik, west of Puale Bay and north of Wide Bay, Alaska Peninsula. <i>Cardioceras</i> cf. <i>C. canadense</i> Whiteaves.
Locality in figure 7	Description
Late Oxfordian to early Kimmeridgian	
1----	USGS Mesozoic loc. 22126 (Imlay, 1955, p. 78, 79). Mesozoic loc. 30071 on Kukpowruk River west of Igloo Mountain, lat 68°43' N., long 163°12'30" W., <i>Buchia concentrica</i> (Sowerby).
2----	USGS Mesozoic loc. 22759 (Imlay, 1955, p. 79, 80). <i>Buchia concentrica</i> (Sowerby).
3----	USGS Mesozoic loc. 21026 (Imlay, 1955, p. 79, 80). <i>Buchia concentrica</i> (Sowerby).
4----	Many USGS Mesozoic localities near the Canning River, northeast Alaska. Mesozoic locs. 24014, 21028, 25598, and 29882 (Imlay, 1955, p. 79, 80; Keller and others, 1961, p. 193) from lat 69°30'45" N. to 69°33' N.; long 146°18' W. to 146°23' W.; Mesozoic locs. 29134-29136 near Shublik Island at lat 69°24' N., long 146°10' W. <i>Amoeboceras</i> (<i>Prionodoceras</i>) and <i>Buchia concentrica</i> (Sowerby).
5----	USGS Mesozoic loc. 29133. North side of central part of Sadlerochit Mountains, lat 69°41' N., long 144°50' W. <i>Buchia</i> cf. <i>B. concentrica</i> (Sowerby).
6----	USGS Mesozoic locs. 5723, 16921, 16922, 18082, 18086, and 18089. Area between Slana and Nabesna in eastern part of Alaska Range (Moffit and Knopf, 1910, p. 29; Moffit, 1938b, p. 32; Moffit, 1954, p. 131). <i>Buchia concentrica</i> (Sowerby).
7----	USGS Mesozoic loc. 5772. Trail along Notch Creek northwest of Chisana, eastern part of Alaska Range (Moffit and Knopf, 1910, p. 29; Moffit, 1954, p. 131). <i>Buchia concentrica</i> (Sowerby).
8----	Many localities in the McCarthy C-5 quadrangle, Wrangell Mountains (MacKevett, 1969, p. A41, A45-49). <i>Buchia concentrica</i> (Sowerby) and <i>Amoeboceras</i> (<i>Prionodoceras</i>) from lower part of Root Glacier Formation.
9----	Many localities in the Talkeetna Mountains A-1 quadrangle, Cook Inlet region. <i>Buchia concentrica</i> (Sowerby).
10----	Richfield Oil Company—Swanson River unit 2 well, SW¼SE¼ sec. 22, T. 8 N., R. 9 W., Seward Meridian, Kenai Peninsula, Alaska. <i>Buchia concentrica</i> (Sowerby) in cores from depths of 11,794-11,804 feet. Also, a species of <i>Pseudolimea</i> , very common in the Naknek Formation, occurs in cores from depths of 11,846-11,856 feet.
11, 12----	Many localities in Snug Harbor Siltstone Member of the Naknek Formation between Iniskin Bay and Tuxedni Bay, Cook Inlet region (Martin, 1926, p. 178-180; Detterman and Hartsock, 1966, p. 51). <i>Amoeboceras</i> (loc. 3046 only) and <i>Buchia concentrica</i> (Sowerby).

Locality in figure 7	Description
Late Oxfordian to early Kimmeridgian—Continued	
13----USGS Mesozoic loc. 28520, 1.22 miles N. 30° E. of tip of Ursus Head; Mesozoic loc. 28521, 0.22 mile N. 62° E. of tip of Ursus Head. Ursus Cove on west side of Cook Inlet. <i>Buchia concentrica</i> (Sowerby).	
14----USGS Mesozoic loc. 29899. On east tip of westernmost island 1 mile N. 40° E. of mouth of Douglas River, lat 59° 64' 20" N., long 153° 44' 50" W. Iliamna quadrangle, Alaska Peninsula. <i>Buchia concentrica</i> (Sowerby) and <i>Perisphinctes</i> (<i>Dichotomosphinctes</i>).	
15----USGS Mesozoic loc. 25836. West side of Kaguyak Bay, Alaska Peninsula. <i>Buchia concentrica</i> (Sowerby).	
16----USGS Mesozoic loc. 29248 at western base of Barrier Range north of Fultons Falls, about 1 mile S. 12° E. of loc. 29247 and 8 miles north of Katmai Bay, Mt. Katmai quadrangle, Alaska Peninsula. <i>Buchia concentrica</i> (Sowerby).	
17----USGS Mesozoic loc. 29245. North side of Kashvik Bay about 1.7 miles east-northeast of head of bay and 3.3 miles S. 27° E. of Atmo Mountain, Karluk quadrangle, Alaska Peninsula. <i>Buchia concentrica</i> (Sowerby).	
18----USGS Mesozoic locs. 3133–3136, 10825, 11333, 12276, 21353, and many others. Puale Bay area, Karluk quadrangle, Alaska Peninsula (Martin, 1926, p. 212–218). <i>Buchia concentrica</i> (Sowerby).	
19----Many localities north and west of Wide Bay, Ugashik quadrangle, Alaska Peninsula (Martin, 1926, p. 212–218). <i>Buchia concentrica</i> (Sowerby).	
20----USGS Mesozoic loc. 25696, 25698, 25708, 25710, 25711, 25713–25715. Chignik Bay area, Chignik quadrangle, Alaska Peninsula. <i>Buchia concentrica</i> (Sowerby).	
21----USGS Mesozoic loc. M 1213. Knife Peak area west of Chignik Bay, Chignik quadrangle, Alaska Peninsula (Burk, 1965, p. 160, 176, 219). <i>Buchia concentrica</i> (Sowerby).	
22----USGS Mesozoic loc. M 1197, M 1224, Staniukovich Mountain, Port Moller quadrangle, Alaska Peninsula (Burk, 1965, p. 160, 166, 219). <i>Buchia concentrica</i> (Sowerby).	

Locality in figure 8	Description
Late Kimmeridgian to early middle Tithonian	
1----USGS Mesozoic loc. 29131, headwaters of tributary to Thetis Creek, lat 68° 42' 30" N., long 164° 40' 30" W. <i>Buchia mosquensis</i> (von Buch); Mesozoic loc. 29132, south-flowing tributary to Epewik River, lat 68° 38' 20" N., long 164° 36' 00" W., northwest Alaska. <i>Buchia</i> cf. <i>B. mosquensis</i> (von Buch).	
2----USGS Mesozoic loc. 22127. Kukpowruk River, lat 68° 42' 30" N., long 163° 14' W. (Imlay, 1955, p. 78, 79). <i>Buchia rugosa</i> (Fischer).	
3----USGS Mesozoic loc. 22776. Easternmost fork of Driftwood Creek, lat 68° 40' N., long 160° 26' 28" W., northwest Alaska (Imlay, 1955, p. 78, 79). <i>Buchia rugosa</i> (Fischer) and <i>B. mosquensis</i> (von Buch).	

Locality in figure 8	Description
Late Kimmeridgian to early middle Tithonian—Continued	
4----USGS Mesozoic loc. 23598. Ipnayik River, 2 miles south of Memorial Creek, lat 68° 23' N., long 157° 15' W., northwest Alaska (Imlay, 1955, p. 78, 79). <i>Buchia</i> cf. <i>B. mosquensis</i> (von Buch) and <i>B. concentrica</i> (Sowerby).	
5----USGS Mesozoic locs. 22750, 22751, 22766, 22768, 22769. Lupine River from lat 68° 49' 30" N. to 68° 51' N., long 148° 17' 30" W. to 148° 22' 30" W. (Keller and others, 1961, p. 195). <i>Buchia rugosa</i> (Fischer), <i>B. mosquensis</i> (von Buch), and <i>B. concentrica</i> (Sowerby).	
6----USGS Mesozoic loc. 22746. Small divide nearly 1 mile east of Nosebleed Creek, lat 68° 52' N., long 148° 08' W., northeast Alaska (Keller and others, 1961, p. 195). <i>Buchia rugosa</i> (Fischer).	
7----USGS Mesozoic loc. 21027, Shaviovik River, main fork of most easterly branch, lat 69° 22' N., long 146° 32' W. (Imlay, 1955, p. 79, 80). <i>Buchia rugosa</i> (Fischer) and <i>B. mosquensis</i> (von Buch). Mesozoic loc. 24028 from Kavik River, lat 69° 24' 45", long 146° 37' 30" W., northeast Alaska. <i>Buchia rugosa</i> (Fischer) and <i>B. concentrica</i> (Sowerby).	
8----USGS Mesozoic locs. 29872, 29897, and 30076. Ridge south of Joe Creek, 11.5 miles N. 75° W. of point where creek crosses International Boundary, Table Mtn. quadrangle, lat 68° 58' N., long 146° 38' W., northeast Alaska. <i>Buchia rugosa</i> (Fischer).	
9----USGS Mesozoic loc. 16085. Head of Lost Creek, lat 62° 36' N., long 142° 59' W., area between Slana and Nebesna, eastern Alaska Range. <i>Buchia rugosa</i> (Fischer).	
10----Many USGS Mesozoic localities in the McCarthy C-5 quadrangle, Wrangell Mountains (MacKevett, 1969, p. A41, A45–A49). <i>Buchia rugosa</i> (Fischer) from upper part of Root Glacier Formation.	
11----USGS Mesozoic locs. 3271, 3309, 8851, 10095, 10169, 27068, 27071–73, 27334, and 27335 from Pybus Bay area in southeastern part of Admiralty Island, southeast Alaska (Loney, 1964, p. 97). <i>Buchia rugosa</i> (Fischer).	
12----USGS Mesozoic locs. 11401 and 11935 from north end of Kupreanoff Island between Hamilton Bay and Keku Strait, southeast Alaska (Martin, 1926, p. 379, 380). <i>Buchia rugosa</i> (Fischer).	
13----USGS Mesozoic locs. 9528–30 on Gravina Island, southeast Alaska. <i>Buchia rugosa</i> (Fischer) and <i>B. cf. B. mosquensis</i> (von Buch).	
14----Many localities in the Talkeetna Mountains A-1 quadrangle, Cook Inlet region. <i>Buchia rugosa</i> (Fischer) and <i>B. mosquensis</i> (von Buch).	
15----USGS Mesozoic locs. 20714 and 27092. Near top of ridge, 2,800 feet west of Discovery Creek, 2 miles S. 69° E. of Mt. Hamilton, and 18.3 miles S. 53° W from confluence of Buckstock and Aniak Rivers. Coordinates 19.6, 3.00 in Russian Mission quadrangle, Kuskokwim region, southwest Alaska. <i>Buchia rugosa</i> (Fischer) and <i>B. mosquensis</i> (von Buch).	

Locality in figure 8	Description
Late Kimmeridgian to early middle Tithonian—Continued	
16----	USGS Mesozoic loc. 24257. Thirty-two miles S. 80°45' E. from Kwinhagak and 55 miles N. 58° 25' E. from Platinum, Goodnews quadrangle, Kuskokwim region, southwest Alaska. <i>Buchia</i> cf. <i>B. rugosa</i> (Fischer).
17----	USGS Mesozoic locs. 24957, 25839, 29250, 29253, M 5341, M 5345, and M 5348. Cliffs on south side of Kamashak Bay extending 2–10 miles west-northwest from the mouth of Douglas River, lat 59°03'45" N. to 59°04'30" N., long 153°49'30" W. to 154°02' W., Iliamna quadrangle, head of Alaska Peninsula. <i>Buchia rugosa</i> (Fischer) and <i>B. mosquensis</i> (von Buch).
18----	USGS Mesozoic loc. 24955 from sea cliff on west side of mouth of Douglas River, coordinates 19.98–1.32; Mesozoic loc. 25832 on westernmost of two offshore islands 0.6 mile N. 35° E. of mouth of Douglas River; Mesozoic loc. 25833 near mouth of Douglas River, coordinates 19.8–0.9, Iliamna quadrangle, head of Alaska Peninsula. <i>Buchia rugosa</i> (Fischer) and <i>B. mosquensis</i> (von Buch).
19----	USGS Mesozoic locs. 24956, 29249, 29254 on west shore of Kaguyak Bay. Mesozoic loc. 24956 is 1 mile north of abandoned village of Kaguyak, coordinates 0.82, 10.38; Mesozoic loc. 29254 is on small island 0.7 mile S. 18° E. of abandoned village, Afognak quadrangle, Alaska Peninsula. <i>Buchia mosquensis</i> (von Buch) and <i>B. rugosa</i> (Fischer).
20----	USGS Mesozoic locs. 25276–25278 and M 2076. On northeast side of Grosvenor Lake, lat 58°41' N., long 155°7'–10' W., Mt. Katmai quadrangle, Alaska Peninsula. <i>Buchia rugosa</i> (Fischer) and <i>B. mosquensis</i> (von Buch) range from just below Chisik Conglomerate Member of Naknek Formation (loc. M 2076) to 530 feet above (loc. 25277).
21----	USGS Mesozoic loc. 29247. About 9 miles north of Katmai Bay at western base of Barrier Range near Fultons Falls and about 7.7 miles N. 54° E. of Topographers Peak, Mt. Katmai quadrangle, Alaska Peninsula. <i>Buchia mosquensis</i> (von Buch).
22----	USGS Mesozoic loc. M 5295. SE¼ sec. 32 (estimated), T. 42 S., R. 59 W., about 8.4 miles N. 30° W. of mouth of Thompson Creek on Chignik Bay, Chignik quadrangle, Alaska Peninsula. <i>Buchia rugosa</i> (Fischer) and <i>B. cf. B. mosquensis</i> (von Buch).
23----	USGS Mesozoic locs. M 1199, M 1244, M 1204 from Staniukovich Mountain between Herendeen Bay and Port Moller, Alaska Peninsula (Burk, 1965, p. 31, 160, 219). <i>Buchia mosquensis</i> (von Buch) and <i>B. rugosa</i> (Fischer).
24----	USGS Mesozoic loc. M 1234 from Lake Creek area southwest of Herendeen Bay, Alaska Peninsula (Burk, 1965, p. 31, 160, 219). <i>Buchia rugosa</i> (Fischer)?
25----	USGS Mesozoic loc. 29898. Black Hills area about 40 miles west of Herendeen Bay on north side of Alaska Peninsula. <i>Buchia rugosa</i> (Fischer).

Locality in figure 9	Description
Late middle to late Tithonian	
1----	USGS Mesozoic loc. 26390. Middle fork of Okpikruak River, lat 68°32'30" N., long 153°30'30" W., north-central Alaska. <i>Buchia</i> cf. <i>B. fischeriana</i> (d'Orbigny).
2----	USGS Mesozoic loc. M2531. Kemik Creek, 3.6 miles upstream from junction with Shavirovik River, Sagavanirtoq quadrangle, lat 69°27' N., long 147°08' W., northeast Alaska. <i>Buchia unschensis</i> (Pavlov).
3----	USGS Mesozoic locs. 30078 and 30079. On Joe Creek, 10.6–11 miles west of Canadian border and 300 feet or less higher stratigraphically than Mesozoic loc. 30076, which has furnished <i>Buchia rugosa</i> (Fischer). Table Mountains quadrangle, northeast Alaska. <i>Buchia unschensis</i> (Pavlov).
4----	USGS Mesozoic loc. 18349. Jacksina River, 1 mile southwest of mouth, just south of Nebesna, eastern part of Alaska Range, eastern Alaska (Moffit, 1943, pl. 2; 1954, p. 131). <i>Buchia</i> cf. <i>B. fischeriana</i> (d'Orbigny).
5----	USGS Mesozoic loc. 8811, near mouth of Gold Run Creek, a southern tributary of Glacier Creek (Capps, 1916, p. 52, fig. 9 on p. 100); Mesozoic loc. M5377, south side of Bonanza Creek, 5.4 miles N. 57° E. of junction with Chathenda Creek in SW¼ sec. 15, T. 4 N., R. 20 E.; Mesozoic loc. M5386, upper part of Chathenda Creek, 6.5 miles N. 75° E. of outlet of Little Beaver Lake at elevation of 5,300 ft.; Mesozoic loc. M5425, Coarse Money Creek, 1 mile upstream from its junction with Bonanza Creek, SW¼ sec. 17, T. 4 N., R. 20 E.; Mesozoic loc. M5563, south side of Chathenda Creek, 5.2 miles N. 46° E. of outlet of Little Beaver Lake, SE¼ sec. 22, T. 4 N., R. 20 E.; Mesozoic loc. M5564, Chathenda Creek, 5.3 miles N. 46° E. of outlet of Little Beaver Lake. All localities in central and west-central part of the Nabesna A-2 quadrangle in eastern part of Alaska Range. <i>Buchia fischeriana</i> (d'Orbigny).
6----	USGS Mesozoic loc. M1215. Northeast Creek area north of Amber Bay, Alaska Peninsula (Burk, 1965, p. 160, 178, 220). <i>Buchia piochii</i> (Gabb).
7----	USGS Mesozoic loc. M 5297. Ten miles N. 76° W. of mouth of Thompson Creek on Chignik Bay, north-central part of sec. 12 near sec. 1 (estimated), T. 43 S., R. 60 W., Chignik quadrangle, Alaska Peninsula (Burk, 1965, p. 160, 176, 219). <i>Buchia piochii</i> (Gabb).
8----	USGS Mesozoic loc. M 1211. Granville Portage area near Ivanoff Bay, Alaska Peninsula (Burk, 1965, p. 160, 170, 219). <i>Buchia piochii</i> (Gabb).
9----	USGS Mesozoic loc. M 1238. Staniukovich Mountain between Herendeen Bay and Port Moller, Alaska Peninsula (Burk, 1965, p. 160, 166, 219). <i>Buchia piochii</i> (Gabb).
10----	USGS Mesozoic loc. M 1236. Lake Creek area west of Herendeen Bay, Alaska Peninsula (Burk, 1965, p. 160, 162, 219). <i>Buchia piochii</i> (Gabb).

REFERENCES

- Arkell, W. J., 1956, *Jurassic geology of the world*: London, Oliver & Boyd, 806 p., 46 pls., 102 figs.
- Atwood, W. W., 1911, *Geology and mineral resources of parts of the Alaska Peninsula*: U.S. Geol. Survey Bull. 467, 137 p.
- Bergquist, H. R., 1966, *Micropaleontology of the Mesozoic rocks of northern Alaska*: U.S. Geol. Survey Prof. Paper 302-D, p. 93-227.
- Bishop, W. F., 1967, Age of the pre-Smackover formations, north Louisiana and south Arkansas: *Am. Assoc. Petroleum Geologists Bull.*, v. 51, no. 2, p. 244-250, 18 figs.
- Brabb, E. E., 1969, Six new Paleozoic and Mesozoic formations in east-central Alaska: U.S. Geol. Survey Bull. 1274-I, p. 1-26.
- 1970, Preliminary geologic map of the Black River quadrangle, east-central Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-601.
- Brabb, E. E., and Churkin, Michael, Jr., 1969, Geologic map of the Charley River quadrangle, east-central Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-573.
- Brew, D. A., Loney, R. A., and Muffler, L. J. P., 1966, Tectonic history of southeastern Alaska: *Canadian Inst. Mining and Metallurgy Spec. Vol. 8*, p. 149-170.
- Brosigé, W. P., and Reiser, H. N., 1964, Geologic map and section of the Chandalar quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-375.
- 1969, Preliminary geologic map of the Coleen quadrangle, Alaska: U.S. Geol. Survey open-file report, scale 1:250,000.
- Brosigé, W. P., and Tailleux, I. L., 1970, Depositional history of northern Alaska, in Adkison, W. L., and Brosigé, M. M., eds., *Geological seminar on the North Slope of Alaska*, Palo Alto, Calif., 1970, Proc.: Los Angeles, Am. Assoc. Petroleum Geologists, Pacific Sec., p. D1-D18.
- Burckhardt, Carlos, 1927, Cefalópodos del Jurásico medio de Oaxaca y Guerrero: *Mexico Inst. Geol. Bol.* 47, 108 p., 34 pls.
- 1930-31, *Étude synthétique sur le Mésozoïque mexicain*: *Schweizer. Paleont. Gesell. Abh.*, v. 49, pl. 1-123; v. 50, p. 123-280, 65 figs.
- Burk, C. A., 1965, Geology of the Alaska Peninsula—Island arc and continental margin: *Geol. Soc. America Mem.* 99, 250 p., 8 pls., 28 figs., 1 geol. map, 1 tectonic map.
- Cady, W. M., Wallace, R. E., Hoare, J. M., and Webber, E. J., 1955, The central Kuskokwim region, Alaska: U.S. Geol. Survey Prof. Paper 268, 132 p., 9 pls., 38 figs.
- Callomon, J. H., 1959, The ammonite zones of the Middle Jurassic beds of East Greenland: *Geol. Mag.*, v. 96, no. 6, p. 505-513, pls. 17, 18.
- Campbell, R. H., 1967, Areal geology in the vicinity of the Chariot site, Lisbourne Peninsula, northwestern Alaska: U.S. Geol. Survey Prof. Paper 395, 71 p.
- Capps, S. R., 1916, The Chisana-White River district, Alaska: U.S. Geol. Survey Bull. 630, 130 p., maps.
- 1923, The Cold Bay district [Alaska]: U.S. Geol. Survey Bull. 739-C, p. 77-116.
- 1934, Notes on the geology of the Alaska Peninsula and Aleutian Islands: U.S. Geol. Survey Bull. 857-D, p. 141-153, 1 pl.
- Carey, S. W., 1958, The tectonic approach to continental drift, in Carey, S. W., *Continental drift; a symposium*: Hobart, Tasmania, Tasmania Univ. Geology Dept., p. 177-355.
- Chapman, R. M., Detterman, R. L., and Mangus, M. D., 1964, *Geology of the Killik-Etiviluk Rivers region, Alaska*: U.S. Geol. Survey Prof. Paper 303-F, p. 325-407.
- Chapman, R. M., and Sable, E. G., 1960, *Geology of the Utukok-Corwin region, northwestern Alaska*: U.S. Geol. Survey Prof. Paper 303-C, p. 47-167.
- Collins, F. R., 1958, Test wells, Topagoruk area, Alaska: U.S. Geol. Survey Prof. Paper 305-D, p. 265-316, pls. 17-18, figs. 14-17.
- 1961, Core tests and test wells Barrow area, Alaska: U.S. Geol. Survey Prof. Paper 305-K, p. 569-643.
- Detterman, R. L., 1963, Revised stratigraphic nomenclature and age of the Tuxedni Group in the Cook Inlet region, Alaska: U.S. Geol. Survey Prof. Paper 475-C, p. C30-C34.
- 1973, Mesozoic sequence in Arctic Alaska, in *International Arctic Symposium*, 2d, Proc.: Am. Assoc. Petroleum Geologists, in press.
- Detterman, R. L., and Hartsock, J. K., 1966, *Geology of the Iniskin-Tuxedni region, Alaska*: U.S. Geol. Survey Prof. Paper 512, 78 p., 6 pls., 7 figs.
- Detterman, R. L., and Reed, B. L., 1964, Preliminary map of the geology of the Iliamna quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-407.
- Detterman, R. L., Reed, B. L., and Lanphere, M. A., 1965, Jurassic plutonism in the Cook Inlet region, Alaska: U.S. Geol. Survey Prof. Paper 525-D, p. D16-D21.
- Erben, H. K., 1956, *El Jurásico Inferior de México y sus amonitas*: Mexico City, Internat. Geol. Cong., 20th, 1956, 393 p., 41 pls., 5 maps.
- 1957, New biostratigraphic correlations in the Jurassic of eastern and south-central Mexico, in *El Mesozoico del Hemisferio Occidental y sus correlaciones mundiales*: Internat. Geol. Cong., 20th, Mexico City, 1956, [Trabajos], sec. 2, p. 43-52.
- Frebold, Hans, 1961, The Jurassic faunas of the Canadian Arctic—Middle and Upper Jurassic ammonites: *Canada Geol. Survey Bull.* 74, 43 p., 21 pls., 3 figs.
- 1964a, Illustrations of Canadian Jurassic fossils—Jurassic of western and Arctic Canada: *Canada Geol. Survey Paper* 63-4, 51 pls.
- 1964b, Outline of the Jurassic system in Canada: *Colloque du Jurassique, Luxembourg 1962, Comptes Rendus et Mém.*, p. 479-483.
- Frebold, Hans, Mountjoy, E. W., and Reed, R. A., 1959, The Oxfordian beds of the Jurassic Fernie group, Alberta and British Columbia: *Canada Geol. Survey Bull.* 53, 47 p., 12 pls., 6 figs.
- Frebold, Hans, Mountjoy, E. W., and Tempelman-Kluit, D. J., 1967, New occurrences of Jurassic rocks and fossils in central and northern Yukon Territory: *Canada Geol. Survey Paper* 67-12, 35 p., 3 pls., 2 figs.
- Frebold, Hans, and Tipper, H. W., 1967, Middle Callovian sedimentary rocks and guide ammonites from southwestern British Columbia: *Canada Geol. Survey Paper* 67-21, 29 p., 3 pls., 2 text-figs.
- 1970, Status of the Jurassic in the Canadian Cordillera of British Columbia, Alberta, and southern Yukon: *Canadian Jour. Earth Sci.*, v. 7, no. 1, p. 1-21, 9 figs.
- Gates, G. O., Grantz, Arthur, and Patton, W. W., Jr., 1968, *Geology and natural gas and oil resources of Alaska—Part 1, Natural gases in rocks of Cenozoic age*: Am. Assoc. Petroleum Geologists Mem. 9, v. 1, p. 3-48.

- Gates, G. O., and Gryc, George, 1963, Structure and tectonic history of Alaska, in *Backbone of the Americas*: Am. Assoc. Petroleum Geologists Mem. 2, p. 264-277.
- Grantz, Arthur, 1960a, Geologic map of Talkeetna Mountains (A-2) quadrangle, Alaska, and the contiguous area to north and northwest: U.S. Geol. Survey Misc. Geol. Inv. Map I-313.
- 1960b, Geologic map of Talkeetna Mountains (A-1) quadrangle, and the south third of Talkeetna Mountains (B-1) quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-314.
- 1961a, Geologic map and cross sections of the Anchorage (D-2) quadrangle and northeasternmost part of the Anchorage (D-3) quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-342.
- 1961b, Geologic map of the north two-thirds of the Anchorage (D-1) quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-343.
- 1966, Strike-slip faults in Alaska: U.S. Geol. Survey open-file report, 82 p., 8 figs.
- Grantz, Arthur, Jones, D. L., and Lanphere, M. A., 1966, Stratigraphy, paleontology, and isotopic ages of upper Mesozoic rocks in the southwestern Wrangell Mountains, Alaska: U.S. Geol. Survey Prof. Paper 550-C, p. C39-C47.
- Grantz, Arthur, Thomas, Herman, Stern, T. W., and Sheffey, N. B., 1963, Potassium-argon and lead-alpha ages for stratigraphically bracketed plutonic rocks in the Talkeetna Mountains, Alaska: U.S. Geol. Survey Prof. Paper 475-B, p. B56-B59.
- Gryc, George, 1951, Paleontology, in Payne, T. G., and others, *Geology of the Arctic slope of Alaska*: U.S. Geol. Survey Oil and Gas Inv., Map OM-126, scale 1:1,000,000, 3 sheets, text.
- Gryc, George, and Lathram, E. H., 1954, Preliminary report on the geology of the Sagavanirktok River area, 1946; U.S. Geol. Survey open-file report, 5 p.
- Gryc, George, and Mangus, M. D., 1954, Preliminary report on the stratigraphy and structure of the area of the Shaviovik and Canning Rivers, Alaska, 1947: U.S. Geol. Survey open-file report, 7 p., 1 pl.
- Hallam, Anthony, 1965, Observations on marine Lower Jurassic stratigraphy of North America, with special reference to the United States: Am. Assoc. Petroleum Geologists Bull., v. 49, no. 9, 1485-1501, 5 figs.
- Hamilton, W. B., 1969, Continental drift in the Arctic [abs.]: *Geol. Soc. America Spec. Paper* 121, p. 510.
- Hoare, J. M., 1961, Geology and tectonic setting of lower Kuskokwim-Bristol Bay region, Alaska: Am. Assoc. Petroleum Geologists Bull., v. 45, no. 5, p. 594-611.
- Hoare, J. M., and Coonrad, W. L., 1959a, Geology of the Bethel quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-285.
- 1959a, Geology of the Russian Mission quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-292.
- 1961a, Geologic map of the Hagemeister Island quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-321.
- 1961b, Geologic map of the Goodnews quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-339.
- Howarth, M. K., 1958, A monograph of the ammonites of the Liassic family Amaltheidae in Britain: London, Palaeontograph. Soc., pt. 1, p. i-xiv, 1-26, pls. 1-4 (v. 111); pt. 2, p. xv-xxxvii, 27-53, pls. 5-10 (v. 112).
- Imlay, R. W., 1943, Jurassic formations of Gulf region: Am. Assoc. Petroleum Geologists Bull., v. 27, no. 11, p. 1407-1533, 14 figs.
- 1952, Correlation of the Jurassic formations of North America, exclusive of Canada: *Geol. Soc. America Bull.*, v. 63, no. 9, p. 953-992, 2 correlation charts.
- 1953a, Callovian (Jurassic) ammonites from the United States and Alaska—Part 1. Western Interior United States: U.S. Geol. Survey Prof. Paper 249-A, p. 1-40, pls. 1-24, figs. 1, 2.
- 1953b, Callovian (Jurassic) ammonites from the United States and Alaska—Part 2. Alaska Peninsula and Cook Inlet region: U.S. Geol. Survey Prof. Paper 249-B, p. 41-108, pls. 25-55, figs. 3-9.
- 1955, Characteristic Jurassic mollusks from northern Alaska: U.S. Geol. Survey Prof. Paper 274-D, p. 69-96, pls. 8-13, fig. 20.
- 1959, Succession and speciation of the pelecypod *Aucella*: U.S. Geol. Survey Prof. Paper 314-G, p. 155-169.
- 1961a, New genera and subgenera of Jurassic (Bajocian) ammonites from Alaska: *Jour. Paleontology*, v. 35, no. 3, p. 467-474.
- 1961b, Late Jurassic ammonites from the western Sierra Nevada, California: U.S. Geol. Survey Prof. Paper 374-D, p. 1-30, 6 pls., 4 figs.
- 1962a, Jurassic (Bathonian or early Callovian) ammonites from Alaska and Montana: U.S. Geol. Survey Prof. Paper 374-C, 32 p.
- 1962b, Late Bajocian ammonites from the Cook Inlet region, Alaska: U.S. Geol. Survey Prof. Paper 418-A, p. 1-15, pls. 1-5, figs. 1-4.
- 1964a, Upper Jurassic mollusks from eastern Oregon and western Idaho: U.S. Geol. Survey Prof. Paper 483-D, p. 1-21, 4 pls., 3 figs.
- 1964b, Middle Bajocian ammonites from the Cook Inlet region, Alaska: U.S. Geol. Survey Prof. Paper 418-B, p. 1-61, pls. 1-29, 5 figs.
- 1965, Jurassic marine faunal differentiation in North America: *Jour. Paleontology*, v. 39, no. 5, p. 1023-1038, 6 figs.
- 1967, The Mesozoic pelecypods *Otapiria* Marwick and *Lupherella* Imlay, new genus, in the United States: U.S. Geol. Survey Prof. Paper 573-B, p. 1-11, 2 pls., 2 figs.
- 1968, Lower Jurassic (Pliensbachian and Toarcian) ammonites from eastern Oregon and California: U.S. Geol. Survey Prof. Paper 593-C, p. 1-51, 9 pls., 8 figs.
- 1970, Jurassic paleogeography of Alaska [abs.]: Am. Assoc. Petroleum Geologists Bull., v. 54, no. 12, p. 2487.
- Imlay, R. W., and Jones, D. L., 1970, Ammonites from the *Buchia* zones in northwestern California and southwestern Oregon: U.S. Geol. Survey Prof. Paper 647-B, p. B1-B59, 15 pls., 8 figs.
- Imlay, R. W., and Reeside, J. B., Jr., 1954, Correlation of the Cretaceous formations of Greenland and Alaska: *Geol. Soc. America Bull.*, v. 65, no. 3, p. 223-246.
- Jeletzky, J. A., 1958, Uppermost Jurassic and Cretaceous rocks of Aklavik Range, northeastern Richardson Mountains, Northwest Territories: Canada Geol. Survey Paper 58-2, 84 p., geol. sketch map, correlation chart.
- 1960, Uppermost Jurassic and Cretaceous rocks, east flank of Richardson Mountains between Stony Creek and lower Donna River, Northwest Territories, 106 M and 107B (parts of): Canada Geol. Survey Paper 59-14 31 p., geol. sketch maps, correlation chart.

- 1962, Pre-Cretaceous Richardson Mountains trough—Its place in the tectonic framework of Arctic Canada and its bearing on some geosynclinal concepts: *Royal Soc. Canada Trans.*, 3d ser., v. 56, sec. 3, pt. 1, p. 55-84.
- 1966, Upper Volgian (latest Jurassic) ammonites and Buchias of Arctic Canada: *Canada Geol. Survey Bull.* 128, 51 p., 8 pls., 2 figs.
- 1967, Jurassic and (?) Triassic rocks of the eastern slope of Richardson Mountains, northwestern District of Mackenzie: *Canada Geol. Survey Paper* 66-50, 171 p., 9 pls., 3 figs.
- Jones, D. L., and Grantz, Arthur, 1964, Stratigraphic and structural significance of Cretaceous fossils from Tiglukpuk Formation, northern Alaska: *Am. Assoc. Petroleum Geologists Bull.*, v. 48, no. 9, p. 1462-1474, 3 pls., 4 figs.
- Juhle, R. W., 1955, Iliamna Volcano and its basement: *U.S. Geol. Survey open-file report*, 74 p., 38 pls.
- Keller, A. S., Morris, R. H., and Detterman, R. L., 1961, Geology of the Shaviovik and Saganvanirktok Rivers region, Alaska: *U.S. Geol. Survey Prof. Paper* 303-D, p. 169-222, pls. 21-26, figs. 26-32.
- Keller, A. S., and Reiser, H. N., 1959, Geology of the Mount Katmai area, Alaska: *U.S. Geol. Survey Bull.* 1058-G, p. 261-298, illus.
- Kellum, L. B., 1945, Jurassic stratigraphy of Alaska and petroleum exploration in northwest America: *New York Acad. Sci. Trans.*, ser. 2, v. 7, no. 8, p. 201-209.
- Kirschner, C. E., and Lyon, C. A., 1973, Stratigraphy and tectonic development of Cook Inlet petroleum province, in *International Arctic Symposium*, 2d, Proc.: *Am. Assoc. Petroleum Geologists*, in press.
- Kirschner, C. E., and Minard, D. L., 1949, Geology of the Iniskin Peninsula, Alaska: *U.S. Geol. Survey Oil and Gas Inv. Prelim. Map* OM-95.
- Knowlton, F. H., 1914, The Jurassic flora of Cape Lisburne, Alaska: *U.S. Geol. Survey Prof. Paper* 85-D, p. 39-64, illus.
- Lathram, E. H., 1965, Preliminary geologic map of northern Alaska: *U.S. Geol. Survey open-file map*, 2 sheets, scale 1:1,000,000.
- Lathram, E. H., Loney, R. A., Berg, H. C., and Pomeroy, J. S., 1960, Progress map of the geology of Admiralty Island, Alaska: *U.S. Geol. Survey Misc. Geol. Inv. Map* I-323.
- Leffingwell, E. de K., 1919, The Canning River region, northern Alaska: *U.S. Geol. Survey Prof. Paper* 109, 251 p.
- Loeblich, H. N., 1954, Summary of microfossil investigations, April 1948 to April 1949, with special reference to the Barrow and Simpson area, 1949: *U.S. Geol. Survey open-file report*, 83 p., 2 pls., 1 fig.
- Loney, R. A., 1964, Stratigraphy and petrography of the Pybus-Gambier area, Admiralty Island, Alaska: *U.S. Geol. Survey Bull.* 1178, 103 p., 2 pls., 33 figs.
- MacKevett, E. M., Jr., 1963, Preliminary geologic map of the McCarthy C-5 quadrangle, Alaska: *U.S. Geol. Survey Misc. Geol. Inv. Map* I-406.
- 1965a, Preliminary geologic map of the McCarthy B-5 quadrangle, Alaska: *U.S. Geol. Survey Misc. Geol. Inv. Map* I-438.
- 1965b, Preliminary geologic map of the McCarthy C-6 quadrangle, Alaska: *U.S. Geol. Survey Misc. Geol. Inv. Map* I-444.
- 1969, Three newly named Jurassic formations in the McCarthy C-5 quadrangle, Alaska: *U.S. Geol. Survey Bull.* 1274-A, p. 35-49, figs. 3-8.
- 1970a, Geologic map of the McCarthy C-4 quadrangle, Alaska: *U.S. Geol. Survey Geol. Quad. Map* GQ-844.
- 1970b, Geologic map of the McCarthy C-5 quadrangle, Alaska: *U.S. Geol. Survey Geol. Quad. Map* GQ-899.
- 1971, Stratigraphy and general geology of the McCarthy C-5 quadrangle, Alaska: *U.S. Geol. Survey Bull.* 1323, 35 p.
- Maddren, A. G., 1912, Geologic investigations along the Canada-Alaska boundary: *U.S. Geol. Survey Bull.* 520, p. 297-314.
- Maddren, A. G., and Harrington, G. L., 1914, [Geology on] Topographic maps 141st Meridian, Porcupine River to Arctic Coast: Washington, D.C., *Internat. Boundary Comm.*
- Mangus, M. D., Detterman, R. L., Lachenbruch, M. C., Jr., Lachenbruch, A. H., 1954, Stratigraphy and structure of the Etivluk and Kuna Rivers area, Alaska, 1950: *U.S. Geol. Survey open-file report*, 15 p., 3 pls., 1 fig.
- Martin, G. C., 1905, The petroleum fields of the Pacific coast of Alaska, with an account of the Bering River coal deposits: *U.S. Geol. Survey Bull.* 250, 64 p.
- 1921, Preliminary report on petroleum in Alaska: *U.S. Geol. Survey Bull.* 719, 83 p., 11 pls., 6 figs.
- 1925, The outlook for petroleum near Chignik [Alaska]: *U.S. Geol. Survey Bull.* 773-D, p. 209-213.
- 1926, The Mesozoic stratigraphy of Alaska: *U.S. Geol. Survey Bull.* 776, 493 p., 13 figs.
- Martin, G. C., and Katz, F. J., 1912, A geologic reconnaissance of the Iliamna region, Alaska: *U.S. Geol. Survey Bull.* 485, 138 p.
- Mather, K. F., 1925, Mineral resources of the Kamishak Bay region [Alaska]: *U.S. Geol. Survey Bull.* 773-D, p. 159-181, illus.
- Mertie, J. B., 1929, Preliminary report on the Sheenjek River district [Alaska]: *U.S. Geol. Survey Bull.* 797, p. 99-123.
- 1930, The Chandalar-Sheenjek district [Alaska]: *U.S. Geol. Survey Bull.* 810-B, p. 87-139, 2 pls., 2 figs.
- Miller, D. J., Payne, T. G., and Gryc, George, 1959, Geology of possible petroleum provinces in Alaska: *U.S. Geol. Survey Bull.* 1094, 131 p.
- Moffit, F. H., 1922a, Geology of the vicinity of Tuxedni Bay, Cook Inlet, Alaska: *U.S. Geol. Survey Bull.* 722-D, p. 141-147, illus.
- 1922b, The Iniskin Bay district [Alaska]: *U.S. Geol. Survey Bull.* 739-C, p. 117-132, 1 pl.
- 1927, The Iniskin-Chinitna Peninsula and the Snug Harbor district, Alaska: *U.S. Geol. Survey Bull.* 789, 71 p., 11 pls., 1 fig.
- 1938a, Geology of the Chitina Valley and adjacent area, Alaska: *U.S. Geol. Survey Bull.* 894, 137 p., 16 pls., 1 fig.
- 1938b, Geology of the Slana-Tok district, Alaska: *U.S. Geol. Survey Bull.* 904, 54 p., 4 pls., 4 figs.
- 1943, Geology of the Nutzotin Mountains, Alaska: *U.S. Geol. Survey Bull.* 933-B, p. 103-299, pls. 2-12, text-figs. 4-7.
- 1954, Geology of the eastern part of the Alaska Range and adjacent area: *U.S. Geol. Survey Bull.* 989-D, p. 65-218, pls. 6, 7, figs. 18-40.

- Moffit, F. H., and Knopf, Adolph, 1910, Mineral resources of the Nabesna-White River district, Alaska; with a section on the Quaternary, by S. R. Capps: U.S. Geol. Survey Bull. 417, 64 p.
- Muller, S. W., and Ferguson, H. G., 1939, Mesozoic stratigraphy of the Hawthorne and Tonopah quadrangles, Nevada: Geol. Soc. America Bull., v. 50, no. 10, p. 1573-1624, 6 pls., 4 figs.
- Paige, Sidney, 1906, The Herendeen Bay coal field: U.S. Geol. Survey Bull. 284, p. 101-108.
- Paige, Sidney, and Knopf, Adolph, 1907, Geologic reconnaissance in the Matanuska and Talkeetna basins, Alaska: U.S. Geol. Survey Bull. 327, 71 p.
- Patton, W. W., Jr., 1956, New formation of Jurassic age, in Gryc, George, and others, Mesozoic sequence in Colville River region, northern Alaska: Am. Assoc. Petroleum Geologists Bull., v. 40, no. 2, p. 213-218.
- Patton, W. W., Jr., and Grantz, Arthur, 1962, Tectonic implications of some new Mesozoic stratigraphic data on Alaska [abs.]: Am. Assoc. Petroleum Geologists Bull., v. 46, no. 2, p. 274-275.
- Patton, W. W., Jr., and Keller, A. S., 1954, Stratigraphy and structure of the upper Siksikpuk-Nanushuk Rivers area, Alaska, 1951: U.S. Geol. Survey open-file report, 20 p., 4 pls., 3 figs.
- Patton, W. W., Jr., and Tailleur, I. L., 1954, Stratigraphy and structure of the Okpikruak and Kiruktagiak Rivers area, Alaska, 1950: U.S. Geol. Survey open-file report, 17 p., 3 pls., 1 fig.
- 1964, Geology of the Killik-Itkillik region, Alaska: U.S. Geol. Survey Prof. Paper 303-G, p. 409-500, pls. 50-51, figs. 66-91.
- Payne, T. G., 1955, (Compiler), Mesozoic and Cenozoic tectonic elements of Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-84.
- Payne, T. G., and others, 1951, Geology of the Arctic slope of Alaska: U.S. Geol. Survey Oil and Gas Inv. Map OM-126, scale 1:1,000,000, 3 sheets, text.
- Reed, B. L., 1968, Geology of the Lake Peters area, northeastern Brooks Range, Alaska: U.S. Geol. Survey Bull. 1236, 132 p.
- Reed, B. L., and Lanphere, M. A., 1969, Age and chemistry of Mesozoic and Tertiary plutonic rocks in south-central Alaska: Geol. Soc. America Bull., v. 80, no. 1, p. 23-44.
- Reiser, H. N., Lanphere, M. A., and Brosgé, W. P., 1965, Jurassic age of mafic igneous complex, Christian quadrangle, Alaska: U.S. Geol. Survey Prof. Paper 525-C, p. C68-C71.
- Reiser, H. N., and Tailleur, I. L., 1969, compilers, Preliminary geologic map of Mt. Michelson quadrangle, Alaska: U.S. Geol. Survey open-file map, scale 1:200,000.
- Reiser, H. N., and others, 1970, Progress map, geology of the Sadlerochit and Shublik Mountains, Mt. Michelson C-1, C-2, C-3, and C-4 quadrangles, Alaska: U.S. Geol. Survey open-file report, 5 sheets (4 maps and explanation).
- Richter, D. H., 1971, Reconnaissance geologic map and section of the Nabesna B-4 quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-656.
- Richter, D. H., and Jones, D. L., 1970, Structure and stratigraphy of the Eastern Alaska Range, Alaska [abs.]: Am. Assoc. Petroleum Geologists Bull., v. 54, no. 12, p. 2502.
- Rickwood, F. K., 1970, The Prudhoe Bay field, in Adkison, W. L., and Brosgé, M. M., eds., Geological seminar on the North Slope of Alaska, Palo Alto, Calif., 1970, Proc.: Los Angeles, Am. Assoc. Petroleum Geologists, Pacific Sec., p. L1-L11.
- Robinson, F. M., 1959, Test wells, Simpson area, Alaska: U.S. Geol. Survey Prof. Paper 305-J, p. 523-568.
- Sable, E. G., 1965, Geology of the Romanzof Mountains, Brooks Range, northeastern Alaska: U.S. Geol. Survey open-file report, 218 p., 7 pls., 23 figs.
- Sable, E. G., and Mangus, M. D., 1954, Stratigraphy and structure of the upper Utukok-Kokolik Rivers area, Alaska, 1951: U.S. Geol. Survey open-file report, 19 p., 2 pls.
- St. Amand, Pierre, 1957, Geological and geophysical synthesis of the tectonics of portions of British Columbia, the Yukon Territory, and Alaska: Geol. Soc. America Bull., v. 68, no. 10, p. 1343-1370.
- Smith, P. S., 1917, The Lake Clark-central Kuskokwim region, Alaska: U.S. Geol. Survey Bull. 655, 162 p.
- 1939, Areal geology of Alaska: U.S. Geol. Survey Prof. Paper 192, 100 p.
- Smith, P. S., and Mertie, J. B., Jr., 1930, Geology and mineral resources of northwestern Alaska: U.S. Geol. Survey Bull. 815, 351 p., 34 pls., 22 figs.
- Smith, W. R., 1925, The Cold Bay-Katmai district [Alaska]: U.S. Geol. Survey Bull. 773-D, p. 183-207, map.
- Smith, W. R., and Baker, A. A., 1924, The Cold Bay-Chignik district, Alaska: U.S. Geol. Survey Bull. 755-D, p. 151-218, illus.
- Spurr, J. E., 1900, A reconnaissance in southwestern Alaska in 1898: U.S. Geol. Survey Ann. Rept. 20, pt. 7, p. 31-264, illus.
- Stanton, T. W., and Martin, G. C., 1905, Mesozoic section on Cook Inlet and Alaska Peninsula: Geol. Soc. America Bull., v. 16, p. 391-410.
- Stone, R. W., 1965, Coal resources of southwestern Alaska: U.S. Geol. Survey Bull. 259, p. 151-171, illus.
- Tailleur, I. L., 1964, Rich oil shale from northern Alaska: U.S. Geol. Survey Prof. Paper 475-D, p. D131-D133.
- 1969, Speculations on North Slope geology: Oil and Gas Jour., v. 67, no. 38, p. 215-220, 225-226; no. 39, p. 128-130.
- Tailleur, I. L., and Brosgé, W. P., 1970, Tectonic history of northern Alaska, in Adkison, W. L., and Brosgé, M. M., eds., Geological seminar on the North Slope of Alaska, Palo Alto, Calif., 1970, Proc.: Los Angeles, Am. Assoc. Petroleum Geologists, Pacific Sec., p. E1-E19.
- Tailleur, I. L., and Kent, B. H., 1954, Stratigraphy and structure of the Southern Foothills section between the Etivluk and Kiligwa Rivers, Alaska, 1951: U.S. Geol. Survey open-file report, 26 p., 4 pls., 2 figs.
- Tailleur, I. L., Kent, B. H., Jr., and Reiser, H. N., 1966, Outcrop geologic maps of the Nuka-Etivluk region, northern Alaska: U.S. Geol. Survey open-file report, 7 sheets (map, scale 1:63,360, and explanation).
- Tappan, Helen, 1951, Micropaleontology, in Payne, T. G., and others, Geology of the Arctic slope of Alaska: U.S. Geol. Survey Oil and Gas Inv. Map OM-126, scale 1:1,000,000, 3 sheets, text.
- 1955, Foraminifera from the Arctic slope of Alaska—Pt. 2, Jurassic Foraminifera: U.S. Geol. Survey Prof. Paper 236-B, p. 21-90, pls. 6-28, figs. 3-9.
- Viniegua O., Francisco, 1971, Age and evolution of salt basins of southeastern Mexico: Am. Assoc. Petroleum Geologists Bull., v. 55, no. 3, p. 478-494, 10 figs.

- Wahrhaftig, Clyde, 1965, Physiographic divisions of Alaska: U.S. Geol. Survey Prof. Paper 482, 52 p.
- Westermann, G. E. G., 1964, The ammonite fauna of the Kialagvik Formation at Wide Bay, Alaska Peninsula. Part 1, Lower Bajocian (Aalenian): *Bulls. Am. Paleontology*, v. 47, no. 216, p. 327-503, pls. 44-76, 37 figs.
- 1969, The ammonite fauna of the Kialagvik Formation at Wide Bay, Alaska Peninsula—Part II, *Sonninia sowerbyi* zone (Bajocian): *Bulls. Am. Paleontology*, v. 57, no. 255, 226 p., pls. 1-47, 56 figs.
- Whittington, C. L., and Sable, E. G., 1954, Preliminary geologic report of Sadlerochit River area, 1948: U.S. Geol. Survey open-file report, 18 p., 1 pl.

INDEX

[Italic page numbers indicate major references]

	Page
Acknowledgments	2
Admiralty Island	20
Alaska Peninsula 2, 8, 9, 14, 15, 17, 18, 20	
Alaska Range	9, 20
Aleutian Range batholith	16
Alinchak Bay	14
Ammonite differentiation, Boreal, Pacific, Tethyan realms ..	20
Ammonites, Alaskan Jurassic, com- pared with those of other parts of world	20
<i>Amoeboceras</i>	19, 21
Arctic Ocean	20
<i>Arcticoceras</i>	20, 21
<i>Arkelloceras</i>	16, 17
<i>Artocephalites</i>	20, 21
Bajocian fossil occurrences	23
Bajocian seas, extent in Alaska	17
Bathonian fossil occurrences	24
Bathonian seas, extent in Alaska	18
Boreal realm	20, 21
Bowser Formation	14
British Columbia	17, 18, 20, 21
Brooks Range	16
<i>Buchia concentrica</i>	19
<i>fischeri</i>	20
<i>fischeriana</i>	15
<i>mosquensis</i>	16, 21
<i>piochii</i>	15, 21
<i>rugosa</i>	15, 21
Early Cretaceous species	16
<i>Cadoceras</i>	16, 21
(<i>Paracadoceras</i>)	21
(<i>Stenocadoceras</i>)	21
California	20, 21
Callovian ammonites, late, absence in Alaska and Canada; presence in California and southern Mexico	16
Callovian fossil occurrences	24
Callovian seas, extent in Alaska	18
Callovian to Oxfordian transition, Southern Alaska	8
Canning River	16
Cape Lisburne	14
<i>Cardioceras</i>	15, 16, 19, 21
Chinitna Bay	14
Chinitna Formation	15
Christian	14
Coleen River	14
Concretions, limestone	9
orange-weathering clay ironstone ..	9
Cook Inlet	8, 15, 17, 18
<i>Cranocephalites</i>	16, 20, 21
Cretaceous rocks, marine	2
<i>Cruciloboceras</i>	16
Depositional environment, Early to Middle Jurassic time	8
<i>Dichotomosphinctes</i>	21
Disconformity, pre-late Oxfordian, Root Glacier Formation	19
Emergence, Callovian, early Oxfordian, late Tithonian	14

	Page
Epieugeosynclinal trough	8
Etivluk River	16
Foraminifera, north-central Alaska, Late Jurassic age questioned ..	16
Fossil occurrences by stages	21
Bajocian	23
Bathonian	24
Callovian	24
early Oxfordian	25
Hettangian	21
late Kimmeridgian to early middle Tithonian	26
late middle to late Tithonian	27
late Oxfordian to early Kimmeridgian	25
Pliensbachian	22
Sinemurian	22
Toarcian	23
Greenland	20, 21
Guerrero, Mexico	16
Gulf of Mexico, Jurassic salt and red beds	21
Hettangian fauna, Porcupine River	14
Hettangian fossil occurrences	21
Iniskin Peninsula	2, 14, 17
<i>Inoceramus ambiguus</i>	14, 17, 18
Introduction	1
Itkillik River	16
Juneau	8
Jurassic rocks, marine, absence in central Alaska	2
Kandik River	14
Katmai	8
<i>Kepplerites</i>	21
Ketchikan	8
Killik River	16
Kimmeridgian, late to early middle Tithonian fossil occurrences ..	26
Kimmeridgian seas, extent in Alaska ..	19
Kuskokwim	8, 17, 18, 20
Late Kimmeridgian to middle Tithonian seas, extent in Alaska	20
Laumontite	8
<i>Leioceras opalinum</i> zone	14
<i>Lilloettia</i>	21
Lithologic, stratigraphic features compared	8
Lubbe Creek Formation	14
Lupine River	16
<i>Lytoceras</i>	21
Marine connections between north and south Alaska in Jurassic time	9
Marine Cretaceous rocks	2
Marine Jurassic rocks, absence in central Alaska	2
Marine Triassic rocks	2
<i>Megasphaeroceras rotundum</i>	17

	Page
Mount Peulik	15
Naknek Formation	15
Nation River	14
Nizina Mountain Formation	14
Oaxaca, Mexico	16
Oil shale	9
Okpikruak River	17
<i>Otapiria tailleuri</i>	16, 17
<i>Otoites sauzei</i> zone	14
Oxfordian, early, fossil occurrences ..	25
late, to early Kimmeridgian fossil occurrences	25
Oxfordian seas, extent in Alaska	19
Pacific realm	20
Paleobiogeographic resemblances, differences during Jurassic, northern and southern Alaska	16
Paleogeographic inferences, Alaska Jurassic seas	16
<i>Parabigotites crassicostratus</i>	17
<i>Parareineckia</i>	20
<i>Phylloceras</i>	21
Pliensbachian fossil occurrences	22
Pliensbachian seas, extent in Alaska ..	16
Plutonism	8
Porcupine River	14
Hettangian fauna	14
<i>Pseudocadoceras</i>	16, 21
Puale Bay	15, 18
Radiometric dating of igneous rocks ..	8
Red beds, Jurassic, deposition in Gulf of Mexico	21
Root Glacier Formation, pre-late Oxfordian disconformity	19
Sagavanirktok River	16, 17
Salt, Jurassic, deposition in Gulf of Mexico	21
Seas, Jurassic, extent in Alaska	16
Sheenjek River	14
Sinemurian fossil occurrences	22
Sinemurian seas, extent in Alaska	16
<i>Stephanoceras humphriesianum</i> zone ..	14, 16
Stratigraphic, lithologic features compared	8
Stratigraphy, Jurassic rocks, east- central Alaska	14
northern Alaska	9
southern Alaska	8
Talkeetna Mountains	2, 8, 14, 15, 20
<i>Teloceras itinsae</i>	17
Tethyan realm	20, 21
Tiglupuk Creek	14
Tiglupuk Formation, name abandoned ..	14
Tithonian, late middle to late, fossil occurrences	27

	Page		Page		Page
Tithonian seas, extent in Alaska	20	Unconformity, Bathonian-Callovian		Unconformity, Toarcian to early	
<i>Tmetoceras</i>	16	boundary, northern Alaska	16	Bajocian	8
Toarcian fossil occurrences	23	late Bathonian, southern Alaska ..	14	top Lower Jurassic, southern Alaska	14
Toarcian seas, extent in Alaska	16	late Callovian, northern Alaska ---	16	Volcanic island-arc system	8
Toarcian to early Bajocian unconformity	8	late Callovian, southern Alaska --	15	Wide Bay	8, 14, 15, 17, 18, 20
Triassic rocks, marine	2	late late Bajocian, southern Alaska	14	Wrangell Mountains ----	8, 9, 14, 15, 17, 18, 19
Tuxedni Bay	14	late middle Bajocian to early		<i>Xenoccephalites</i>	20, 21
Twist Creek Siltstone	14	Bathonian, northern Alaska	16	Yukon Territory	18, 20
Unconformities, Jurassic, occurrences..	14	late Tithonian, Berriasian,			
		southern Alaska	16		
		middle Bajocian to early Bathonian	9		

